


(51) International Patent Classification ⁶ : C12N 15/29, 15/62, 15/70, 15/86, A61K 38/16, 48/00	A2	(11) International Publication Number: WO 98/49311
		(43) International Publication Date: 5 November 1998 (05.11.98)

Without international search report and to be republished upon receipt of that report.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**Title: RICIN-LIKE TOXIN VARIANTS FOR TREATMENT OF
CANCER, VIRAL OR PARASITIC INFECTIONS**

FIELD OF THE INVENTION

The invention relates to proteins useful as therapeutics
5 against cancer, viral infections, parasitic and fungal infections. The
proteins contain A and B chains of a ricin-like toxin linked by a linker
sequence that is specifically cleaved and activated by proteases specific to
disease-associated pathogens or cells.

BACKGROUND OF THE INVENTION

10 Bacteria and plants are known to produce cytotoxic
proteins which may consist of one, two or several polypeptides or
subunits. Those proteins having a single subunit may be loosely
classified as Type I proteins. Many of the cytotoxins which have
evolved two subunit structures are referred to as type II proteins
15 (Saelinger, C.B. in Trafficking of Bacterial Toxins (eds. Saelinger, C.B.)
1-13 (CRC Press Inc., Boca Raton, Florida, 1990). One subunit, the A
chain, possesses the toxic activity whereas the second subunit, the B
chain, binds cell surfaces and mediates entry of the toxin into a target
cell. A subset of these toxins kill target cells by inhibiting protein
20 biosynthesis. For example, bacterial toxins such as diphtheria toxin or
Pseudomonas exotoxin inhibit protein synthesis by inactivating
elongation factor 2. Plant toxins such as ricin, abrin, and bacterial toxin
Shiga toxin, inhibit protein synthesis by directly inactivating the
ribosomes (Olsnes, S. & Phil, A. in Molecular action of toxins and
25 viruses (eds. Cohen, P. & vanHeyningen, S.) 51-105 Elsevier Biomedical
Press, Amsterdam, 1982).

Ricin, derived from the seeds of *Ricinus communis*
(castor oil plant), may be the most potent of the plant toxins. It is
estimated that a single ricin A chain is able to inactivate ribosomes at a

rate of 1500 ribosomes/minute. Consequently, a single molecule of ricin is enough to kill a cell (Olsnes, S. & Phil, A. in *Molecular action of toxins and viruses* (eds. Cohen, P. & vanHeyningen, S.) (Elsevier Biomedical Press, Amsterdam, 1982). The ricin toxin is a glycosylated
5 heterodimer consisting of A and B chains with molecular masses of 30,625 Da and 31,431 Da linked by a disulphide bond. The A chain of ricin has an N-glycosidase activity and catalyzes the excision of a specific adenine residue from the 28S rRNA of eukaryotic ribosomes (Endo, Y. & Tsurugi, K. J., *Biol. Chem.* 262:8128 (1987)). The B chain of ricin,
10 although not toxic in itself, promotes the toxicity of the A chain by binding to galactose residues on the surface of eukaryotic cells and stimulating receptor-mediated endocytosis of the toxin molecule (Simmons et al., *Biol. Chem.* 261:7912 (1986)). Once the toxin molecule consisting of the A and B chains is internalized into the cell via
15 clathrin-dependent or independent mechanisms, the greater reduction potential within the cell induces a release of the active A chain, eliciting its inhibitory effect on protein synthesis and its cytotoxicity (Emmanuel, F. et al., *Anal. Biochem.* 173: 134-141 (1988); Blum, J.S. et al., *J. Biol. Chem.* 266: 22091-22095 (1991); Fiani, M.L. et al., *Arch. Biochem. Biophys.* 307: 225-230 (1993)). Empirical evidence suggests that activated toxin (e.g. ricin, shiga toxin and others) in the endosomes is transcytosed through the trans-Golgi network to the endoplasmic reticulum by retrograde transport before the A chain is translocated into the cytoplasm to elicit its action (Sandvig, K. & van Deurs, B., *FEBS Lett.*
25 346: 99-102 (1994).

Protein toxins are initially produced in an inactive, precursor form. Ricin is initially produced as a single polypeptide (preproricin) with a 35 amino acid N-terminal presequence and 12 amino acid linker between the A and B chains. The pre-sequence is
30 removed during translocation of the ricin precursor into the endoplasmic reticulum (Lord, J.M., *Eur. J. Biochem.* 146:403-409 (1985) and Lord, J.M., *Eur. J. Biochem.* 146:411-416 (1985)). The proricin is then

translocated into specialized organelles called protein bodies where a plant protease cleaves the protein at a linker region between the A and B chains (Lord, J.M. et al., *FASAB Journal* 8:201-208 (1994)). The two chains, however, remain covalently attached by an interchain disulfide bond (cysteine 259 in the A chain to cysteine 4 in the B chain) and mature disulfide linked ricin is stored in protein bodies inside the plant cells. The A chain is inactive in proricin (O'Hare, M. et al., *FEBS Lett.* 273:200-204 (1990)) and it is inactive in the disulfide-linked mature ricin (Richardson, P.T. et al., *FEBS Lett.* 255:15-20 (1989)). The ribosomes of the castor bean plant are themselves susceptible to inactivation by ricin A chain; however, as there is no cell surface galactose to permit B chain recognition the A chain cannot re-enter the cell. The exact mechanism of A chain release and activation in target cell cytoplasm is not known (Lord, J.M. et al., *FASAB Journal* 8:201-208 (1994)). However, it is known that for activation to take place the disulfide bond between the A and B chains must be reduced and, hence, the linkage between subunits broken.

Diphtheria toxin is produced by *Corynebacterium diphtheriae* as a 535 amino acid polypeptide with a molecular weight of approximately 58kD (Greenfield, L. et al., *Proc. Natl. Acad. Sci. USA* 80:6853-6857 (1983); Pastan, I. et al., *Annu. Rev. Biochem.* 61:331-354 (1992); Collier, R.J. & Kandel, J., *J. Biol. Chem.* 246:1496-1503 (1971)). It is secreted as a single-chain polypeptide consisting of 2 functional domains. Similar to proricin, the N-terminal domain (A-chain) contains the cytotoxic moiety whereas the C-terminal domain (B-chain) is responsible for binding to the cells and facilitates toxin endocytosis. Conversely, the mechanism of cytotoxicity for diphtheria toxin is based on ADP-ribosylation of EF-2 thereby blocking protein synthesis and producing cell death. The 2 functional domains in diphtheria toxin are linked by an arginine-rich peptide sequence as well as a disulphide bond. Once the diphtheria toxin is internalized into the cell, the arginine-rich peptide linker is cleaved by trypsin-like enzymes and the

- 4 -

disulphide bond (Cys 186-201) is reduced. The cytotoxic domain is subsequently translocated into the cytosol substantially as described above for ricin and elicits ribosomal inhibition and cytotoxicity.

Pseudomonas exotoxin is also a 66kD single-chain toxin protein secreted by *Pseudomonas aeruginosa* with a similar mechanism of cytotoxicity to that of diphtheria toxin (Pastan, I. et al., *Annu. Rev. Biochem.* 61:331-354 (1992); Ogata, M. et al., *J. Biol. Chem.* 267:25396-25401 (1992); Vagil, M.L. et al., *Infect. Immunol.* 16:353-361 (1977)). *Pseudomonas* exotoxin consists of 3 conjoint functional domains. The first domain Ia (amino acids 1-252) is responsible for cell binding and toxin endocytosis, a second domain II (amino acids 253-364) is responsible for toxin translocation from the endocytic vesicle to the cytosol, and a third domain III (amino acids 400-613) is responsible for protein synthesis inhibition and cytotoxicity. After *Pseudomonas* exotoxin enters the cell, the liberation of the cytotoxic domain is effected by both proteolytic cleavage of a polypeptide sequence in the second domain (near Arg 279) and the reduction of the disulphide bond (Cys 265-287) in the endocytic vesicles. In essence, the overall pathway to cytotoxicity is analogous to diphtheria toxin with the exception that the toxin translocation domain in *Pseudomonas* exotoxin is structurally distinct.

Other toxins possessing distinct functional domains for cytotoxicity and cell binding/toxin translocation include abrin, modeccin and volkensin (Sandvig, K. et al., *Biochem. Soc. Trans.* 21:707-711 (1993)). Some toxins such as Shiga toxin and cholera toxin also have multiple polypeptide chains responsible for receptor binding and endocytosis.

The ricin gene has been cloned and sequenced, and the X-ray crystal structures of the A and B chains have been described (Rutenber, E. et al. *Proteins* 10:240-250 (1991); Weston et al., *Mol. Bio.* 244:410-422, 1994; Lamb and Lord, *Eur. J. Biochem.* 14:265 (1985); Halling, K. et al. *Nucleic Acids Res.* 13:8019 (1985)). Similarly, the genes for

diphtheria toxin and *Pseudomonas* exotoxin have been cloned and sequenced, and the 3-dimensional structures of the toxin proteins have been elucidated and described (Columblatti, M. et al., *J. Biol. Chem.* 261:3030-3035 (1986); Allured, V.S. et al., *Proc. Natl. Acad. Sci. USA* 83:1320-1324 (1986); Gray, G.L. et al., *Proc. Natl. Acad. Sci. USA* 81:2645-2649 (1984); Greenfield, L. et al., *Proc. Natl. Acad. Sci. USA* 80:6853-6857 (1983); Collier, R.J. et al., *J. Biol. Chem.* 257:5283-5285 (1982)).

The potential of bacterial and plant toxins for inhibiting mammalian retroviruses, particularly acquired immunodeficiency syndrome (AIDS), has been investigated. Bacterial toxins such as *Pseudomonas* exotoxin-A and subunit A of diphtheria toxin; dual chain ribosomal inhibitory plant toxins such as ricin, and single chain ribosomal inhibitory proteins such as trichosanthin and pokeweed antiviral protein have been used for the elimination of HIV infected cells (Olson et al., *AIDS Res. and Human Retroviruses* 7:1025-1030 (1991)). The high toxicity of these toxins for mammalian cells, combined with a lack of specificity of action poses a major problem to the development of pharmaceuticals incorporating the toxins, such as immunotoxins.

Due to their extreme toxicity there has been much interest in making ricin-based immunotoxins as therapeutic agents for specifically destroying or inhibiting infected or tumourous cells or tissues (Vitetta et al., *Science* 238:1098-1104(1987)). An immunotoxin is a conjugate of a specific cell binding component, such as a monoclonal antibody or growth factor and the toxin in which the two protein components are covalently linked. Generally, the components are chemically coupled. However, the linkage may also be a peptide or disulfide bond. The antibody directs the toxin to cell types presenting a specific antigen thereby providing a specificity of action not possible with the natural toxin. Immunotoxins have been made both with the entire ricin molecule (i.e. both chains) and with the ricin A chain alone (Spooner et al., *Mol. Immunol.* 31:117-125, (1994)).

- 6 -

Immunotoxins made with the ricin dimer (IT-Rs) are more potent toxins than those made with only the A chain (IT-As). The increased toxicity of IT-Rs is thought to be attributed to the dual role of the B chains in binding to the cell surface and in translocating the A chain to the cytosolic compartment of the target cell (Vitetta et al., *Science* 238:1098-1104 (1987); Vitetta & Thorpe, *Seminars in Cell Biology* 2:47-58 (1991)). However, the presence of the B chain in these conjugates also promotes the entry of the immunotoxin into nontarget cells. Even small amounts of B chain may override the specificity of the cell-binding component as the B chain will bind nonspecifically to galactose associated with N-linked carbohydrates, which is present on most cells. IT-As are more specific and safer to use than IT-Rs. However, in the absence of the B chain the A chain has greatly reduced toxicity. Due to the reduced potency of IT-As as compared to IT-Rs, large doses of IT-As must be administered to patients. The large doses frequently cause immune responses and production of neutralizing antibodies in patients (Vitetta et al., *Science* 238:1098-1104 (1987)). IT-As and IT-Rs both suffer from reduced toxicity as the A chain is not released from the conjugate into the target cell cytoplasm.

A number of immunotoxins have been designed to recognize antigens on the surfaces of tumour cells and cells of the immune system (Pastan et al., *Annals New York Academy of Sciences* 758:345-353 (1995)). A major problem with the use of such immunotoxins is that the antibody component is its only targeting mechanism and the target antigen is often found on non-target cells (Vitetta et al., *Immunology Today* 14:252-259 (1993)). Also, the preparation of a suitable specific cell binding component may be problematic. For example, antigens specific for the target cell may not be available and many potential target cells and infective organisms can alter their antigenic make up rapidly to avoid immune recognition. In view of the extreme toxicity of proteins such as ricin, the lack of

specificity of the immunotoxins may severely limit their usefulness as therapeutics for the treatment of cancer and infectious diseases.

The insertion of intramolecular protease cleavage sites between the cytotoxic and cell-binding components of a toxin can mimic the way that the natural toxin is activated. European patent application no. 466,222 describes the use of maize-derived pro-proteins which can be converted into active form by cleavage with extracellular blood enzymes such as factor Xa, thrombin or collagenase. Garred, O. et al. (*J. Biol. Chem.* 270:10817-10821 (1995)) documented the use of a ubiquitous calcium-dependent serine protease, furin, to activate shiga toxin by cleavage of the trypsin-sensitive linkage between the cytotoxic A-chain and the pentamer of cell-binding B-units. Westby et al. (*Bioconjugate Chem.* 3:375-381 (1992)) documented fusion proteins which have a specific cell binding component and proricin with a protease sensitive cleavage site specific for factor Xa within the linker sequence. O'Hare et al. (*FEBS Lett.* 273:200-204 (1990)) also described a recombinant fusion protein of RTA and staphylococcal protein A joined by a trypsin-sensitive cleavage site. In view of the ubiquitous nature of the extracellular proteases utilized in these approaches, such artificial activation of the toxin precursor or immunotoxin does not confer a mechanism for intracellular toxin activation and the problems of target specificity and adverse immunological reactions to the cell-binding component of the immunotoxin remain.

In a variation of the approach of insertion of intramolecular protease cleavage sites on proteins which combine a binding chain and a toxic chain, Leppla, S.H. et al. (*Bacterial Protein Toxins zbl.bakt.suppl.* 24:431-442 (1994)) suggest the replacement of the native cleavage site of the protective antigen (PA) produced by *Bacillus anthracis* with a cleavage site that is recognized by cells that contain a particular protease. PA, recognizes, binds, and thereby assists in the internalization of lethal factor (LF) and edema toxin (ET). also produced by *Bacillus anthracis*. However, this approach is wholly dependent on

- 8 -

the availability of LF, or ET and PA all being localized to cells wherein the modified PA can be activated by the specific protease. It does not confer a mechanism for intracellular toxin activation and presents a problem of ensuring sufficient quantities of toxin for internalization in
5 target cells.

The *in vitro* activation of a *Staphylococcus*-derived pore-forming toxin, α -hemolysin by extracellular tumour-associated proteases has been documented (Panchel, R.G. et al., *Nature Biotechnology* 14:852-857 (1996)). Artificial activation of α -hemolysin *in*
10 *vitro* by said proteases was reported but the actual activity and utility of α -hemolysin in the destruction of target cells were not demonstrated.

Hemolysin does not inhibit protein synthesis but is a heptameric transmembrane pore which acts as a channel to allow leakage of molecules up to 3 kD thereby disrupting the ionic balances of
15 the living cell. The α -hemolysin activation domain is likely located on the outside of the target cell (for activation by extracellular proteases). The triggering mechanism in the disclosed hemolysin precursor does not involve the intracellular proteolytic cleavage of 2 functionally distinct domains. Also, the proteases used for the α -hemolysin
20 activation are ubiquitously secreted extracellular proteases and toxin activation would not be confined to activation in the vicinity of diseased cells. Such widespread activation of the toxin does not confer target specificity and limits the usefulness of said α -hemolysin toxin as therapeutics due to systemic toxicity.

25 A variety of proteases specifically associated with malignancy, viral infections and parasitic infections have been identified and described. For example, cathepsin is a family of serine, cysteine or aspartic endopeptidases and exopeptidases which has been implicated to play a primary role in cancer metastasis (Schwartz, M.K.,
30 *Clin. Chim. Acta* 237:67-78 (1995); Spiess, E. et al., *J. Histochem.*

Cytochem. 42:917-929 (1994); Scarborough, P.E. et al., *Protein Sci.* 2:264-276 (1993); Sloane, B.F. et al., *Proc. Natl. Acad. Sci. USA* 83:2483-2487 (1986); Mikkelsen, T. et al., *J. Neurosurg* 83:285-290 (1995)). Matrix metalloproteinases (MMPs or matrixins) are zinc-dependent proteinases consisting of collagenases, matrilysin, stromelysins, gelatinases and macrophage elastase (Krane, S.M., *Ann. N.Y. Acad. Sci.* 732:1-10 (1994); Woessner, J.F., *Ann. N.Y. Acad. Sci.* 732:11-21 (1994); Carvalho, K. et al., *Biochem. Biophys. Res. Comm.* 191:172-179 (1993); Nakano, A. et al. *J. of Neurosurg*, 83:298-307 (1995); Peng, K-W, et al. *Human Gene Therapy*, 8:729-738 (1997); More, D.H. et al. *Gynaecologic Oncology*, 65:78-82 (1997)). These proteases are involved in pathological matrix remodeling. Under normal physiological conditions, regulation of matrixin activity is effected at the level of gene expression. Enzymatic activity is also controlled stringently by tissue inhibitors of metalloproteinases (TIMPs) (Murphy, G. et al., *Ann. N.Y. Acad. Sci.* 732:31-41 (1994)). The expression of MMP genes is reported to be activated in inflammatory disorders (e.g. rheumatoid arthritis) and malignancy.

In malaria, parasitic serine and aspartic proteases are involved in host erythrocyte invasion by the *Plasmodium* parasite and in hemoglobin catabolism by intraerythrocytic malaria (O'Dea, K.P. et al., *Mol. Biochem. Parasitol.* 72:111-119 (1995); Blackman, M.J. et al., *Mol. Biochem. Parasitol.* 62:103-114 (1993); Cooper, J.A. et al., *Mol. Biochem. Parasitol.* 56:151-160 (1992); Goldberg, D.E. et al., *J. Exp. Med.* 173:961-969 (1991)). *Schistosoma mansoni* is also a pathogenic parasite which causes schistosomiasis or bilharzia. Elastinolytic proteinases have been associated specifically with the virulence of this particular parasite (McKerrow, J.H. et al., *J. Biol. Chem.* 260:3703-3707 (1985)).

Welch, A.R. et al. (*Proc. Natl. Acad. Sci. USA* 88:10797-10800 (1991)) has described a series of viral proteases which are specifically associated with human cytomegalovirus, human herpesviruses, Epstein-Barr virus, varicella zoster virus-I. and

- 10 -

infectious laryngotracheitis virus. These proteases possess similar substrate specificity and play an integral role in viral scaffold protein restructuring in capsid assembly and virus maturation. Other viral proteases serving similar functions have also been documented for

5 human T-cell leukemia virus (Blaha, I. et al., *FEBS Lett.* 309:389-393 (1992); Pettit, S.C. et al., *J. Biol. Chem.* 266:14539-14547 (1991)), hepatitis viruses (Hirowatari, Y. et al., *Anal. Biochem.* 225:113-120 (1995); Hirowatari, Y. et al., *Arch. Virol.* 133:349-356 (1993); Jewell, D.A. et al., *Biochemistry* 31:7862-7869 (1992)), poliomyelitis virus (Weidner, J.R. et al., *Arch. Biochem. Biophys.* 286:402-408 (1991)), and human rhinovirus

10 (Long, A.C. et al., *FEBS Lett.* 258:75-78 (1989)).

Candida yeasts are dimorphic fungi which are responsible for a majority of opportunistic infections in AIDS patients (Holmberg, K. and Myer, R., *Scand. J. Infect. Dis.* 18:179-192 (1986)). Aspartic

15 proteinases have been associated specifically with numerous virulent strains of *Candida* including *Candida albican*, *Candida tropicalis*, and *Candida parapsilosis* (Abad-Zapatero, C. et al., *Protein Sci.* 5:640-652 (1996); Cutfield, S.M. et al., *Biochemistry* 35:398-410 (1995); Ruchel, R. et al., *Zentralbl. Bakteriол. Mikrobiol Hyg. I Abt. Orig. A.* 255:537-548 (1983);

20 Remold, H. et al., *Biochim. Biophys. Acta* 167:399-406 (1968)), and the levels of these enzymes have been correlated with the lethality of the strain (Schreiber, B. et al., *Diagn. Microbiol. Infect. Dis.* 3:1-5 (1985)).

SUMMARY OF THE INVENTION

The invention relates to novel recombinant toxic

25 proteins which are specifically toxic to diseased cells but do not depend for their specificity of action on a specific cell binding component. The recombinant proteins of the invention have an A chain of a ricin-like toxin linked to a B chain by a synthetic linker sequence which may be cleaved specifically by a protease localised in cells or tissues affected by a

30 specific disease to liberate the toxic A chain thereby selectively inhibiting or destroying the diseased cells or tissues. The term diseased

- 11 -

cells as used herein, includes cells affected by cancer, or infected by fungi, or viruses, including retroviruses, or parasites.

Toxin targeting using the recombinant toxic proteins of the invention takes advantage of the fact that many DNA viruses exploit host cellular transport mechanisms to escape immunological destruction. This is achieved by enhancing the retrograde translocation of host major histocompatibility complex (MHC) type I molecules from the endoplasmic reticulum into the cytoplasm (Bonifacino, J.S., *Nature* 384: 405-406 (1996); Wiertz, E.J. et al., *Nature* 384: 432-438 (1996)). The facilitation of retrograde transport in diseased cells by the virus can enhance the transcytosis and cytotoxicity of a recombinant toxic protein of the present invention thereby further reducing non-specific cytotoxicity and improving the overall safety of the product.

The recombinant toxic proteins of the present invention may be used to treat diseases including various forms of cancer such as T- and B-cell lymphoproliferative diseases, ovarian cancer, pancreatic cancer, head and neck cancer, squamous cell carcinoma, gastrointestinal cancer, breast cancer, prostate cancer, non small cell lung cancer, malaria, and diverse viral disease states associated with infection with human cytomegalovirus, hepatitis virus, herpes virus, human rhinovirus, infectious laryngotracheitis virus, poliomyelitis virus, or varicella zoster virus.

In one aspect, the present invention provides a purified and isolated nucleic acid having a nucleotide sequence encoding an A chain of a ricin-like toxin, a B chain of a ricin-like toxin and a heterologous linker amino acid sequence, linking the A and B chains. The linker sequence is not a native linker sequence of a ricin-like toxin, but rather a synthetic heterologous linker sequence containing a cleavage recognition site for a disease-specific protease. The A and or the B chain may be those of ricin.

In an embodiment, of the invention the cleavage recognition site is the cleavage recognition site for a cancer-associated

protease. In particular embodiments, the linker amino acid sequence comprises SLLKSRMVPNFN or SLLIARRMPNFN cleaved by cathepsin B; SKLVQASASGVN or SSYLKASDAPDN cleaved by an Epstein-Barr virus protease; RPKPQQFFGLMN cleaved by MMP-3 (stromelysin);

5 SLRPLALWRSFN cleaved by MMP-7 (matrilysin); SPQGIAGQRNFN cleaved by MMP-9; DVDERDVRGFASFL cleaved by a thermolysin-like MMP; SLPLGLWAPNFN cleaved by matrix metalloproteinase 2(MMP-2) ; SLLIFRSWANFN cleaved by cathepsin L; SGVVIATVIVIT cleaved by cathepsin D; SLGPQGIWGQFN cleaved by matrix metalloproteinase

10 1(MMP-1); KKSPGRVVGGSV cleaved by urokinase-type plasminogen activator; PQGLLGAPGILG cleaved by membrane type 1 matrixmetalloproteinase (MT-MMP);

HGPEGLRVGFYESDVMGRGHARLVHVEEPHT cleaved by stromelysin 3 (or MMP-11), thermolysin, fibroblast collagenase and stromelysin-1;

15 GPQGLAGQRGIV cleaved by matrix metalloproteinase 13 (collagenase-3); GGSGQRGRKALE cleaved by tissue-type plasminogen activator(tPA); SLSALLSSDIFN cleaved by human prostate-specific antigen; SLPRFKIIGGFN cleaved by kallikrein (hK3); SLLGIAVPGNFN cleaved by neutrophil elastase; and FFKNIVTPRTPP cleaved by calpain

20 (calcium activated neutral protease). The nucleic acid sequences for ricin A and B chains with each of the linker sequences are shown in Figures 2D, 35C, 3D, 4D, 5D, 6D, 16D, 17D, 34C, 36C , 37C, 38C , 39C, 40C, 41C, 42C , 43C, 44C, 45C, 46C and 47C, respectively.

In another embodiment, the cleavage recognition site is

25 the cleavage recognition site for a protease associated with the malaria parasite, *Plasmodium falciparum*. In particular embodiments, the linker amino acid sequence comprises QVVQLQNYDEED; LPIFGESEDNDE; QVVTGEAISVTM; ALERTFLSFPTN or KFQDMLNISQHQ. The nucleic nucleotide sequences for ricin A and B

30 chains with each of the linker sequences are shown in Figures 7D, 8D, 9D, 10D, and 11D.

In a another embodiment, the cleavage recognition site is the cleavage recognition site for a viral protease. The linker sequences preferably comprise the sequence Y-X-Y-A-Z wherein X is valine or leucine, Y is a polar amino acid, and Z is serine, asparagine or valine.

5 In particular embodiments, the linker amino acid sequence comprises SGVVNASCRAN or SSYVKASVSPEN cleaved by a human cytomegalovirus protease; SALVNASSAHVN or STYLQASEKFKN cleaved by a herpes simplex 1 virus protease; SSILNASVPNFN cleaved by a human herpes virus 6 protease; SQDVNAVEASSN or
10 SVYLQASTGYGN cleaved by a varicella zoster virus protease; or SKYLQANEVITN cleaved by an infectious laryngotracheitis virus protease. The nucleic nucleotide sequences for ricin A and B chains with each of the linker sequences are shown in Figures 12D, 13D, 14D, 15D, 18D, 19D, 20D, and 22D.

15 In another embodiment, the cleavage recognition site is the cleavage recognition site for a hepatitis A viral protease. In particular embodiments, the linker amino acid sequence comprises SELRTQSFSNWN or SELWSQGIDDDN cleaved by a hepatitis A virus protease. The nucleic nucleotide sequences for ricin A and B chains
20 with each of the linker sequences are shown in Figures 23D or 24D.

In another embodiment, the cleavage recognition site is the cleavage recognition site for a hepatitis C viral protease. In particular embodiments, the linker amino acid sequence comprises DLEVVTSTWVFN, DEMEECASHLFN, EDVVCCSMSYFN or
25 KGWRLAPITAY cleaved by a hepatitis C virus protease. The nucleic nucleotide sequences for ricin A and B chains with each of the linker sequences are shown in Figures 30C, 31C, 32C and 33C.

In another embodiment, the cleavage recognition site is the cleavage recognition site for a *Candida* fungal protease. In particular
30 embodiments, the linker amino acid sequence is SKPAKFFRLNFN, SKPIEFFRLNFN or SKPAEFFALNFN cleaved by *Candida* aspartic

- 14 -

protease. The nucleic nucleotide sequences for ricin A and B chains with the first linker sequence are shown in Figures 25D.

The present invention also provides a plasmid incorporating the nucleic acid of the invention. In an embodiment, the plasmid has the restriction map as shown in Figures 2A, 3A, 4A, 5A, 6A, 7A, 8A, 9A, 10A, 11A, 12A, 13A, 14A, 15A, 16A, 17A, 18A, 19A, 20A, 21A, 22A, 23A, 24A, or 25A.

In another embodiment, the present invention provides a baculovirus transfer vector incorporating the nucleic acid of the invention. In particular embodiments, the invention provides a baculovirus transfer vector having the DNA sequence as shown in Figure 1.

In a further embodiment, the present invention provides a baculovirus transfer vector incorporating the nucleic acid of the invention. In particular embodiments, the invention provides a baculovirus transfer vector having the restriction map as shown in Figures 2C, 3C, 4C, 5C, 6C, 7C, 8C, 9C, 10C, 11C, 12C, 13C, 14C, 15C, 16C, 17C, 18C, 19C, 20C, 21C, 22C, 23C, 24C, 25C, 30A, 31A, 32A, 33A, 34A, 35A, 36A, 37A, 38A, 39A, 40A, 41A, 42A, 43A, 44A, 45A, 46A, or 47A. or having the DNA sequence as shown in Figure 1.

In a further aspect, the present invention provides a recombinant protein comprising an A chain of a ricin-like toxin, a B chain of a ricin-like toxin and a heterologous linker amino acid sequence, linking the A and B chains, wherein the linker sequence contains a cleavage recognition site for a disease-specific protease (e.g., a cancer, viral, parasitic, or fungal protease). The A and/or the B chain may be those of ricin. In an embodiment, the cleavage recognition site is the cleavage recognition site for a cancer, viral or parasitic protease substantially as described above. In a particular embodiment, the cancer is T-cell or B-cell lymphoproliferative disease. In another particular embodiment, the virus is human cytomegalovirus, Epstein-Barr virus, hepatitis virus, herpes virus, human rhinovirus, infectious

laryngotracheitis virus, poliomyelitis virus, or varicella zoster virus. In a further particular embodiment, the parasite is *Plasmodium falciparum*.

5 In a further aspect, the invention provides a pharmaceutical composition for treating a fungal infection, such as *Candida*, in a mammal comprising the recombinant protein of the invention and a pharmaceutically acceptable carrier, diluent or excipient.

10 In yet another aspect, the invention provides a method of inhibiting or destroying cells affected by a disease, which cells are associated with a disease specific protease, including cancer or infection with a virus, fungus, or a parasite each of which has a specific protease, comprising the steps of preparing a recombinant protein of the invention having a heterologous linker sequence which contains a
15 cleavage recognition site for the disease-specific protease and administering the recombinant protein to the cells. In an embodiment, the cancer is T-cell or B-cell lymphoproliferative disease, ovarian cancer, pancreatic cancer, head and neck cancer, squamous cell carcinoma, gastrointestinal cancer, breast cancer, prostate cancer, non
20 small cell lung cancer. In another embodiment, the virus is human cytomegalovirus, Epstein-Barr virus, hepatitis virus, herpes virus, human rhinovirus, human T-cell leukemia virus, infectious laryngotracheitis virus, poliomyelitis virus, or varicella zoster virus. In another embodiment, the parasite is *Plasmodium falciparum*.

25 The present invention also relates to a method of treating a mammal with disease wherein cells affected by the disease are associated with a disease specific protease, including cancer or infection with a virus, fungus, or a parasite each of which has a specific protease by administering an effective amount of one or more recombinant
30 proteins of the invention to said mammal.

Still further, a process is provided for preparing a pharmaceutical for treating a mammal with disease wherein cells

- 16 -

affected by the disease are associated with a disease specific protease, including cancer or infection with a virus, fungus, or a parasite each of which has a specific protease comprising the steps of preparing a purified and isolated nucleic acid having a nucleotide sequence encoding an A chain of a ricin-like toxin, a B chain of a ricin-like toxin and a heterologous linker amino acid sequence, linking the A and B chains, wherein the linker sequence contains a cleavage recognition site for the disease-specific protease; introducing the nucleic acid into a host cell; expressing the nucleic acid in the host cell to obtain a recombinant protein comprising an A chain of a ricin-like toxin, a B chain of a ricin-like toxin and a heterologous linker amino acid sequence, linking the A and B chains wherein the linker sequence contains the cleavage recognition site for the disease-specific protease; and suspending the protein in a pharmaceutically acceptable carrier, diluent or excipient.

In an embodiment, a process is provided for preparing a pharmaceutical for treating a mammal with disease wherein cells affected by the disease are associated with a disease specific protease, including cancer or infection with a virus, fungus, or a parasite each of which has a specific protease comprising the steps of identifying a cleavage recognition site for the protease; preparing a recombinant protein comprising an A chain of a ricin-like toxin, a B chain of a ricin-like toxin and a heterologous linker amino acid sequence, linking the A and B chains wherein the linker sequence contains the cleavage recognition site for the protease and suspending the protein in a pharmaceutically acceptable carrier, diluent or excipient.

In a further aspect, the invention provides a pharmaceutical composition for treating for treating a mammal with disease wherein cells affected by the disease are associated with a disease specific protease, including cancer or infection with a virus, fungus, or a parasite comprising the recombinant protein of the invention and a pharmaceutically acceptable carrier, diluent or excipient.

Other features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples while indicating preferred embodiments of the invention are
5 given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

DESCRIPTION OF THE DRAWINGS

The invention will be better understood with reference to
10 the drawings in which:

Figure 1 shows the DNA sequence of the baculovirus transfer vector, pVL1393;

Figure 2A summarizes the cloning strategy used to generate the pAP-213 construct;

15 Figure 2B shows the nucleotide sequence of the Cathepsin B linker regions of pAP-213;

Figure 2C shows the subcloning of the Cathepsin B linker variant into a baculovirus transfer vector;

Figure 2D shows the DNA sequence of the pAP-214 insert
20 containing ricin and the Cathepsin B linker;

Figure 3A summarizes the cloning strategy used to generate the pAP-215 construct;

Figure 3B shows the nucleotide sequence of the MMP-3 linker regions of pAP-215;

25 Figure 3C shows the subcloning of the MMP-3 linker variant into a baculovirus transfer vector;

Figure 3D shows the DNA sequence of the pAP-216 insert containing ricin and the MMP-3 linker;

Figure 4A summarizes the cloning strategy used to
30 generate the pAP-217 construct;

Figure 4B shows the nucleotide sequence of the MMP-7 linker regions of pAP-217;

- 18 -

Figure 4C shows the subcloning of the MMP-7 linker variant into a baculovirus transfer vector;

Figure 4D shows the DNA sequence of the pAP-218 insert containing ricin and the MMP-7 linker;

5 Figure 5A summarizes the cloning strategy used to generate the pAP-219 construct;

Figure 5B shows the nucleotide sequence of the MMP-9 linker regions of pAP-219;

10 Figure 5C shows the subcloning of the MMP-9 linker variant into a baculovirus transfer vector;

Figure 5D shows the DNA sequence of the pAP-220 insert containing ricin and the MMP-9 linker.

Figure 6A summarizes the cloning strategy used to generate the pAP-221 construct;

15 Figure 6B shows the nucleotide sequence of the thermolysin-like MMP linker regions of pAP-221;

Figure 6C shows the subcloning of the thermolysin-like MMP linker variant into a baculovirus transfer vector.

20 Figure 6D shows the DNA sequence of the pAP-222 insert containing ricin and the thermolysin-like MMP linker;

Figure 7A summarizes the cloning strategy used to generate the pAP-223 construct;

Figure 7B shows the nucleotide sequence of the Plasmodium falciparum-A linker regions of pAP-223;

25 Figure 7C shows the subcloning of the Plasmodium falciparum-A linker variant into a baculovirus transfer vector;

Figure 7D shows the DNA sequence of the pAP-224 insert containing ricin and the Plasmodium falciparum-A linker;

30 Figure 8A summarizes the cloning strategy used to generate the pAP-225 construct;

Figure 8B shows the nucleotide sequence of the Plasmodium falciparum-B linker regions of pAP-225;

- 19 -

Figure 8C shows the subcloning of the Plasmodium falciparum-B linker variant into a baculovirus transfer vector;

Figure 8D shows the DNA sequence of the pAP-226 insert containing ricin and the Plasmodium falciparum-B linker;

5 Figure 9A summarizes the cloning strategy used to generate the pAP-227 construct;

Figure 9B shows the nucleotide sequence of the Plasmodium falciparum-C linker regions of pAP-227;

10 Figure 9C shows the subcloning of the Plasmodium falciparum-C linker variant into a baculovirus transfer vector;

Figure 9D shows the DNA sequence of the pAP-228 insert containing ricin and the Plasmodium falciparum-C linker;

Figure 10A summarizes the cloning strategy used to generate the pAP-229 construct;

15 Figure 10B shows the nucleotide sequence of the Plasmodium falciparum-D linker regions of pAP-229;

Figure 10C shows the subcloning of the Plasmodium falciparum-D linker variant into a baculovirus transfer vector;

20 Figure 10D shows the DNA sequence of the pAP-230 insert containing ricin and the Plasmodium falciparum-D linker;

Figure 11A summarizes the cloning strategy used to generate the pAP-231 construct;

Figure 11B shows the nucleotide sequence of the Plasmodium falciparum-E linker regions of pAP-231;

25 Figure 11C shows the subcloning of the Plasmodium falciparum-E linker variant into a baculovirus transfer vector;

Figure 11D shows the DNA sequence of the pAP-232 insert containing ricin and the Plasmodium falciparum-E linker;

30 Figure 12A summarizes the cloning strategy used to generate the pAP-233 construct;

Figure 12B shows the nucleotide sequence of the HSV-A linker regions of pAP-233;

- 20 -

Figure 12C shows the subcloning of the HSV-A linker variant into a baculovirus transfer vector;

Figure 12D shows the DNA sequence of the pAP-234 insert containing ricin and the HSV-A linker;

5 Figure 13A summarizes the cloning strategy used to generate the pAP-235 construct;

Figure 13B shows the nucleotide sequence of the HSV-B linker regions of pAP-235;

10 Figure 13C shows the subcloning of the HSV-B linker variant into a baculovirus transfer vector;

Figure 13D shows the DNA sequence of the pAP-236 insert containing ricin and the HSV-B linker;

Figure 14A summarizes the cloning strategy used to generate the pAP-237 construct;

15 Figure 14B shows the nucleotide sequence of the VZV-A linker regions of pAP-237;

Figure 14C shows the subcloning of the VZV-A linker variant into a baculovirus transfer vector;

20 Figure 14D shows the DNA sequence of the pAP-238 insert containing ricin and the VZV-A linker;

Figure 15A summarizes the cloning strategy used to generate the pAP-239 construct;

Figure 15B shows the nucleotide sequence of the VZV-B linker regions of pAP-239;

25 Figure 15C shows the subcloning of the VZV-B linker variant into a baculovirus transfer vector;

Figure 15D shows the DNA sequence of the pAP-240 insert containing ricin and the VZV-B linker;

30 Figure 16A summarizes the cloning strategy used to generate the pAP-241 construct;

Figure 16B shows the nucleotide sequence of the EBV-A linker regions of pAP-241;

- 21 -

Figure 16C shows the subcloning of the EBV-A linker variant into a baculovirus transfer vector;

Figure 16D shows the DNA sequence of the pAP-242 insert containing ricin and the EBV-A linker;

5 Figure 17A summarizes the cloning strategy used to generate the pAP-243 construct;

Figure 17B shows the nucleotide sequence of the EBV-B linker regions of pAP-243;

10 Figure 17C shows the subcloning of the EBV-B linker variant into a baculovirus transfer vector;

Figure 17D shows the DNA sequence of the pAP-244 insert containing ricin and the EBV-B linker;

Figure 18A summarizes the cloning strategy used to generate the pAP-245 construct;

15 Figure 18B shows the nucleotide sequence of the CMV-A linker regions of pAP-245;

Figure 18C shows the subcloning of the CMV-A linker variant into a baculovirus transfer vector;

20 Figure 18D shows the DNA sequence of the pAP-246 insert containing ricin and the CMV-A linker;

Figure 19A summarizes the cloning strategy used to generate the pAP-247 construct;

Figure 19B shows the nucleotide sequence of the CMV-B linker regions of pAP-247;

25 Figure 19C shows the subcloning of the CMV-B linker variant into a baculovirus transfer vector;

Figure 19D shows the DNA sequence of the pAP-248 insert containing ricin and the CMV-B linker.

30 Figure 20A summarizes the cloning strategy used to generate the pAP-249 construct;

Figure 20B shows the nucleotide sequence of the HHV-6 linker regions of pAP-249;

- 22 -

Figure 20C shows the subcloning of the HHV-6 linker variant into a baculovirus transfer vector;

Figure 20D shows the DNA sequence of the pAP-250 insert containing ricin and the HHV-6 linker;

5 Figure 21 shows the amino acid sequences of the wild type ricin linker and cancer protease-sensitive amino acid linkers contained in pAP-213 to pAP-222 and linkers pAP-241 to pAP-244;

Figure 22A summarizes the cloning strategy used to generate the pAP-253 construct;

10 Figure 22B shows the nucleotide sequence of the ILV linker regions of pAP-253;

Figure 22C shows the subcloning of the ILV linker variant into a baculovirus transfer vector;

15 Figure 22D shows the DNA sequence of the pAP-254 insert containing ricin and the ILV linker;

Figure 23A summarizes the cloning strategy used to generate the pAP-257 construct;

Figure 23B shows the nucleotide sequence of the HAV-A linker regions of pAP-257;

20 Figure 23C shows the subcloning of the HAV-A linker variant into a baculovirus transfer vector;

Figure 23D shows the DNA sequence of the pAP-258 insert containing ricin and the HAV-A linker;

25 Figure 24A summarizes the cloning strategy used to generate the pAP-255 construct;

Figure 24B shows the nucleotide sequence of the HAV-B linker regions of pAP-255;

Figure 24C shows the subcloning of the HAV-B linker variant into a baculovirus transfer vector;

30 Figure 24D shows the DNA sequence of the pAP-256 insert containing ricin and the HAV-B linker;

Figure 25A summarizes the cloning strategy used to generate the pAP-259 construct;

Figure 25B shows the nucleotide sequence of the CAN linker regions of pAP-259;

5 Figure 25C shows the subcloning of the CAN linker variant into a baculovirus transfer vector;

Figure 25D shows the DNA sequence of the pAP-260 insert containing ricin and the CAN linker;

10 Figure 26 shows the amino acid sequences of the wild type ricin linker and *Plasmodium falciparum* protease-sensitive amino acid linkers contained in linkers pAP-223 to pAP-232;

 Figure 27 shows the amino acid sequences of the wild type ricin linker and the viral protease-sensitive amino acid linkers contained in pAP-233 to pAP-240, pAP-245-pAP-248, pAP-253 to pAP-
15 258;

Figure 28 shows the amino acid sequences of the wild type ricin linker and the *Candida* aspartic protease-sensitive amino acid linker contained in pAP-259 to pAP-264;

20 Figure 29 describes an alternative mutagenesis and subcloning strategy to provide a baculovirus transfer vector containing the ricin-like toxin variant gene; and

Figure 30A summarizes the cloning strategy used to generate the pAP-262 construct;

25 Figure 30B shows the nucleotide sequence of the HCV-A linker region of pAP-262;

Figure 30C shows the DNA sequence of the pAP-262 insert;

Figure 30D shows the amino acid sequence comparison of mutant preproricin linker region HCV-A to wild type;

30 Figure 31A summarizes the cloning strategy used to generate the pAP-264 construct;

- 24 -

Figure 31B shows the nucleotide sequence of the HCV-B linker region of pAP-264;

Figure 31C shows the DNA sequence of the pAP-264 insert;

5 Figure 31D shows the amino acid sequence comparison of mutant preproricin linker region HCV-B to wild type;

Figure 32A summarizes the cloning strategy used to generate the pAP-266 construct;

10 Figure 32B shows the nucleotide sequence of the HCV-C linker region of pAP-266;

Figure 32C shows the DNA sequence of the pAP-266 insert;

Figure 32D shows the amino acid sequence comparison of mutant preproricin linker region HCV-C to wild type;

15 Figure 33A summarizes the cloning strategy used to generate the pAP-268 construct;

Figure 33B shows the nucleotide sequence of the HCV-D linker region of pAP-268;

20 Figure 33C shows the DNA sequence of the pAP-268 insert;

Figure 33D shows the amino acid sequence comparison of mutant preproricin linker region HCV-D to wild type;

Figure 34A summarizes the cloning strategy used to generate the pAP-270 construct;

25 Figure 34B shows the nucleotide sequence of the MMP-2 linker region of pAP-270;

Figure 34C shows the DNA sequence of the pAP-270 insert;

30 Figure 34D shows the amino acid sequence comparison of mutant preproricin linker region of MMP-2 to wild type;

Figure 35A summarizes the cloning strategy used to generate the pAP-272 construct;

- 25 -

Figure 35B shows the nucleotide sequence of the Cathepsin B (Site 2) linker region of pAP-272;

Figure 35C shows the DNA sequence of the pAP-272 insert;

5 Figure 35D shows the amino acid sequence comparison of mutant preproricin linker region of Cathepsin B (Site 2) to wild type;

Figure 36A summarizes the cloning strategy used to generate the pAP-274 construct;

10 Figure 36B shows the nucleotide sequence of the Cathepsin L linker region of pAP-274;

Figure 36C shows the DNA sequence of the pAP-274 insert;

Figure 36D shows the amino acid sequence comparison of mutant preproricin linker region of Cathepsin L to wild type;

15 Figure 37A summarizes the cloning strategy used to generate the pAP-276 construct;

Figure 37B shows the nucleotide sequence of the Cathepsin D linker region of pAP-276;

20 Figure 37C shows the DNA sequence of the pAP-276 insert;

Figure 37D shows the amino acid sequence comparison of mutant preproricin linker region of Cathepsin D to wild type;

Figure 38A summarizes the cloning strategy used to generate the pAP-278 construct;

25 Figure 38B shows the nucleotide sequence of the MMP-1 linker region of pAP-278;

Figure 38C shows the DNA sequence of the pAP-278 insert;

30 Figure 38D shows the amino acid sequence comparison of mutant preproricin linker region of MMP-1 to wild type;

Figure 39A summarizes the cloning strategy used to generate the pAP-280 construct;

- 26 -

Figure 39B shows the nucleotide sequence of the Urokinase-Type Plasminogen Activator linker region of pAP-280;

Figure 39C shows the DNA sequence of the pAP-280 insert;

5 Figure 39D shows the amino acid sequence comparison of mutant preproricin linker region of Urokinase-Type Plasminogen Activator to wild type;

Figure 40A summarizes the cloning strategy used to generate the pAP-282 construct;

10 Figure 40B shows the nucleotide sequence of the MT-MMP linker region of pAP-282;

Figure 40C shows the DNA sequence of the pAP-282 insert;

15 Figure 40D shows the amino acid sequence comparison of mutant preproricin linker region of MT-MMP to wild type;

Figure 41A summarizes the cloning strategy used to generate the pAP-284 construct;

Figure 41B shows the nucleotide sequence of the MMP-11 linker region of pAP-284;

20 Figure 41C shows the DNA sequence of the pAP-284 insert;

Figure 41D shows the amino acid sequence comparison of mutant preproricin linker region of MMP-11 to wild type;

25 Figure 42A summarizes the cloning strategy used to generate the pAP-286 construct;

Figure 42B shows the nucleotide sequence of the MMP-13 linker region of pAP-286;

Figure 42C shows the DNA sequence of the pAP-286 insert;

30 Figure 42D shows the amino acid sequence comparison of mutant preproricin linker region of MMP-13 to wild type;

Figure 43A summarizes the cloning strategy used to generate the pAP-288 construct;

Figure 43B shows the nucleotide sequence of the Tissue-type Plasminogen Activator linker region of pAP-288;

5 Figure 43C shows the DNA sequence of the pAP-288 insert;

Figure 43D shows the amino acid sequence comparison of mutant preproricin linker region of Tissue-type Plasminogen Activator to wild type;

10 Figure 44A summarizes the cloning strategy used to generate the pAP-290 construct;

Figure 44B shows the nucleotide sequence of the human Prostate-Specific Antigen linker region of pAP-290;

15 Figure 44C shows the DNA sequence of the pAP-290 insert;

Figure 44D shows the amino acid sequence comparison of mutant preproricin linker region of the human Prostate-Specific Antigen to wild type;

20 Figure 45A summarizes the cloning strategy used to generate the pAP-292 construct;

Figure 45B shows the nucleotide sequence of the kallikrein linker region of pAP-292;

Figure 45C shows the DNA sequence of the pAP-292 insert;

25 Figure 45D shows the amino acid sequence comparison of mutant preproricin linker region of the kallikrein to wild type;

Figure 46A summarizes the cloning strategy used to generate the pAP-294 construct;

30 Figure 46B shows the nucleotide sequence of the neutrophil elastase linker region of pAP-294;

Figure 46C shows the DNA sequence of the pAP-294 insert;

- 28 -

Figure 46D shows the amino acid sequence comparison of mutant preproricin linker region of neutrophil elastase to wild type;

Figure 47A summarizes the cloning strategy used to generate the pAP-296 construct;

5 Figure 47B shows the nucleotide sequence of the calpain linker region of pAP-296;

Figure 47C shows the DNA sequence of the pAP-296 insert;

10 Figure 47D shows the amino acid sequence comparison of mutant preproricin linker region of calpain to wild type;

Figure 48 is a blot showing cleavage of pAP-214 by Cathepsin B;

Figure 49 is a blot showing cleavage of pAP-220 with MMP-9;

15 Figure 50 is a blot showing activation of pAP-214; and Figure 51 is a blot showing activation of pAP-220.

Figure 52 is a blot showing cleavage of pAP-248 with HCMV.

20 Figure 53 is a blot showing activation of pAP-248.

Figure 54 is a blot showing cleavage of pAP-256 by HAV 3C.

Figure 55 is a blot showing activation of pAP-256.

25 Figure 56 is a semi-logithmic graph illustrating the cytotoxicity to COS-1 cells of undigested pAP-214 and pAP-214 digested with Cathepsin B.

Figure 57 is a semi-logithmic graph illustrating the cytotoxicity of pAP-220 digested with MMP-9 compared to freshly thawed pAP-220 and ricin on COS-1 cells.

30 Figure 58 is a blot showing cleavage of pAP-270 with MMP-2.

Figure 59 is a blot showing activation of pAP-270.

Figure 60 is a blot showing cleavage of pAP-288 by t-PA.

- 29 -

Figure 61 is a blot showing activation of pAP-288.

Figure 62 is a blot showing cleavage of pAP-294 with human neutrophil elastase.

Figure 63 is a blot showing activation of pAP-294.

5 Figure 64 is a blot showing cleavage of pAP-296 with calpain.

Figure 65 is a blot showing activation of pAP-296.

Figure 66 is a blot showing cleavage of pAP-222 with MMP-2.

10 Figure 67 is a blot showing activation of pAP-222.

DETAILED DESCRIPTION OF THE INVENTION

Nucleic Acid Molecules of the Invention

As mentioned above, the present invention relates to novel nucleic acid molecules comprising a nucleotide sequence
15 encoding an A chain of a ricin-like toxin, a B chain of a ricin-like toxin and a heterologous linker amino acid sequence, linking the A and B chains. The heterologous linker sequence contains a cleavage recognition site for a disease-specific protease (e.g. a viral protease, parasitic protease, cancer-associated protease, or a fungal protease).

20 The term "isolated and purified" as used herein refers to a nucleic acid substantially free of cellular material or culture medium when produced by recombinant DNA techniques, or chemical precursors, or other chemicals when chemically synthesized. An "isolated and purified" nucleic acid is also substantially free of
25 sequences which naturally flank the nucleic acid (*i.e.* sequences located at the 5' and 3' ends of the nucleic acid) from which the nucleic acid is derived. The term "nucleic acid" is intended to include DNA and RNA and can be either double stranded or single stranded.

The term "linker sequence" as used herein refers to an
30 internal amino acid sequence within the protein encoded by the nucleic acid molecule of the invention which contains residues linking the A and B chain so as to render the A chain incapable of exerting its toxic

- 30 -

effect, for example catalytically inhibiting translation of a eukaryotic ribosome. By heterologous is meant that the linker sequence is not a sequence native to the A or B chain of a ricin-like toxin or precursor thereof. However, preferably, the linker sequence may be of a similar
5 length to the linker sequence of a ricin-like toxin and should not interfere with the role of the B chain in cell binding and transport into the cytoplasm. When the linker sequence is cleaved the A chain becomes active or toxic.

The nucleic acid molecule of the invention is cloned by
10 subjecting a preproricin cDNA clone to site-directed mutagenesis in order to generate a series of variants differing only in the sequence between the A and B chains (linker region). Oligonucleotides, corresponding to the extreme 5' and 3' ends of the preproricin gene are synthesized and used to PCR amplify the gene. Using the cDNA
15 sequence for preproricin (Lamb et al., *Eur. J. Biochem.* 145:266-270 (1985)), several oligonucleotide primers are designed to flank the start and stop codons of the preproricin open reading frame.

The preproricin cDNA is amplified using the upstream primer Ricin-99 or Ricin-109 and the downstream primer Ricin1729C
20 with Vent DNA polymerase (New England Biolabs) using standard procedures (Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, (Cold Spring Harbor Laboratory Press, 1989)). The purified PCR fragment encoding the preproricin cDNA is then ligated into an Eco RI-digested pBluescript II SK plasmid (Stratagene), and is
25 used to transform competent XL1-Blue cells (Stratagene). The cloned PCR product containing the putative preproricin gene is confirmed by DNA sequencing of the entire cDNA clone. The sequences and location of oligonucleotide primers used for sequencing are shown in Table 1.

30 The preproricin cDNA clone is subjected to site directed mutagenesis in order to generate a series of variants differing only in the sequence between the A and B chains (linker region). The wild-type

- 31 -

preproricin linker region is replaced with the heterogenous linker sequences that are cleaved by the various disease-specific proteases as shown in Figures 21, 26, 27, 28, and Part D of Figures 30-47. Linker identification as used herein in connection with the sequences provided
5 in these figures have been assigned the sequence ID numbers as discussed below.

The linker regions of the variants encode a cleavage recognition sequence for a disease-specific protease associated with for example, cancer, viruses, parasites, or fungi. The mutagenesis and
10 cloning strategy used to generate the disease-specific protease-sensitive linker variants are summarized in Part A of Figures 2-20, and Part A of Figures 22-25. The first step involves a DNA amplification using a set of mutagenic primers in combination with the two flanking primers Richin-99Eco or Ricin-109Eco and Ricin1729C Pst I. Restriction digested
15 PCR fragments are gel purified and then ligated with PBluescript SK which has been digested with Eco RI and Pst I. Ligation reactions are used to transform competent XL1-Blue cells (Stratagene). Recombinant clones are identified by restriction digests of plasmid miniprep DNA and the mutant linker sequences are confirmed by DNA sequencing.
20 With respect to the nucleotide sequences and amino acid sequences prepared as a result of the implementation of this strategy the following sequences have been assigned the sequence ID numbers as indicated.

SEQ ID NO. 1 is used herein in connection with the DNA sequence of the baculovirus transfer vector, pVL1393.

25 The nucleotide sequence of Cathepsin B linker regions of pAP-213 are referred to herein as SEQ ID NO. 2.

The nucleotide sequence of Cathepsin B linker regions of pAP-214 are referred to herein as SEQ ID NO. 3.

The nucleotide sequence of MMP-3 linker regions of pAP-
30 215 are referred to herein as SEQ ID NO. 4.

The DNA sequence of the pAP-216 insert containing ricin and the MMP-3 linker are referred to herein as SEQ ID NO. 5.

The nucleotide sequence of MMP-7 linker regions of pAP-217 are referred to herein as SEQ ID NO. 6.

The DNA sequence of the pAP-218 insert containing ricin and the MMP-7 linker are referred to herein as SEQ ID NO. 7.

5 The nucleotide sequence of MMP-9 linker regions of pAP-219 are referred to herein as SEQ ID NO. 8.

The DNA sequence of the pAP-220 insert containing ricin and the MMP-9 are referred to herein as SEQ ID NO. 9.

10 The nucleotide sequence of thermolysin-like MMP linker regions of pAP-221 are referred to herein as SEQ ID NO. 10.

The DNA sequence of of pAP-222 insert containing ricin and the thermolysin-like MMP linker are referred to herein as SEQ ID NO. 11.

15 The nucleotide sequence of Plasmodium falciparum-A linker regions of pAP-223 are referred to herein as SEQ ID NO. 12.

The DNA sequence of the pAP-224 insert containing ricin and the Plasmodium falciparum-A linker are referred to herein as SEQ ID NO. 13.

20 The nucleotide sequence of Plasmodium falciparum-B linker regions of pAP-225 are referred to herein as SEQ ID NO. 14.

The DNA sequence of the pAP-226 insert containing ricin and the Plasmodium falciparum-B linker are referred to herein as SEQ ID NO. 15.

25 The nucleotide sequence of Plasmodium falciparum-C linker regions of pAP-227 are referred to herein as SEQ ID NO. 16.

The DNA sequence of the pAP-228 insert containing ricin and the Plasmodium falciparum-C linker are referred to herein as SEQ ID NO. 17.

30 The nucleotide sequence of the the Plasmodium falciparum-D linker regions of pAP-229 is referred to herein as SEQ ID NO. 18.

The DNA sequence of the pAP-230 insert containing ricin and the Plasmodium falciparum-D linker is referred to herein as SEQ ID NO. 19.

The nucleotide sequence of the Plasmodium falciparum-E linker regions of pAP-231 is referred to herein as SEQ ID NO. 20.

The DNA sequence of the pAP-232 insert containing ricin and the Plasmodium falciparum-E linker is referred to herein as SEQ ID NO. 21.

The nucleotide sequence of the HSV-A linker regions of pAP-233 is referred to herein as SEQ ID NO. 22.

The DNA sequence of the pAP-234 insert containing ricin and the HSV-A linker is referred to herein as SEQ ID NO. 23.

The nucleotide sequence of the HSV-B linker regions of pAP-235 is referred to herein as SEQ ID NO. 24.

The DNA sequence of the pAP-236 insert containing ricin and the HSV-B linker is referred to herein as SEQ ID NO. 25.

The nucleotide sequence of the VZV-A linker regions of pAP-237 are referred to herein as SEQ ID NO. 26.

The DNA sequence of the pAP-238 insert containing ricin and the VZV-A linker are referred to herein as SEQ ID NO. 27.

The nucleotide sequence of the VZV-B linker regions of PAP-239 is referred to herein as SEQ ID NO. 28.

The DNA sequence of the pAP-240 insert containing ricin and the VZV-B linker is referred to herein as SEQ ID NO. 29.

The nucleotide sequence of the EBV-A linker regions of pAP-241 is referred to herein as SEQ ID NO. 30.

The DNA sequence of the pAP-242 insert containing ricin and the EBV-A linker is referred to herein as SEQ ID NO. 31.

The nucleotide sequence of the EBV-B linker regions of pAP-243 is referred to herein as SEQ ID NO. 32.

The DNA sequence of the pAP-244 insert containing ricin and the EBV-B linker is referred to herein as SEQ ID NO. 33.

- 34 -

The nucleotide sequence of the CMV-A linker regions of pAP-245 is referred to herein as SEQ ID NO. 34.

The DNA sequence of the pAP-246 insert containing ricin and the CMV-A linker is referred to herein as SEQ ID NO. 35.

5 The nucleotide sequence of the CMV-B linker regions of pAP-247 is referred to herein as SEQ ID NO. 36.

The DNA sequence of the pAP-248 insert containing ricin and the CMV-B linker is referred to herein as SEQ ID NO. 37.

10 The nucleotide sequence of the HHV-6 linker regions of pAP-249 is referred to herein as SEQ ID NO. 38.

The DNA sequence of the pAP-250 insert containing ricin and the HHV-6 linker is referred to herein as SEQ ID NO. 39.

15 The amino acid sequences of the cancer protease-sensitive amino acid linkers contained in the following pAP proteins have the sequence ID numbers as indicated: pAP-213 and pAP-214 (SEQ ID NO. 40); pAP-215 and pAP-216 (SEQ ID NO. 41); pAP-217 and pAP-218; (SEQ ID NO. 42); pAP-219 and pAP-220 (SEQ ID NO. 43); and pAP-221 and pAP-222 (SEQ ID NO. 44).

20 The amino acid sequences of the following cancer protease-sensitive linkers are referred to herein with the corresponding sequence ID numbers: pAP-241 and pAP-242 (SEQ ID NO. 45); and pAP-243 and pAP-244 (SEQ ID NO. 46).

The nucleotide sequence of the ILV linker regions of pAP-253 is referred to herein as SEQ ID NO. 47.

25 The DNA sequence of the pAP-254 insert containing ricin and the ILV linker is referred to herein as SEQ ID NO. 48.

The nucleotide sequence of the HAV-A linker regions of pAP-257 is referred to herein as SEQ ID NO. 49.

30 The DNA sequence of the pAP-258 insert containing ricin and HAV-A linker is referred to herein as SEQ ID NO. 50.

The nucleotide sequence of the HAV-B linker regions of pAP-255 is referred to herein as SEQ ID NO. 51.

The DNA sequence of the pAP-256 insert containing ricin and the HAV-B linker is referred to herein as SEQ ID NO. 52.

The nucleotide sequence of the CAN linker regions of pAP-259 is referred to herein as SEQ ID NO. 53.

5 The DNA sequence of the pAP-260 insert containing ricin and the CAN linker is referred to herein as SEQ ID NO. 54.

 The amino acid sequences of *Plasmodium falciparum* protease-sensitive linkers are referred to herein by the sequence ID numbers as follows: pAP-223 and pAP-224 (SEQ ID NO 55); pAP-225 and
10 pAP-226 (SEQ ID NO 56); pAP-227 and pAP-228 (SEQ ID NO 57); pAP-229 and pAP-230 (SEQ ID NO 58); and pAP-231 and pAP-232 (SEQ ID NO 59) (see Figure 26).

 The amino acid sequences of the viral protease-sensitive linkers which follow are referred to herein by the sequence ID numbers
15 indicated: pAP-233 and pAP 234 (SEQ ID NO 60); pAP-235 and pAP-236 (SEQ ID NO 61); and pAP-249 and pAP-250 (SEQ ID NO 62) (see Figure 27).

 The amino acid sequences of the viral protease-sensitive linkers which follow are referred to herein by the sequence ID numbers
20 indicated: pAP-245 and pAP-246 (SEQ ID NO 63) ; and pAP-247 and pAP-248 (SEQ ID NO 64) (see Figure 27).

 The amino acid sequences of the viral protease-sensitive linkers which follow are referred to herein by the sequence ID numbers indicated: pAP-237 and pAP-238 (SEQ ID NO 65); and pAP-239 and pAP-
25 240 (SEQ ID NO 66); pAP-253 and pAP-254 (SEQ ID NO 67); pAP-255 and pAP-256 (SEQ ID NO 68); and pAP-257 and pAP-258 (SEQ ID NO 69) (see Figure 27).

 The amino acid sequences of the *Candida* aspartic protease-sensitive linkers are referred to herein by the sequence ID
30 numbers indicated: pAP-259 and pAP-260 (SEQ ID NO 70); pAP-261 and pAP-262 (SEQ ID NO 71); and pAP-263 and pAP-264 (SEQ ID NO 72).

An alternative mutagenesis and cloning strategy that can be used to generate the disease-specific protease-sensitive linker variants is summarized in Figure 29. The first step of this method involves a DNA amplification using a set of mutagenic primers in combination with the two flanking primers Ricin-109Eco and Ricin1729Pst. Restriction digested PCR fragments (Eco RI and Pst I) are gel purified. Preproricin variants produced from this method can be subcloned directly into the baculovirus transfer vector digested with Eco RI and Pst I and intermediate ligation steps involving pBluescript SK and pSB2 are circumvented. The cloning strategies used to generate disease-specific protease-sensitive linker variants are summarized in Part A of Figures 30 to 47. With respect to the nucleotide sequences and amino acid sequences prepared as a result of the implementation of this strategy the following sequences have been assigned the sequence ID numbers as indicated.

The nucleotide sequence of the HCV-A linker region of pAP-262 is referred to herein as SEQ ID NO. 73.

The DNA sequence of the pAP-262 insert is referred to herein as SEQ ID NO. 74.

The amino acid sequence of the mutant preproricin linker region for HCV-A, pAP-262, is referred to herein as SEQ ID NO. 75.

The nucleotide sequence of the HCV-B linker region of pAP-264 is referred to herein as SEQ ID NO. 76.

The DNA sequence of the pAP-264 insert is referred to herein as SEQ ID NO. 77.

The amino acid sequence of the mutant preproricin linker region for HCV-B, pAP-264, is referred to herein as SEQ ID NO. 78.

The nucleotide sequence of the HCV-C linker region of pAP-266 is referred to herein as SEQ ID NO. 79.

The DNA sequence of the pAP-266 insert is referred to herein as SEQ ID NO. 80.

The amino acid sequence of the mutant preprorin linker region for HCV-C, pAP-266, is referred to herein as SEQ ID NO. 81.

The nucleotide sequence of the HCV-D linker region of pAP-268 is referred to herein as SEQ ID NO. 82.

The DNA sequence of the pAP-268 insert is referred to herein as SEQ ID NO. 83.

The amino acid sequence of the mutant preprorin linker region for HCV-D, pAP-268, is referred to herein as SEQ ID NO. 84.

The nucleotide sequence of the MMP-2 linker region of pAP-270 is referred to herein as SEQ ID NO. 85.

The DNA sequence of the pAP-270 insert is referred to herein as SEQ ID NO. 86.

The amino acid sequence of the mutant preprorin linker region for MMP-2, pAP-270, is referred to herein as SEQ ID NO. 87.

The nucleotide acid sequence of the Cathepsin B (Site 2) linker region of pAP-272 is referred to herein as SEQ ID NO. 88.

The DNA sequence of the pAP-272 insert is referred to herein as SEQ ID NO. 89.

The amino acid sequence of the mutant preprorin linker region for Cathepsin B (Site 2), pAP-272, is referred to herein as SEQ ID NO. 90.

The nucleotide sequence of the Cathepsin L linker region of pAP-274 is referred to herein as SEQ ID NO. 91.

The DNA sequence of the pAP-274 insert is referred to herein as SEQ ID NO. 92.

- 38 -

The amino acid sequence of the mutant preproricin linker region of Cathepsin L, pAP-274, is referred to herein as SEQ ID NO. 93.

5 The nucleotide sequence of Cathepsin D linker region of pAP-276 is referred to herein as SEQ ID NO. 94.

The DNA sequence of the pAP-276 insert is referred to herein as SEQ ID NO. 95.

10 The amino acid sequence of the mutant preproricin linker region for Cathepsin D, pAP-276, is referred to herein as SEQ ID NO. 96.

The nucleotide sequence of the MMP-1 linker region of pAP-278 is referred to herein as SEQ ID NO. 97.

The DNA sequence of the pAP-278 insert is referred to herein as SEQ ID NO. 98.

15 The amino acid sequence of the mutant preproricin linker region for MMP-1, pAP-278, is referred to herein as SEQ ID NO. 99.

20 The nucleotide sequence of the Urokinase-Type Plasminogen Activator linker region of pAP-280 is referred to herein as SEQ ID NO. 100.

The DNA sequene of the pAP-280 insert is referred to herein as SEQ ID NO. 101.

25 The amino acid sequence of the mutant preproricin linker region for Urokinase-Type Plasminogen Activator, pAP-280, is referred to herein as SEQ ID NO. 102.

The nucleotide sequence of MT-MMP linker region of pAP-282 is referred to herein as SEQ ID NO. 103.

The DNA sequence of the pAP-282 insert is referred to herein as SEQ ID NO. 104.

30 The amino acid sequence of the mutant preproricin linker region for MT-MMP, pAP-282, is referred to herein as SEQ ID NO. 105.

The nucleotide sequence of the MMP-11 linker region of pAP-284 is referred to herein as SEQ ID NO. 106.

The DNA sequence of the pAP-284 insert is referred to herein as SEQ ID NO. 107.

5 The amino acid sequence of the mutant preprorcin linker region for MMP-11, pAP-284, is referred to herein as SEQ ID NO. 108.

The nucleotide sequence of the MMP-13 linker region of pAP-286 is referred to herein as SEQ ID NO. 109.

10 The DNA sequence of the pAP-286 insert is referred to herein as SEQ ID NO. 110.

The amino acid sequence of the mutant preprorcin linker region for MMP-13, pAP-286, is referred to herein as SEQ ID NO. 111.

15 The nucleotide sequence of the Tissue-type Plasminogen Activator linker region of pAP-288 is referred to herein as SEQ ID NO. 112.

The DNA sequence of the pAP-288 insert is referred to herein as SEQ ID NO. 113.

20 The amino acid sequence of the mutant preprorcin linker region for Tissue-type Plasminogen Activator, pAP-288, is referred to herein as SEQ ID NO. 114.

The nucleotide sequence of the human Prostate-Specific Antigen linker region of pAP-290 is referred to herein as SEQ ID NO.

25 115.

The DNA sequence of the pAP-290 insert is referred to herein as SEQ ID NO. 116.

The amino acid sequence of the mutant preprorcin linker region for the human Prostate-Specific Antigen, pAP-290, is referred to herein as SEQ ID NO. 117.

30

The nucleotide sequence of the kallikrein linker region of pAP-292 is referred to herein as SEQ ID NO. 118.

- 40 -

The DNA sequence of the pAP-292 insert is referred to herein as SEQ ID NO. 119.

The amino acid sequence of the mutant preproricin linker region for the kallikrein, pAP-292, is referred to herein as SEQ ID
5 NO. 120.

The nucleotide sequence of the neutrophil elastase linker region of pAP-294 is referred to herein as SEQ ID NO. 121.

The DNA sequence of the pAP-294 insert is referred to herein as SEQ ID NO. 122.

10 The amino acid sequence of the mutant preproricin linker region for neutrophil elastase, pAP-294, is referred to herein as SEQ ID NO. 123.

The nucleotide sequence of the calpain linker region of pAP-296 is referred to herein as SEQ ID NO. 124.

15 The DNA sequence of the pAP-296 insert is referred to herein as SEQ ID NO. 125.

The amino acid sequence of the mutant preproricin linker region for calpain, pAP-296, is referred to herein as SEQ ID NO.
126.

20 The amino acid sequence of the wild type linker region is referred to herein as SEQ ID NO. 127.

The nucleic acid molecule of the invention has sequences encoding an A chain of a ricin-like toxin, a B chain of a ricin-like toxin and a heterologous linker sequence containing a cleavage recognition
25 site for a disease-specific protease. The nucleic acid may be expressed to provide a recombinant protein having an A chain of a ricin-like toxin, a B chain of a ricin-like toxin and a heterologous linker sequence containing a cleavage recognition site for a disease-specific protease.

30 The nucleic acid molecule may comprise the A and/or B chain of ricin. The ricin gene has been cloned and sequenced, and the X-ray crystal structures of the A and B chains are published (Rutenber, E., et al. *Proteins* 10:240-250 (1991); Weston et al., *Mol. Biol.* 244:410-422

(1994); Lamb and Lord, *Eur. J. Biochem.* 14:265 (1985); Halling, K., et al., *Nucleic Acids Res.* 13:8019 (1985)). It will be appreciated that the invention includes nucleic acid molecules encoding truncations of A and B chains of ricin like proteins and analogs and homologs of A and B chains of ricin-like proteins and truncations thereof (i.e., ricin-like proteins), as described herein. It will further be appreciated that variant forms of the nucleic acid molecules of the invention which arise by alternative splicing of an mRNA corresponding to a cDNA of the invention are encompassed by the invention.

Another aspect of the invention provides a nucleotide sequence which hybridizes under high stringency conditions to a nucleotide sequence encoding the A and/or B chains of a ricin-like protein. Appropriate stringency conditions which promote DNA hybridization are known to those skilled in the art, or can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1 6.3.6. For example, 6.0 x sodium chloride/sodium citrate (SSC) at about 45°C, followed by a wash of 2.0 x SSC at 50°C may be employed. The stringency may be selected based on the conditions used in the wash step. By way of example, the salt concentration in the wash step can be selected from a high stringency of about 0.2 x SSC at 50°C. In addition, the temperature in the wash step can be at high stringency conditions, at about 65°C.

The nucleic acid molecule may comprise the A and/or B chain of a ricin-like toxin. Methods for cloning ricin-like toxins are known in the art and are described, for example, in E.P. 466,222. Sequences encoding ricin or ricin-like A and B chains may be obtained by selective amplification of a coding region, using sets of degenerative primers or probes for selectively amplifying the coding region in a genomic or cDNA library. Appropriate primers may be selected from the nucleic acid sequence of A and B chains of ricin or ricin-like toxins. It is also possible to design synthetic oligonucleotide primers from the nucleotide sequences for use in PCR. Suitable primers may be selected

- 42 -

from the sequences encoding regions of ricin-like proteins which are highly conserved, as described for example in U.S. Patent No 5,101,025 and E.P. 466,222.

A nucleic acid can be amplified from cDNA or genomic
5 DNA using these oligonucleotide primers and standard PCR
amplification techniques. The nucleic acid so amplified can be cloned
into an appropriate vector and characterized by DNA sequence analysis.
It will be appreciated that cDNA may be prepared from mRNA, by
isolating total cellular mRNA by a variety of techniques, for example, by
10 using the guanidinium-thiocyanate extraction procedure of Chirgwin et
al., *Biochemistry* 18, 5294-5299 (1979). cDNA is then synthesized from
the mRNA using reverse transcriptase (for example, Moloney MLV
reverse transcriptase available from Gibco/BRL, Bethesda, MD, or AMV
reverse transcriptase available from Seikagaku America, Inc., St.
15 Petersburg, FL). It will be appreciated that the methods described above
may be used to obtain the coding sequence from plants, bacteria or
fungi, preferably plants, which produce known ricin-like proteins and
also to screen for the presence of genes encoding as yet unknown
ricin-like proteins.

20 A sequence containing a cleavage recognition site for a
specific protease may be selected based on the disease or the pathogen
which is to be targeted by the recombinant protein. The cleavage
recognition site may be selected from sequences known to encode a
cleavage recognition site for the cancer, viral or parasitic protease.
25 Sequences encoding cleavage recognition sites may be identified by
testing the expression product of the sequence for susceptibility to
cleavage by the respective protease.

A sequence containing a cleavage recognition site for a
viral, fungal, parasitic or cancer associated protease may be selected
30 based on the retrovirus which is to be targeted by the recombinant
protein. The cleavage recognition site may be selected from sequences
known to encode a cleavage recognition site for the viral, fungal,

parasitic or cancer associated protease. Sequences encoding cleavage recognition sites may be identified by testing the expression product of the sequence for susceptibility to cleavage by a viral, fungal, parasitic or cancer associated protease. A polypeptide containing the suspected
5 cleavage recognition site may be incubated with a protease and the amount of cleavage product determined (DiIannit, 1990, J. Biol. Chem. 285: 17345-17354 (1990)).

The protease may be prepared by methods known in the art and used to test suspected cleavage recognition sites.

10 In one embodiment, the preparation of tumour-associated cathepsin B, its substrates and enzymatic activity assay methodology have been described by Sloane, B.F. et al. (*Proc. Natl. Acad. Sci. USA* 83:2483-2487 (1986)), Schwartz, M.K. (*Clin. Chim. Acta* 237:67-78 (1995)), and Panchal, R.G. et al. (*Nature Biotechnol.* 14:852-856 (1996)).
15 The preparation of Epstein-Barr virus protease, its substrates and enzymatic activity assay methodology have been described by Welch, A.R. (*Proc. Natl. Acad. Sci. USA* 88:10792-10796 (1991)).

In another embodiment, the preparation of *Plasmodium falciparum* proteases, their substrates and enzymatic activity assay
20 methodology have been described by Goldberg, D.E. et al. (*J. Exp. Med.* 173:961-969 (1991)), Cooper & Bujard (*Mol. Biochem. Parasitol.* 56:151-160 (1992)), Nwagwu, M. et al. (*Exp. Parasitol.* 75:399-414 (1992)), Rosenthal, P.J. et al. (*J. Clin. Invest.* 91:1052-1056 (1993)), Blackman, M.J. et al. (*Mol. Biochem. Parasitol.* 62:103-114 (1995)).

25 In a further embodiment, the preparation of proteases from human cytomegalovirus, human herpes virus, varicella zoster virus and infectious laryngotracheitis virus have been taught by Liu F. & Roizman, B. (*J. Virol.* 65:5149-5156 (1991)) and Welch, A.R. (*Proc. Natl. Acad. Sci. USA* 88:10792-10796 (1991)). In addition, their respective
30 substrates and enzymatic activity assay methodologies are also described.

In another embodiment, the preparation of hepatitis A virus protease, its substrates and enzymatic activity assay methodology have been described by Jewell, D.A. et al. (*Biochemistry* 31:7862-7869 (1992)). The preparation of poliovirus protease, its substrates and enzymatic activity assay methodology have been described by Weidner, J.R. et al. (*Arch. Biochem. Biophys.* 286:402-408 (1991)). The preparation of human rhinovirus protease, its substrates and enzymatic activity assay methodology have been described by Long, A.C. et al. (*FEBS Lett.* 258:75-78 (1989)).

10 In another embodiment of the invention, the preparation of proteases associated with *Candida* yeasts their substrates and enzymatic activity are contemplated, including the aspartic proteinases which have been associated specifically with numerous virulent strains of *Candida* including *Candida albican*, *Candida tropicalis*, and *Candida*
15 *parapsilosis* (Abad-Zapatero, C. et al., *Protein Sci.* 5:640-652 (1996); Cutfield, S.M. et al., *Biochemistry* 35:398-410 (1995); Ruchel, R. et al, *Zentralbl. Bakteriol. Mikrobiol Hyg. I Abt. Orig. A.* 255:537-548 (1983); Remold, H. et al., *Biochim. Biophys. Acta* 167:399-406 (1968)).

The nucleic acid molecule of the invention may be
20 prepared by site directed mutagenesis. For example, the cleavage site of a disease-specific protease may be prepared by site directed mutagenesis of the homologous linker sequence of a proricin-like toxin. Procedures for cloning proricin-like genes, encoding a linker sequence are described in EP 466,222. Site directed mutagenesis may be accomplished by DNA
25 amplification of mutagenic primers in combination with flanking primers. Suitable procedures using the mutagenic primers are shown in Parts A and B of Figures 1-4, Figures 13-16, Figures 18-36, Figures 38-41, and Figures 50-67.

The nucleic acid molecule of the invention may also
30 encode a fusion protein. A sequence encoding a heterologous linker sequence containing a cleavage recognition site for a disease-specific protease may be cloned from a cDNA or genomic library or chemically

synthesized based on the known sequence of such cleavage sites. The heterologous linker sequence may then be fused in frame with the sequences encoding the A and B chains of the ricin-like toxin for expression as a fusion protein. It will be appreciated that a nucleic acid molecule encoding a fusion protein may contain a sequence encoding an A chain and a B chain from the same ricin-like toxin or the encoded A and B chains may be from different toxins. For example, the A chain may be derived from ricin and the B chain may be derived from abrin. A protein may also be prepared by chemical conjugation of the A and B chains and linker sequence using conventional coupling agents for covalent attachment.

An isolated and purified nucleic acid molecule of the invention which is RNA can be isolated by cloning a cDNA encoding an A and B chain and a linker into an appropriate vector which allows for transcription of the cDNA to produce an RNA molecule which encodes a protein of the invention. For example, a cDNA can be cloned downstream of a bacteriophage promoter, (e.g. a T7 promoter) in a vector, cDNA can be transcribed in vitro with T7 polymerase, and the resultant RNA can be isolated by standard techniques.

20 Recombinant Protein of the Invention

As previously mentioned, the invention provides novel recombinant proteins which incorporate the A and B chains of a ricin like toxin linked by a heterologous linker sequence containing a cleavage recognition site for a disease-specific protease. It is an advantage of the recombinant proteins of the invention that they are non-toxic until the A chain is liberated from the B chain by specific cleavage of the linker by the target protease.

Thus the protein may be used to specifically target cancer cells or cells infected with a virus or parasite in the absence of additional specific cell-binding components to target infected cells. It is a further advantage that the disease-specific protease cleaves the heterologous linker intracellularly thereby releasing the toxic A chain directly into

the cytoplasm of the cancer cell or infected cell. As a result, said cells are specifically targeted and non-infected normal cells are not directly exposed to the activated free A chain.

Ricin is a plant derived ribosome inhibiting protein which blocks protein synthesis in eukaryotic cells. Ricin may be derived from the seeds of *Ricinus communis* (castor oil plant). The ricin toxin is a glycosylated heterodimer with A and B chain molecular masses of 30,625 Da and 31,431 Da respectively. The A chain of ricin has an N-glycosidase activity and catalyzes the excision of a specific adenine residue from the 28S rRNA of eukaryotic ribosomes (Endo, Y; & Tsurugi, K. J. Biol. Chem. 262:8128 (1987)). The B chain of ricin, although not toxic in itself, promotes the toxicity of the A chain by binding to galactose residues on the surface of eukaryotic cells and stimulating receptor-mediated endocytosis of the toxin molecule (Simmons et al., Biol. Chem. 261:7912 (1986)).

All protein toxins are initially produced in an inactive, precursor form. Ricin is initially produced as a single polypeptide (preproricin) with a 35 amino acid N-terminal presequence and 12 amino acid linker between the A and B chains. The pre-sequence is removed during translocation of the ricin precursor into the endoplasmic reticulum (Lord, J.M., Eur. J. Biochem. 146:403-409 (1985) and Lord, J.M., Eur. J. Biochem. 146:411-416 (1985)). The proricin is then translocated into specialized organelles called protein bodies where a plant protease cleaves the protein at a linker region between the A and B chains (Lord, J.M. et al., FASAB Journal 8:201-208 (1994)). The two chains, however, remain covalently attached by an interchain disulfide bond (cysteine 259 in the A chain to cysteine 4 in the B chain) and mature disulfide linked ricin is stored in protein bodies inside plant cells. The A chain is inactive in the proricin (O'Hare, M., et al., FEBS Lett. 273:200-204 (1990)) and it is inactive in the disulfide-linked mature ricin (Richardson, P.T. et al., FEBS Lett. 255:15-20 (1989)). The ribosomes of the castor bean plant are themselves susceptible to inactivation by

- 47 -

ricin A chain; however, as there is no cell surface galactose to permit B chain recognition the A chain cannot re-enter the cell.

Ricin-like proteins include, but are not limited to, bacterial, fungal and plant toxins which have A and B chains and inactivate ribosomes and inhibit protein synthesis. The A chain is an active polypeptide subunit which is responsible for the pharmacologic effect of the toxin. In most cases the active component of the A chain is an enzyme. The B chain is responsible for binding the toxin to the cell surface and is thought to facilitate entry of the A chain into the cell cytoplasm. The A and B chains in the mature toxins are linked by disulfide bonds. The toxins most similar in structure to ricin are plant toxins which have one A chain and one B chain. Examples of such toxins include abrin which may be isolated from the seeds of *Abrus precatorius* and modeccin.

Ricin-like bacterial proteins include diphtheria toxin, which is produced by *Corynebacterium diphtheriae*, *Pseudomonas enterotoxin A* and cholera toxin. It will be appreciated that the term ricin-like toxins is also intended to include the A chain of those toxins which have only an A chain. The recombinant proteins of the invention could include the A chain of these toxins conjugated to, or expressed as, a recombinant protein with the B chain of another toxin. Examples of plant toxins having only an A chain include trichosanthin, MMC and pokeweed antiviral proteins, dianthin 30, dianthin 32, croton II, curcin II and wheat germ inhibitor. Examples of fungal toxins having only an A chain include alpha-sarcin, restrictocin, mitogillin, enomycin, phenomycin. Examples of bacterial toxins having only an A chain include cytotoxin from *Shigella dysenteriae* and related Shiga-like toxins. Recombinant trichosanthin and the coding sequence thereof is disclosed in U.S. Patents 5,101,025 and 5,128,460.

In addition to the entire A or B chains of a ricin-like toxin, it will be appreciated that the recombinant protein of the invention may contain only that portion of the A chain which is

necessary for exerting its cytotoxic effect. For example, the first 30 amino acids of the ricin A chain may be removed resulting in a truncated A chain which retains toxic activity. The truncated ricin or ricin-like A chain may be prepared by expression of a truncated gene or by
5 proteolytic degradation, for example with Nagarase (Funmatsu et al., *Jap. J. Med. Sci. Biol.* 23:264-267 (1970)). Similarly, the recombinant protein of the invention may contain only that portion of the B chain necessary for galactose recognition, cell binding and transport into the cell cytoplasm. Truncated B chains are described for example in E.P.
10 145,111. The A and B chains may be glycosylated or non-glycosylated. Glycosylated A and B chains may be obtained by expression in the appropriate host cell capable of glycosylation. Non-glycosylated chains may be obtained by expression in nonglycosylating host cells or by treatment to remove or destroy the carbohydrate moieties.

15 The proteins of the invention may be prepared using recombinant DNA methods. Accordingly, the nucleic acid molecules of the present invention may be incorporated in a known manner into an appropriate expression vector which ensures good expression of the protein. Possible expression vectors include but are not limited to
20 cosmids, plasmids, or modified viruses (e.g. replication defective retroviruses, adenoviruses and adeno-associated viruses), so long as the vector is compatible with the host cell used. The expression vectors are "suitable for transformation of a host cell", which means that the expression vectors contain a nucleic acid molecule of the invention and
25 regulatory sequences selected on the basis of the host cells to be used for expression, which is operatively linked to the nucleic acid molecule. Operatively linked is intended to mean that the nucleic acid is linked to regulatory sequences in a manner which allows expression of the nucleic acid.

30 The invention therefore contemplates a recombinant expression vector of the invention containing a nucleic acid molecule of the invention, or a fragment thereof, and the necessary regulatory

sequences for the transcription and translation of the inserted protein-sequence.

Suitable regulatory sequences may be derived from a variety of sources, including bacterial, fungal, viral, mammalian, or insect genes (For example, see the regulatory sequences described in Goeddel, Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, CA (1990). Selection of appropriate regulatory sequences is dependent on the host cell chosen as discussed below, and may be readily accomplished by one of ordinary skill in the art. Examples of such regulatory sequences include: a transcriptional promoter and enhancer or RNA polymerase binding sequence, a ribosomal binding sequence, including a translation initiation signal. Additionally, depending on the host cell chosen and the vector employed, other sequences, such as an origin of replication, additional DNA restriction sites, enhancers, and sequences conferring inducibility of transcription may be incorporated into the expression vector. It will also be appreciated that the necessary regulatory sequences may be supplied by the native A and B chains and/or its flanking regions.

The recombinant expression vectors of the invention may also contain a selectable marker gene which facilitates the selection of host cells transformed or transfected with a recombinant molecule of the invention. Examples of selectable marker genes are genes encoding a protein such as G418 and hygromycin which confer resistance to certain drugs, β -galactosidase, chloramphenicol acetyltransferase, firefly luciferase, or an immunoglobulin or portion thereof such as the Fc portion of an immunoglobulin preferably IgG. Transcription of the selectable marker gene is monitored by changes in the concentration of the selectable marker protein such as β -galactosidase, chloramphenicol acetyltransferase, or firefly luciferase. If the selectable marker gene encodes a protein conferring antibiotic resistance such as neomycin resistance transformant cells can be selected with G418. Cells that have

incorporated the selectable marker gene will survive, while the other cells die. This makes it possible to visualize and assay for expression of recombinant expression vectors of the invention and in particular to determine the effect of a mutation on expression and phenotype. It will
5 be appreciated that selectable markers can be introduced on a separate vector from the nucleic acid of interest.

The recombinant expression vectors may also contain genes which encode a fusion moiety which provides increased expression of the recombinant protein; increased solubility of the
10 recombinant protein; and aid in the purification of the target recombinant protein by acting as a ligand in affinity purification. For example, a proteolytic cleavage site may be added to the target recombinant protein to allow separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein.
15 Typical fusion expression vectors include pGEX (Amrad Corp., Melbourne, Australia), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the recombinant protein.

20 Recombinant expression vectors can be introduced into host cells to produce a transformant host cell. The term "transformant host cell" is intended to include prokaryotic and eukaryotic cells which have been transformed or transfected with a recombinant expression vector of the invention. The terms "transformed with", "transfected
25 with", "transformation" and "transfection" are intended to encompass introduction of nucleic acid (e.g. a vector) into a cell by one of many possible techniques known in the art. Prokaryotic cells can be transformed with nucleic acid by, for example, electroporation or calcium-chloride mediated transformation. Nucleic acid can be
30 introduced into mammalian cells via conventional techniques such as calcium phosphate or calcium chloride co-precipitation, DEAE-dextran mediated transfection, lipofectin, electroporation or microinjection.

Suitable methods for transforming and transfecting host cells can be found in Sambrook et al. (Molecular Cloning: A Laboratory Manual, 2nd Edition, Cold Spring Harbor Laboratory press (1989)), and other laboratory textbooks.

5 Suitable host cells include a wide variety of prokaryotic and eukaryotic host cells. For example, the proteins of the invention may be expressed in bacterial cells such as *E. coli*, insect cells (using baculovirus), yeast cells or mammalian cells. Other suitable host cells can be found in Goeddel, Gene Expression Technology: Methods in
10 Enzymology 185, Academic Press, San Diego, CA (1991).

 More particularly, bacterial host cells suitable for carrying out the present invention include *E. coli*, *B. subtilis*, *Salmonella typhimurium*, and various species within the genus' *Pseudomonas*, *Streptomyces*, and *Staphylococcus*, as well as many other bacterial
15 species well known to one of ordinary skill in the art. Suitable bacterial expression vectors preferably comprise a promoter which functions in the host cell, one or more selectable phenotypic markers, and a bacterial origin of replication. Representative promoters include the β -lactamase (penicillinase) and lactose promoter system (see Chang et al., *Nature*
20 275:615 (1978)), the trp promoter (Nichols and Yanofsky, *Meth in Enzymology* 101:155, (1983) and the tac promoter (Russell et al., *Gene* 20: 231, (1982)). Representative selectable markers include various antibiotic resistance markers such as the kanamycin or ampicillin resistance genes. Suitable expression vectors include but are not limited
25 to bacteriophages such as lambda derivatives or plasmids such as pBR322 (Bolivar et al., *Gene* 2:9S, (1977)), the pUC plasmids pUC18, pUC19, pUC118, pUC119 (see Messing, *Meth in Enzymology* 101:20-77, 1983 and Vieira and Messing, *Gene* 19:259-268 (1982)), and pNH8A, pNH16a, pNH18a, and Bluescript M13 (Stratagene, La Jolla, Calif.).
30 Typical fusion expression vectors which may be used are discussed above, e.g. pGEX (Amrad Corp., Melbourne, Australia), pMAL (New

England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ). Examples of inducible non-fusion expression vectors include pTrc (Amann et al., *Gene* 69:301-315 (1988)) and pET 11d (Studier et al., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California, 60-89 (1990)).

Yeast and fungi host cells suitable for carrying out the present invention include, but are not limited to *Saccharomyces cerevisiae*, the genera *Pichia* or *Kluyveromyces* and various species of the genus *Aspergillus*. Examples of vectors for expression in yeast *S. cerevisiae* include pYepSec1 (Baldari. et al., *Embo J.* 6:229-234 (1987)), pMFa (Kurjan and Herskowitz, *Cell* 30:933-943 (1982)), pJRY88 (Schultz et al., *Gene* 54:113-123 (1987)), and pYES2 (Invitrogen Corporation, San Diego, CA). Protocols for the transformation of yeast and fungi are well known to those of ordinary skill in the art.(see Hinnen et al., *Proc. Natl. Acad. Sci. USA* 75:1929 (1978); Itoh et al., *J. Bacteriology* 153:163 (1983), and Cullen et al. (*Bio/Technology* 5:369 (1987)).

Mammalian cells suitable for carrying out the present invention include, among others: COS (e.g., ATCC No. CRL 1650 or 1651), BHK (e.g. ATCC No. CRL 6281), CHO (ATCC No. CCL 61), HeLa (e.g., ATCC No. CCL 2), 293 (ATCC No. 1573) and NS-1 cells. Suitable expression vectors for directing expression in mammalian cells generally include a promoter (e.g., derived from viral material such as polyoma, Adenovirus 2, cytomegalovirus and Simian Virus 40), as well as other transcriptional and translational control sequences. Examples of mammalian expression vectors include pCDM8 (Seed, B., *Nature* 329:840 (1987)) and pMT2PC (Kaufman et al., *EMBO J.* 6:187-195 (1987)).

Given the teachings provided herein, promoters, terminators, and methods for introducing expression vectors of an appropriate type into plant, avian, and insect cells may also be readily accomplished. For example, within one embodiment, the proteins of the invention may be expressed from plant cells (see Sinkar et al., *J. Biosci* (Bangalore) 11:47-58 (1987), which reviews the use of

Agrobacterium rhizogenes vectors; see also Zambryski et al., Genetic Engineering, Principles and Methods, Hollaender and Setlow (eds.), Vol. VI, pp. 253-278, Plenum Press, New York (1984), which describes the use of expression vectors for plant cells, including, among others, 5 pAS2022, pAS2023, and pAS2034).

Insect cells suitable for carrying out the present invention include cells and cell lines from *Bombyx*, *Trichoplusia* or *Spodoptera* species. Baculovirus vectors available for expression of proteins in cultured insect cells (SF 9 cells) include the pAc series (Smith et al., *Mol.* 10 *Cell Biol.* 3:2156-2165 (1983)) and the pVL series (Lucklow, V.A., and Summers, M.D., *Virology* 170:31-39 (1989)). Some baculovirus-insect cell expression systems suitable for expression of the recombinant proteins of the invention are described in PCT/US/02442.

Alternatively, the proteins of the invention may also be 15 expressed in non-human transgenic animals such as, rats, rabbits, sheep and pigs (Hammer et al. *Nature* 315:680-683 (1985); Palmiter et al. *Science* 222:809-814 (1983); Brinster et al. *Proc. Natl. Acad. Sci. USA* 82:4438-4442 (1985); Palmiter and Brinster *Cell* 41:343-345 (1985) and U.S. Patent No. 4,736,866).

20 The proteins of the invention may also be prepared by chemical synthesis using techniques well known in the chemistry of proteins such as solid phase synthesis (Merrifield, *J. Am. Chem. Assoc.* 85:2149-2154 (1964)) or synthesis in homogenous solution (Houbenweyl, *Methods of Organic Chemistry*, ed. E. Wansch, Vol. 15 I and II, Thieme, 25 Stuttgart (1987)).

The present invention also provides proteins comprising an A chain of a ricin-like toxin, a B chain of a ricin-like toxin and a heterologous linker amino acid sequence linking the A and B chains, wherein the linker sequence contains a cleavage recognition site for a 30 disease-specific protease. Such a protein could be prepared other than by recombinant means, for example by chemical synthesis or by conjugation of A and B chains and a linker sequence isolated and

- 54 -

purified from their natural plant, fungal or bacterial source. Such A and B chains could be prepared having the glycosylation pattern of the native ricin-like toxin.

N-terminal or C-terminal fusion proteins comprising the protein of the invention conjugated with other molecules, such as proteins may be prepared by fusing, through recombinant techniques. The resultant fusion proteins contain a protein of the invention fused to the selected protein or marker protein as described herein. The recombinant protein of the invention may also be conjugated to other proteins by known techniques. For example, the proteins may be coupled using heterobifunctional thiol-containing linkers as described in WO 90/10457, N-succinimidyl-3-(2-pyridyldithio-propionate) or N-succinimidyl-5 thioacetate. Examples of proteins which may be used to prepare fusion proteins or conjugates include cell binding proteins such as immunoglobulins, hormones, growth factors, lectins, insulin, low density lipoprotein, glucagon, endorphins, transferrin, bombesin, asialoglycoprotein glutathione-S-transferase (GST), hemagglutinin (HA), and truncated myc.

Utility of the Nucleic Acid Molecules and Proteins of the Invention

The proteins of the invention may be used to specifically inhibit or destroy mammalian cells affected by a disease or infection which have associated with such cells a specific protease, i.e., disease-specific, for example cancer cells or cells infected with a virus, fungus or parasite, all of which are encompassed within the term "disease-specific." It is an advantage of the recombinant proteins of the invention that they have specificity for said cells without the need for a cell binding component. The ricin-like B chain of the recombinant proteins recognize galactose moieties on the cell surface and ensure that the protein is taken up by the diseased cell and released into the cytoplasm. When the protein is internalized into a non-infected cell, cleavage of the heterologous linker would not occur in the absence of the disease-specific protease and the A chain will remain inactive bound to the B

chain. Conversely, when the protein is internalized into a diseased cell, the disease-specific protease will cleave the cleavage recognition site in the linker thereby releasing the toxic A chain.

The specificity of a recombinant protein of the invention
5 may be tested by treating the protein with the disease-specific protease which is thought to be specific for the cleavage recognition site of the linker and assaying for cleavage products. Disease-specific proteases may be isolated from cancer cells or infected cells, or they may be prepared recombinantly, for example following the procedures in
10 Darket et al. (*J. Biol. Chem.* 254:2307-2312 (1988)). The cleavage products may be identified for example based on size, antigenicity or activity. The toxicity of the recombinant protein may be investigated by subjecting the cleavage products to an *in vitro* translation assay in cell lysates, for example using Brome Mosaic Virus mRNA as a template.
15 Toxicity of the cleavage products may be determined using a ribosomal inactivation assay (Westby et al., *Bioconjugate Chem.* 3:377-382 (1992)). The effect of the cleavage products on protein synthesis may be measured in standardized assays of *in vitro* translation utilizing partially defined cell free systems composed for example of a
20 reticulocyte lysate preparation as a source of ribosomes and various essential cofactors, such as mRNA template and amino acids. Use of radiolabelled amino acids in the mixture allows quantitation of incorporation of free amino acid precursors into trichloroacetic acid precipitable proteins. Rabbit reticulocyte lysates may be conveniently
25 used (O'Hare, *FEBS Lett.* 273:200-204 (1990)).

The ability of the recombinant proteins of the invention to selectively inhibit or destroy animal cancer cells or cells infected with a virus or parasite may be readily tested *in vitro* using animal cancer cell lines or cell cultures infected with the virus or parasite of interest.
30 The selective inhibitory effect of the recombinant proteins of the invention may be determined, for example, by demonstrating the selective inhibition of viral antigen expression in infected mammalian

- 56 -

cells, the selective inhibition of general mRNA translation and protein synthesis in diseased cells, or selective inhibition of cellular proliferation in cancer cells or infected cells.

Toxicity may also be measured based on cell viability, for example the viability of infected and non-infected cell cultures exposed to the recombinant protein may be compared. Cell viability may be assessed by known techniques, such as trypan blue exclusion assays.

In another example, a number of models may be used to test the cytotoxicity of recombinant proteins having a heterologous linker sequence containing a cleavage recognition site for a cancer-associated matrix metalloprotease. Thompson, E.W. et al. (*Breast Cancer Res. Treatment* 31:357-370 (1994)) has described a model for the determination of invasiveness of human breast cancer cells *in vitro* by measuring tumour cell-mediated proteolysis of extracellular matrix and tumour cell invasion of reconstituted basement membrane (collagen, laminin, fibronectin, Matrigel or gelatin). Other applicable cancer cell models include cultured ovarian adenocarcinoma cells (Young, T.N. et al. *Gynecol. Oncol.* 62:89-99 (1996); Moore, D.H. et al. *Gynecol. Oncol.* 65:78-82 (1997)), human follicular thyroid cancer cells (Demeure, M.J. et al., *World J. Surg.* 16:770-776 (1992)), human melanoma (A-2058) and fibrosarcoma (HT-1080) cell lines (Mackay, A.R. et al. *Lab. Invest.* 70:781-783 (1994)), and lung squamous (HS-24) and adenocarcinoma (SB-3) cell lines (Spiess, E. et al. *J. Histochem. Cytochem.* 42:917-929 (1994)). An *in vivo* test system involving the implantation of tumours and measurement of tumour growth and metastasis in athymic nude mice has also been described (Thompson, E.W. et al., *Breast Cancer Res. Treatment* 31:357-370 (1994); Shi, Y.E. et al., *Cancer Res.* 53:1409-1415 (1993)).

A further model may be used to test the cytotoxicity of recombinant proteins having a heterologous linker sequence containing a cleavage recognition site for a cancer-associated Cathepsin

B protease is provided in human glioma (Mikkelsen, T. et al. *J. Neurosurg.*, 83:285-290 (1995)).

Similarly, the cytotoxicity of recombinant proteins having a heterologous linker sequence containing a cleavage recognition site
5 for a malarial protease may be tested by a Plasmodium invasion assay using human erythrocytes infected with mature-stage merozoite parasites as described by McPherson, R.A. et al. (*Mol. Biochem. Parasitol.* 62:233-242 (1993)). Alternatively, in vitro cultures of human hepatic parenchymal cells may be used to evaluate schizont infectivity and
10 Plasmodium merozoite generation.

With respect to models of viral infection and replication, suitable animal cells which can be cultured *in vitro* and which are capable of maintaining viral replication can be used as hosts. The toxicity of the recombinant protein for infected and non-infected
15 cultures may then be compared. The ability of the recombinant protein of the invention to inhibit the expression of these viral antigens may be an important indicator of the ability of the protein to inhibit viral replication. Levels of these antigens may be measured in assays using labelled antibodies having specificity for the antigens. Inhibition of
20 viral antigen expression has been correlated with inhibition of viral replication (U.S. Patent No. 4,869,903). Toxicity may also be assessed based on a decrease in protein synthesis in target cells, which may be measured by known techniques, such as incorporation of labelled amino acids, such as [3H] leucine (O'Hare et al., *FEBS Lett.* 273:200-204
25 (1990)). Infected cells may also be pulsed with radiolabelled thymidine and incorporation of the radioactive label into cellular DNA may be taken as a measure of cellular proliferation. Toxicity may also be measured based on cell death or lysis, for example, the viability of infected and non-infected cell cultures exposed to the recombinant
30 protein may be compared. Cell viability may be assessed by known techniques, such as trypan blue exclusion assays.

- 58 -

Although the primary specificity of the proteins of the invention for diseased cells is mediated by the specific cleavage of the cleavage recognition site of the linker, it will be appreciated that specific cell binding components may optionally be conjugated to the proteins of the invention. Such cell binding components may be expressed as fusion proteins with the proteins of the invention or the cell binding component may be physically or chemically coupled to the protein component. Examples of suitable cell binding components include antibodies to cancer, viral or parasitic proteins.

Antibodies having specificity for a cell surface protein may be prepared by conventional methods. A mammal, (e.g. a mouse, hamster, or rabbit) can be immunized with an immunogenic form of the peptide which elicits an antibody response in the mammal. Techniques for conferring immunogenicity on a peptide include conjugation to carriers or other techniques well known in the art. For example, the peptide can be administered in the presence of adjuvant. The progress of immunization can be monitored by detection of antibody titers in plasma or serum. Standard ELISA or other immunoassay procedures can be used with the immunogen as antigen to assess the levels of antibodies. Following immunization, antisera can be obtained and, if desired, polyclonal antibodies isolated from the sera.

To produce monoclonal antibodies, antibody producing cells (lymphocytes) can be harvested from an immunized animal and fused with myeloma cells by standard somatic cell fusion procedures thus immortalizing these cells and yielding hybridoma cells. Such techniques are well known in the art, (e.g. the hybridoma technique originally developed by Kohler and Milstein (*Nature* 256:495-497 (1975)) as well as other techniques such as the human B-cell hybridoma technique (Kozbor et al., *Immunol. Today* 4:72 (1983)), the EBV-hybridoma technique to produce human monoclonal antibodies (Cole et al., *Monoclonal Antibodies in Cancer Therapy* Allen R., Bliss,

- 59 -

Inc., pages 77-96 (1985)), and screening of combinatorial antibody libraries (Huse et al., *Science* 246:1275 (1989)). Hybridoma cells can be screened immunochemically for production of antibodies specifically reactive with the peptide and the monoclonal antibodies can be isolated.

5 The term "antibody" as used herein is intended to include fragments thereof which also specifically react with a cell surface component. Antibodies can be fragmented using conventional techniques and the fragments screened for utility in the same manner as described above. For example, F(ab')₂ fragments can be generated by
10 treating antibody with pepsin. The resulting F(ab')₂ fragment can be treated to reduce disulfide bridges to produce Fab' fragments.

 Chimeric antibody derivatives, i.e., antibody molecules that combine a non-human animal variable region and a human constant region are also contemplated within the scope of the
15 invention. Chimeric antibody molecules can include, for example, the antigen binding domain from an antibody of a mouse, rat, or other species, with human constant regions. Conventional methods may be used to make chimeric antibodies containing the immunoglobulin variable region which recognizes a cell surface antigen (See, for
20 example, Morrison et al., *Proc. Natl Acad. Sci. U.S.A.* 81:6851 (1985); Takeda et al., *Nature* 314:452 (1985), Cabilly et al., U.S. Patent No. 4,816,567; Boss et al., U.S. Patent No. 4,816,397; Tanaguchi et al., E.P. Patent No. 171,496; European Patent No. 173,494, United Kingdom Patent No. GB 2177096B). It is expected that chimeric antibodies would
25 be less immunogenic in a human subject than the corresponding non-chimeric antibody.

 Monoclonal or chimeric antibodies specifically reactive against cell surface components can be further humanized by producing human constant region chimeras, in which parts of the variable
30 regions, particularly the conserved framework regions of the antigen-binding domain, are of human origin and only the hypervariable regions are of non-human origin. Such

- 60 -

immunoglobulin molecules may be made by techniques known in the art, (e.g. Teng et al., *Proc. Natl. Acad. Sci. U.S.A.*, 80:7308-7312 (1983); Kozbor et al., *Immunology Today* 4:7279 (1983); Olsson et al., *Meth. Enzymol.*, 92:3-16 (1982), and PCT Publication WO92/06193 or EP
5 239,400). Humanized antibodies can also be commercially produced (Scotgen Limited, 2 Holly Road, Twickenham, Middlesex, Great Britain.)

Specific antibodies, or antibody fragments, reactive against cell surface components may also be generated by screening expression libraries encoding immunoglobulin genes, or portions
10 thereof, expressed in bacteria with cell surface components. For example, complete Fab fragments, VH regions and FV regions can be expressed in bacteria using phage expression libraries (See for example Ward et al., *Nature* 341:544-546 (1989); Huse et al., *Science* 246:1275-1281 (1989); and McCafferty et al., *Nature* 348:552-554 (1990)). Alternatively, a
15 SCID-hu mouse, for example the model developed by Genpharm, can be used to produce antibodies, or fragments thereof.

The proteins of the invention may be formulated into pharmaceutical compositions for administration to subjects in a biologically compatible form suitable for administration *in vivo*. By
20 "biologically compatible form suitable for administration in vivo" is meant a form of the substance to be administered in which any toxic effects are outweighed by the therapeutic effects. The substances may be administered to living organisms including humans, and animals. Administration of a therapeutically active amount of the
25 pharmaceutical compositions of the present invention is defined as an amount effective, at dosages and for periods of time necessary to achieve the desired result. For example, a therapeutically active amount of a substance may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of antibody
30 to elicit a desired response in the individual. Dosage regime may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered daily or the dose may be

proportionally reduced as indicated by the exigencies of the therapeutic situation.

The nucleic acid molecules of the invention may be formulated into pharmaceutical compositions for administration to subjects in a biologically compatible form suitable for administration *in vivo*. By "biologically compatible form suitable for administration *in vivo*" is meant a form of the substance to be administered in which any toxic effects are outweighed by the therapeutic effects. The substances may be administered to living organisms including humans, and animals. Administration of a therapeutically active amount of the pharmaceutical compositions of the present invention is defined as an amount effective, at dosages and for periods of time necessary to achieve the desired result. For example, a therapeutically active amount of a substance may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of antibody to elicit a desired response in the individual. Dosage regime may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation.

The active substance may be administered in a convenient manner such as by injection (subcutaneous, intravenous, intramuscular, etc.), oral administration, inhalation, transdermal administration (such as topical cream or ointment, etc.), or suppository applications. Depending on the route of administration, the active substance may be coated in a material to protect the compound from the action of enzymes, acids and other natural conditions which may inactivate the compound.

The compositions described herein can be prepared by *per se* known methods for the preparation of pharmaceutically acceptable compositions which can be administered to subjects, such that an effective quantity of the active substance is combined in a mixture with

a pharmaceutically acceptable vehicle. Suitable vehicles are described, for example, in Remington's Pharmaceutical Sciences (Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa., USA 1985). On this basis, the compositions include, albeit not exclusively, solutions of the substances in association with one or more pharmaceutically acceptable vehicles or diluents, and contained in buffered solutions with a suitable pH and iso-osmotic with the physiological fluids.

The pharmaceutical compositions may be used in methods for treating animals, including mammals, preferably humans, with cancer or infected with a virus or a parasite. It is anticipated that the compositions will be particularly useful for treating patients with B-cell lymphoproliferative disease, (melanoma), mononucleosis, cytomegalic inclusion disease, malaria, herpes, shingles, hepatitis, poliomyelitis, or infectious laryngotracheitis. The dosage and type of recombinant protein to be administered will depend on a variety of factors which may be readily monitored in human subjects. Such factors include the etiology and severity (grade and stage) of neoplasia, the stage of malarial infection (e.g. exoerythrocytic *vs.* erythrocytic), or antigen levels associated with viral load in patient tissues or circulation.

As mentioned above, the novel recombinant toxic proteins and nucleic acid molecules of the present invention are useful in treating cancerous or infected cells wherein the cells contain a specific protease that can cleave the linker region of the recombinant toxic protein. One skilled in the art can appreciate that many different recombinant toxic proteins can be prepared once a disease associated protease has been identified. For example, the novel recombinant toxic proteins and nucleic acid molecules of the invention may be used to treat CNS tumors. Muller et al. (1993) describe increased activity of Insulin-type Growth Factor Binding Protein-3 (IGFBP-3) protease in the Cerebral Spinal Fluid of patients with CNS tumors. Cohen et al. (1992) claim that prostate-specific antigen (PSA) is an IGFBP-3 protease. The

pAP290 construct described above is a substrate for PSA. Conover et al. (1994) claim that cathepsin D is IGFBP-3 protease. The pAP276 described herein is a substrate for cathepsin D. Another example of a specific use of the invention is treatment of human glioma which has been shown to produce cathepsin D (Mikkelsen, T. et al. *J. Neurosurg*, 83:285-290 (1995)). The pAP 214 and 272 define herein are substrates for cathepsin B.

In addition, the novel proteins and nucleic acid molecules of the present invention may be used to treat cystic fibrosis. Hansen et al. (1995) describe how CF airway disease is characterized by neutrophil-dominated chronic inflammation with an excess of uninhibited neutrophil elastase (NE). NE levels in CF sputum are 350 times higher than that found in normal sputum. The pAP294 described herein is a substrate for neutrophil elastase.

As well, the novel proteins and nucleic acid molecules of the present invention may also be used to treat multiple sclerosis. Bever Jr. et al. (1994) implicate cathepsin B (possibly from inflammatory cells of hematogenous origin) in the demyelination found in multiple sclerosis. pAPs 214 and 272 defined herein present substrates for cathepsin B.

The term "animal" as used herein includes all members of the animal kingdom including mammals, preferably humans.

The following non-limiting examples are illustrative of the present invention:

EXAMPLES

Example 1

Cloning and Expression of Proricin Variants Activated by Disease-Specific Proteases

Isolation of total RNA

The proricin gene was cloned from new foliage of the castor bean plant. Total messenger RNA was isolated according to established procedures (Sambrook et al., *Molecular Cloning: A Lab*

Manual (Cold Spring Harbour Press, Cold Spring Harbour, (1989)) and cDNA generated using reverse transcriptase.

cDNA Synthesis:

Oligonucleotides, corresponding to the extreme 5' and 3' ends of the preproricin gene were synthesized and used to PCR amplify the gene. Using the cDNA sequence for preproricin (Lamb et al., Eur. J. Biochem., 145:266-270, 1985), several oligonucleotide primers were designed to flank the start and stop codons of the preproricin open reading frame. The oligonucleotides were synthesized using an Applied Biosystems Model 392 DNA/RNA Synthesizer. First strand cDNA synthesis was primed using the oligonucleotide Ricin1729C (Table 1). Three micrograms of total RNA was used as a template for oligo Ricin1729C primed synthesis of cDNA using Superscript II Reverse Transcriptase (BRL) following the manufacturer's protocol.

DNA Amplification and Cloning

The first strand cDNA synthesis reaction was used as template for DNA amplification by the polymerase chain reaction (PCR). The preproricin cDNA was amplified using the upstream primer Ricin-99 and the downstream primer Ricin1729C with Vent DNA polymerase (New England Biolabs) using standard procedures (Sambrook et al., Molecular Cloning: A Laboratory Manual, Second Edition, (Cold Spring Harbor Laboratory Press, 1989)). Amplification was carried out in a Biometra thermal cycler (TRIO-Thermalcycler) using the following cycling parameters: denaturation 95°C for 1 min., annealing 52°C for 1 min., and extension 72°C for 2 min., (33 cycles), followed by a final extension cycle at 72°C for 10 min. The 1846bp amplified product was fractionated on an agarose gel (Sambrook et al., Molecular Cloning: A Laboratory Manual, Second Edition, (Cold Spring Harbor Laboratory Press, 1989), and the DNA purified from the gel slice using Qiaex resin (Qiagen) following the manufacturer's protocol. The purified PCR fragment encoding the preproricin cDNA was then ligated (Sambrook et al., Molecular Cloning: A Laboratory Manual, Second

Edition, (Cold Spring Harbor Laboratory Press, 1989)) into an Eco RV-digested pBluescript II SK plasmid (Stratagene), and used to transform competent XL1-Blue cells (Stratagene). Positive clones were confirmed by restriction digestion of purified plasmid DNA. Plasmid DNA was
5 extracted using a Qiaprep Spin Plasmid Miniprep Kit (Qiagen).

DNA Sequencing

The cloned PCR product containing the putative preproricin gene was confirmed by DNA sequencing of the entire cDNA clone (pAP-144). Sequencing was performed using an Applied
10 Biosystems 373A Automated DNA Sequencer, and confirmed by double-stranded dideoxy sequencing by the Sanger method using the Sequenase kit (USB). The oligonucleotide primers used for sequencing were as follows: Ricin267, Ricin486, Ricin725, Ricin937, Ricin1151, Ricini1399, Ricin1627, T3 primer
15 (5'AATTAACCCTCACTAAAGGG-3') (SEQ ID NO. 128) and T7 primer (5'GTAATACGACTCACTATAGGGC-3) (SEQ ID NO. 129). Sequence data was compiled and analyzed using PC Gene software package (intelligenetics). The sequences and location of oligonucleotide primers is shown in Table 1. The oligonucleotide primers shown in Table 1
20 have been assigned the following sequence ID numbers:
Ricin-109 is referred to herein as SEQ ID NO. 130;
Ricin-99Eco is referred to herein as SEQ ID NO. 131;
Ricin267 is referred to herein as SEQ ID NO. 132;
Ricin486 is referred to herein as SEQ ID NO. 133;
25 Ricin725 is referred to herein as SEQ ID NO. 134;
Ricin 937 is referred to herein as SEQ ID NO. 135;
Ricin 1151 is referred to herein as SEQ ID NO. 136;
Ricin 1399 is referred to herein as SEQ ID NO. 137;
Ricin 1627 is referred to herein as SEQ ID NO. 138;
30 Ricin 1729C is referred to herein as SEQ ID NO. 139; and
Ricin 1729C Xba is referred to herein as SEQ ID NO. 140.

Production and Cloning of Linker Variants

- 66 -

pAP144 cut with EcoRI was used as target for PCR pairs employing the Ricin109-Eco oligonucleotide (Ricin-109Eco primer: 5-GGAGGAATCCGGAGATGAAACCGGGAGGAAATACTATTGTAAT-3 (SEQ ID No. 141)) and a mutagenic primer for the 5' half of the linker as well as the Ricin1729PstI primer (Ricin1729-PstI: 5-GTAGGCGCTGCAGATAACTTGCTGTCCTTTCAG-3 (SEQ ID No. 142)) and a mutagenic primer for the 3' half of the linker. The cycling conditions used for the PCRs were 98 degrees C for 2 min.; 98C 1 min., 52C 1 min., 72C 1 min. 15 sec. (30 cycles); 72 degrees C 10min.; 4 degrees C soak. The PCR products were then digested by EcoRI and PstI respectively, electrophoresed on an agarose gel, and the bands purified by via glass wool spin columns. Triple ligations comprising the PCR product pairs (corresponding halves of the new linker) and pVL1393 vector digested with EcoRI and PstI were carried out. Recombinant clones were identified by restriction digests of plasmid miniprep DNA and the altered linkers confirmed by DNA sequencing. See Figure 45 as an example of the cloning strategy. Recombinant clones were identified by restriction digests of plasmid miniprep DNA and the altered linkers confirmed by DNA sequencing. Note that since all altered linker variants were cloned directly into the pVL1393 vector odd-numbered pAPs were no longer required or produced.

Isolation of Recombinant Baculoviruses

Insect cells *S. frugiperda* (Sf9), and *Trichoplusia ni* (Tn368 and BTI-TN-581-4 (High Five)) were maintained on EX-CELL 405 medium (JRH Biosciences) supplemented with 10% total calf serum (Summers et al., A Manual of Methods of Baculovirus Vectors and Insect Cell Culture Procedures, (Texas Agricultural Experiment Station, 1987)). Two micrograms of recombinant pVL1393 DNA was co-transfected with 0.5 microgram of BaculoGold AcNPV DNA (Pharmingen) into 2×10^6 Tn368 insect cells following the manufacturer's protocol (Gruenwald et al., Baculovirus Expression Vector System: Procedures and Methods Manual, 2nd Edition, (San

- 67 -

Diego, CA, 1993)). On day 5 post-transfection, media were centrifuged and the supernatants tested in limiting dilution assays with Tn368 cells (Summers et al., A Manual of Methods of Baculovirus Vectors and Insect Cell Culture Procedures, (Texas Agricultural Experiment Station, 1987)). Recombinant viruses in the supernatants were then amplified by infecting Tn368 cells at a multiplicity of infection (moi) of 0.1, followed by collection of day 3 to 5 supernatants. A total of three rounds of amplification were performed for each recombinant following established procedures (Summers et al., A Manual of Methods of Baculovirus Vectors and Insect Cell Culture Procedures, (Texas Agricultural Experiment Station, 1987 and Gruenwald et al., Baculovirus Expression Vector System: Procedures and Methods Manual, 2nd Edition, (San Diego, CA, 1993)).

Expression of Mutant Proricin

Recombinant baculoviruses were used to infect 1×10^7 Tn368 or sf9 cells at an moi of 9 in EX-CELL 405 media (JRH Biosciences) with 25mM α -lactose in spinner flasks. Media supernatants containing mutant proricins were collected 3 or 4 days post-infection.

EXAMPLE 2

Harvesting and affinity column purification of pro-ricin variants

Protein samples were harvested three days post transfection. The cells were removed by centrifuging the media at 8288 g for ten minutes using a GS3 (Sorvall) centrifuge rotor. The supernatant was further clarified by centrifuging at 25400 g using a SLA-1500 rotor (Sorvall) for 45 minutes. Protease inhibitor phenylmethylsulfonyl fluoride (Sigma) was slowly added to a final concentration of 1mM. The samples were further prepared by adding lactose to a concentration of 20 mM (not including the previous lactose contained in the expression medium). The samples were concentrated to 700 mL using a Prep/Scale-TFF Cartridge (2.5ft, 10K regenerated cellulose (Millipore)) and a Masterflex pump. The samples were then

- 68 -

dialysed for 2 days in 1X Column Buffer (50 mM Tris, 100 mM NaCl, 0.02% NaN₃, pH 7.5) using dialysis tubing (10 K MWCO, 32 mm flat width(Spectra/Por)). Subsequently, the samples were clarified by centrifuging at 25400 g using a SLA-1500 rotor (Sorvall) for 45 minutes.

- 5 Following centrifugation, the samples were degassed and applied at 4 degrees C to a XK26/20 (Pharmacia) column (attached to a Pharmacia peristaltic pump, Pharmacia Single-path Monitor UV-1 Control and Optical Units, and Bromma LKB 2210 2-Channel Recorder) containing 20 mL of α -Lactose Agarose Resin (Sigma). The column was
- 10 washed for 3 hours with 1X Column buffer. Elution of pro-ricin variant was performed by eluting with buffer (1X Column buffer (0.1% NaN₃), 100 mM Lactose) until the baseline was again restored. The samples were concentrated using an Amicon 8050 concentrator (Amicon) with a YM10 76 mm membrane, utilizing argon gas to pressurize the chamber.
- 15 The samples were further concentrated in Centricon 10 (Millipore) concentrators according to manufacturer's specifications.

Purification of Variant pAP-Protein by gel filtration chromatography

- In order to purify the pro-ricin variant from processed material produced during fermentation, the protein was applied to a
- 20 SUPERDEX 75 (16/60) column and SUPERDEX 200 (16/60) column (Pharmacia) connected in series equilibrated with 50 mM Tris, 100mM NaCl, pH 7.5 containing 100 mM Lactose and 0.1% β -mercaptoethanol (β ME). The flow rate of the column was 0.15 mL/min and fractions were collected every 25 minutes. The UV (280 nm) trace was used to
- 25 determine the approximate location of the purified pAP-protein and thus determine the samples for Western analysis.

Western analysis of column fractions

- Fractions eluted from the SUPERDEX columns (Pharmacia) were analyzed for purity using standard Western blotting
- 30 techniques. An aliquot of 10 μ L from each fraction was boiled in 1X sample buffer (62.6 mM Tris-C1, pH 6.8, 4.4% β ME, 2% sodium dodecyl

- 69 -

sulfate (SDS), 5% glycerol (all from Sigma) and 0.002% bromophenol blue (Biorad)) for five minutes. Denatured samples were loaded on 12% Tris-Glycine Gels (Biorad) along with 50 ng of RCA₆₀ (Sigma) and 5 µL of kaleidoscope prestained standards (Biorad). Electrophoresis was carried out for ninety minutes at 100V in 25 mM Tris-Cl, pH 8.3, 0.1% SDS, and 192 mM glycine using the BioRad Mini Protean II cells (Biorad).

Following electrophoresis gels were equilibrated in transfer buffer (48 mM Tris, 39 mM glycine, 0.0375% SDS, and 20% Methanol) for a few minutes. PVDF Biorad membrane was presoaked for one minute in 100% methanol, rinsed in ddH₂O and two minutes in transfer buffer. Whatman paper was soaked briefly in transfer buffer. Five pieces of Whatman paper, membrane, gel, and another five pieces of Whatman paper were arranged on the bottom cathode (anode) of the Pharmacia Novablot transfer apparatus (Pharmacia). Transfer was for one hour at constant current (2 mA/cm²).

Transfer was confirmed by checking for the appearance of the prestained standards on the membrane. Non-specific sites on the membrane were blocked by incubating the blot for thirty minutes in 1X Phosphate Buffered Saline (1X PBS; 137 mM NaCl, 2.7 mM KCl, 8 mM Na₂HPO₄, 1.5 mM KH₂PO₄, pH 7.4) with 5% skim milk powder (Carnation). Primary antibody (Rabbit α-ricin, Sigma) was diluted 1:3000 in 1X PBS containing 0.1% Tween 20 (Sigma) and 2.5% skim milk and incubated with blot for forty five minutes on a orbital shaker (VWR). Non-specifically bound primary antibody was removed by washing the blot for ten minutes with 1X PBS containing 0.2% Tween 20. This was repeated four times. Secondary antibody donkey anti-rabbit (Amersham) was incubated with the blot under the same conditions as the primary antibody. Excess secondary antibody was washed as described above. Blots were developed with the ECL Western Blotting detection reagents according to the manufacturer's instructions. Blots

- 70 -

were exposed to Medtec's Full Speed Blue Film (Medtee) or Amersham's ECL Hyperfilm (Amersham) for one second to five minutes. Film was developed in a KODAK Automatic Developer.

Determination of lectin binding ability of pro-ricin variant

- 5 An Immulon 2 plate (VDVR) was coated with 100 μ L per well of 10 μ g/ml of asialofetuin and left overnight at 4°C. The plate was washed with 3X 300 μ L per well with ddH₂O using an automated plate washer (BioRad). The plate was blocked for one hour at 37°C by adding 300 μ L per well of PBS containing 1% ovalbumin. The plate was
- 10 washed again as above. Pro-ricin variant pAP-protein was added to the plate in various dilutions in 1X Baculo. A standard curve of RCA₆₀ (Sigma) from 1-10 ng was also included. The plate was incubated for 1 h at 37°C. The plate was washed as above. Anti-ricin monoclonal antibody (Sigma) was diluted 1:3000 in 1X PBS containing 0.5%
- 15 ovalbumin and 0.1% tween-20, added at 100 μ L per well and incubated for 1 h at 37°C. The plate was washed as above. Donkey-anti rabbit polyclonal antibody was diluted 1:3000 in 1X PBS containing 0.5% ovalbumin, 0.1% Tween-20, and added at 100 μ L per well and incubated for 1 h at 37°C. The plate was given a final wash as described above.
- 20 Substrate was added to plate at 100 μ L per well (1 mg/ml o-phenylenediamine (Sigma), 1 μ L/ml H₂O₂, 25 μ L of stop solution (20% H₂SO₄) was added and the absorbance read (A490nm-A630nm) using a SPECTRA MAX 340 plate reader (Molecular Devices).

Determination of pAP -Protein activity using the rabbit reticulocyte assay

Ricin samples were prepared for reduction.

A) RCA₆₀ = 3,500 ng/ μ L of RCA₆₀ + 997 μ L 1xEndo buffer (25mM Tris, 25mM KCl, 5mM MgCl₂, pH 7.6)

Reduction = 95 μ L of 10ng/ μ L + 5 μ L β -mercaptoethanol

- 71 -

B) Ricin variants

Reduction = 40 μ L variant + 2 μ L β -mercaptoethanol

The ricin standard and the variants were incubated for 30 minutes at room temperature.

5 **Ricin - Rabbit Reticulocyte lysate reaction**

- The required number of 0.5 mL tubes were labelled. (2 tubes for each sample, + and - aniline). To each of the sample tubes 20 μ L of 1X endo buffer was added, and 30 μ L of buffer was added to the controls. To the sample tubes either 10 μ L of 10ng/ μ L Ricin or 10 μ L of variant was added. Finally, 30 μ L of rabbit reticulocyte lysate was added to all the tubes. The samples were incubated for 30 minutes at 30°C using the thermal block. Samples were removed from the eppendorf tube and contents added into a 1.5 mL tube containing 1 mL of TRIZOL (Gibco). Samples were incubated for 15 minutes at room temperature.
- 15 After the incubation, 200 μ L of chloroform was added, and the sample was vortexed and spun at 12,000 g for 15 minutes at 4°C. The top aqueous layer from the samples was removed and contents added to a 1 mL tube containing 500 μ L of isopropanol. Samples were incubated for 15 minutes at room temperature and then centrifuged at 12,000 for 15
- 20 minutes at 4°C. Supernatant was removed and the pellets were washed with 1 mL of 70% ethanol. Centrifugation at 12,000 g for 5 minutes at 4°C precipitated the RNA. All but approximately 20 μ L of the supernatant was removed and air dried. Pellets from the other samples (+aniline samples) were dissolved in 20 μ L of DEPC treated ddH₂O. An
- 25 80 μ L aliquot of 1 M aniline (distilled) with 2.8 M acetic acid was added to these RNA samples and transferred to a fresh 0.5 mL tube. The samples were incubated in the dark for 3 minutes at 60°C. RNA was precipitated by adding 100 μ L of 95% ethanol and 5 μ L of 3M sodium acetate, pH 5.2 to each tube and centrifuging at 12,000 g for 30 minutes at

- 72 -

4°C. Pellets were washed with 1 mL 70% ethanol and centrifuged again at 12,000g for 5 minutes at 4°C to precipitate RNA. The supernatant was removed and air dried. These pellets were dissolved in 10µL of 0.1 X E buffer. To all samples, 10 µL of formamide loading dye was added. The

5 RNA ladder (8 µL of ladder + 8 µL of loading dye) was also included. Samples were incubated for 2 minutes at 70°C on the thermal block. Electrophoresis was carried out on the samples using 1.2% agarose, 50% formamide gels in 0.1X E buffer + 0.2% SDS. The gel was run for 90 minutes at 75 watts. RNA was visualized by staining the gel in 1 µg/µL

10 ethidium bromide in running buffer for 45 minutes. The gel was examined on a 302 nm UV box, photographed using the gel documentation system and saved to a computer disk.

Results:

Protein Expression Yields

15 Aliquots were taken at each stop of the harvesting/purification and tested. Yields of functional ricin variant were determined by ELISA. Typical results of an 2400 mL prep of infected *T. ni* cells are given below.

<u>Aliquot</u>	<u>µg pAP 220</u>
20 Before concentration and dialysis	6000
After concentration and dialysis	4931
alpha- Lactose agarose column flow through	219
alpha- Lactose agarose column elution	1058

25 Yield: $1058/6000 = 17.6\%$

Purification of pAP -Protein and Western Analysis of column fractions

Partially purified pAP-protein was applied to Superdex 75 and 200 (16/60) columns connected in series in order to remove the

contaminating non-specifically processed pAP-protein. Eluted fractions were tested via Western analysis as described above and the fractions containing the most pure protein were pooled, concentrated and re-applied to the column. The variant was applied a total of three times to the column. Final purified pAP-protein has less than 1% processed variant.

The purified pAP-protein was tested for susceptibility to cleavage by the particular protease and for activation of the A-chain of the proricin variant, (inhibition of protein synthesis). Typically, pAP-protein was incubated with and without protease for a specified time period and then electrophoresed and blotted. Cleaved pAP will run as two 30 kDa proteins (B is slightly larger) under reducing (SDS-PAGE) conditions. Unprocessed pAP-protein, which contains the linker region, will run at 60 kDa.

15 Activation of pAP -Protein variant with Specific Protease

Activation of protease treated pAP-protein is based on the method of *May et al.* (EMBO Journal. 8 301-8, 1989). Activation of ricin A chain upon cleavage of the intermediary linker results in catalytic depurination of the adenosine 4325 residue of 28S or 26S rRNA. This depurination renders the molecule susceptible to amine-catalyzed hydrolysis by aniline of the phosphodiester bond on either side of the modification site. The result is a diagnostic 390 base band. As such, reticulocyte ribosomes incubated with biochemically purified ricin A chain, released the characteristic RNA fragment upon aniline treatment of isolated rRNA (May, M.J. et al. Embo. Journal, 8:301-308 at 302-303 (1989)). It is on this basis that the assay allows for the determination of activity of a ricin A chain which has been cleaved from the intact unit containing a particular variant linker sequence.

EXAMPLE 3

30 In Vitro Protease Digestion of Proricin Variants:

Affinity-purified proricin variant is treated with individual disease-specific proteases to confirm specific cleavage in the linker

region. Ricin-like toxin variants are eluted from the lactose-agarose matrix in protease digestion buffer (50mM NaCl, 50mM Na-acetate, pH 5.5, 1mM dithiothreitol) containing 100mM lactose. Proricin substrate is then incubated at 37°C for 60 minutes with a disease-specific protease.

- 5 The cleavage products consisting ricin A and B chains are identified using SDS/PAGE (Sambrook et al., *Molecular Cloning: a Laboratory Manual*, 2nd. ed., Cold Spring Harbor Press, 1989), followed by Western blot analysis using anti-ricin antibodies (Sigma).

Cathepsin B may be obtained from Medcor or Calbiochem.

- 10 Matrix metalloproteinases may be prepared substantially as described by Lark, M.W. et al. (*Proceedings of the 4th International Conference of the Inflammation Research Association* Abstract 145 (1988)) and Welch, A.R. et al. (*Arch. Biochem. Biophys.* 324:59-64 (1995)). Candida acid protease may be prepared substantially as described in Remold, H.H. et al. (*Biochim. Biophys. Acta* 167:399-406 (1968)), Ray, T.L. and Payne, C.D. (15 *Infect. Immunol.* 58:508-514 (1990)) and Fusek, M. et al. (*FEBS Lett.* 327:108-112 (1993)). Hepatitis A protease may be prepared as described in Jewell, D.A. et al. (*Biochemistry* 31:7862-7869 (1992)). Plasmodium proteases may be prepared as described in Goldberg, D.E. et al. (*J. Exp.* 20 *Med.* 173:961-969 (1991)) and Cooper, J.A. and Bujard, H. (*Mol. Biochem. Parasitol.* 56:151-160 (1992)).

In Vitro Cytotoxicity Assay:

- Human ovarian cancer cells (e.g. MA148) are seeded in 96-well flat-bottom plates and are exposed to ricin-like toxin variants or control medium at 37°C for 16 h. The viability of the cancer cells is determined by measuring [³⁵S]methionine incorporation and is significantly lower in wells treated with the toxin variants than those with control medium.

In Vivo Tumour Growth Inhibition Assay:

- 30 Human breast cancer (e.g. MCF-7) cells are maintained in suitable medium containing 10% fetal calf serum. The cells are grown, harvested and subsequently injected subcutaneously into

ovariectomized athymic nude mice. Tumour size is determined at intervals by measuring two right-angle measurements using calipers. In animals that received ricin-like toxin variants containing the matrix metalloproteinase-sensitive linkers, tumour size and the rate of
5 tumour growth are lower than animals in the control group.

In Vivo Tumour Metastasis Assay:

The metastasis study is performed substantially as described in Honn, K.V. et al. (*Biochem. Pharmacol.* 34:235-241 (1985)). Viable B16a melanoma tumour cells are prepared and injected subcutaneously into
10 the left axillary region of syngeneic mice. The extent of tumour metastasis is measured after 4 weeks. The lungs are removed from the animals and are fixed in Bouin's solution and macroscopic pulmonary metastases are counted using a dissecting microscope. In general without therapeutic intervention, injection of 10^5 viable tumour cells
15 forms approximately 40-50 pulmonary metastases. The number of metastases in animal treated with proricin variants containing cathepsin B-sensitive linkers is substantially lower.

EXAMPLE 4

In Vitro Protease Digestion of Proricin Variants by Cancer Proteases
20 Cathepsin B or MMP-9

The general protocol for proricin digestion by cancer proteases is described in Examples 2 and 3.

In Vitro Protease Digestion of Cathepsin B Proricin Variant

Affinity-purified mutant proricin is treated with individual
25 disease-specific proteases to confirm specific cleavage in the linker region. The proricin substrate is digested in a Cathepsin B protease buffer (50 mM Sodium acetate, 2 mM EDTA, 0.05% Triton) at 40°C. Two hours and overnight (16 hr) digestion reactions are carried out using 100ng of proricin substrate and 100 and 618 ng of Cathepsin B protease
30 per reaction (CALBIOCHEM, USA). The cleavage products of proricin (ricin A and B chains) are identified using SDS/PAGE (Sambrook et al., Molecular cloning: a laboratory Manual, 2nd. ed., Cold Spring Harbor

Press, 1989), followed by Western blot analysis using anti-ricin antibodies (Sigma).

In Vitro Protease Digestion of MMP-9 Proricin Variant

Affinity-purified mutant proricin is treated with individual
5 disease-specific proteases to confirm specific cleavage in the linker
region. The proricin substrate is digested in 1X column buffer (100 mM
NaCl, 50 mM Tris, PH 7.5) at 37°C. Two hours and overnight (16 hr)
digestion reactions are set up using 50 ng of MMP-9 proricin substrate
and 20 and 200 ng of MMP-9 protease per reaction (CALBIOCHEM,
10 USA). The cleavage products of proricin (ricin A and B chains) are
identified using SDS/PAGE (Sambrook et al., Molecular cloning: a
laboratory Manual, 2nd. ed., Cold Spring Harbor Press, 1989), followed
by Western blot analysis using anti-ricin antibodies (Sigma).

The protocol for Western analysis of ricin chains is described in
15 Example 2.

Results

Figures 48 and 49 illustrate Western blots showing the cleavage
of the protease-sensitive linkers by cathepsin B (pAP 214) and MMP-9
(pAP 220) respectively. Without protease digestion, the proricin variant
20 appears as a single band at approximately 60 kDa (Lane B of Figure 48
and Lane A of Figure 49). Wild type ricin A chain and B chain appear as
two disparate bands at approximately 30 kDa (Lane A of Figure 48 and
Lane E of Figure 49). Increasing extent of proricin cleavage can clearly be
observed with increasing protease concentration (Lanes C and D of
25 Figure 48 and Lanes B-C of Figure 49).

EXAMPLE 5

In vitro protease digestion of various proricin variants by their
corresponding proteases.

The general protocol for proricin digestion by coresponding
30 proteases was as desribed in Examples 2 and 3 and should be considered
in connection with the digestions described below.

Cleavage of pAP-222 protein with the Matrix Metalloproteinase 2 (MMP-2)

Affinity-purified mutant proricin is treated with individual disease-specific proteases to confirm specific cleavage in the linker
5 region.

The pAP-222 protein sample (1.0 ug) was digested with the MMP-2 protease (1.0 ug) overnight at 37° C. The total volume of the digestion reaction was 21.5 ul, and 0.250 ug of the reaction sample was loaded on a protein gel. The MMP-2 protease was purchased from
10 Calbiochem-Novabiochem Corporation, USA.

Cleavage of pAP-248 protein with the Human Cytomegalovirus (HCMV) protease

Affinity-purified mutant proricin is treated with individual disease-specific proteases to confirm specific cleavage in the linker
15 region.

The pAP-248 protein sample (1.19 ug) was digested with the HCMV protease (1.13 ug) overnight at 37°C. The total volume of the digestion was 10.5 ul, and 0.279 ug of the reaction sample was loaded on a protein gel. The HCMV was purchased from BACHEM Bioscience Inc., USA.

Cleavage of pAP-256 protein with the Hepatitis A virus 3C (HAV 3C) protease

Affinity-purified mutant proricin is treated with individual disease-specific proteases to confirm specific cleavage in the linker
region.

25 The pAP-256 protein sample (1.26 ug) was digested with the HAV 3C protease (5 ug) overnight at 37°C. The total volume of the digestion was 12.5 ul, and 0.302 ug of the digestion sample was loaded on a protein gel. The HAV 3C protease was a gift from Dr. G. Lawson from Bates Collage, Main, USA.

Cleavage of pAP-270 protein with the Matrix Metalloproteinase 2 (MMP-2)

- 78 -

Affinity-purified mutant proricin is treated with individual disease-specific proteases to confirm specific cleavage in the linker region.

5 The pAP-270 protein sample (0.120 ug) was digested with the MMP-2 protease (0.25 ug) overnight at 37° C. The total volume of the digestion reaction was 22.5 ul, and 0.106 ug of the reaction sample was loaded on a protein gel. The MMP-2 protease was purchased from Calbiochem-Novabiochem Corporation, USA.

Cleavage of pAP-288 protein with tPA plasminogen tissue activator

10 Affinity-purified mutant proricin is treated with individual disease-specific proteases to confirm specific cleavage in the linker region. The pAP-288 protein sample (1.65 ug) was digested with the t-PA protease (0.5 ug) overnight at 37° C. The total volume of the digestion reaction was 55 ul, and 0.6 ug of the reaction sample was
15 loaded on a protein gel. The t-PA was purchased from Sigma Chemical Co., USA.

Cleavage of pAP-294 protein with human neutrophil elastase

Affinity-purified mutant proricin is treated with individual disease-specific proteases to confirm specific cleavage in the linker
20 region.

The pAP-256 protein sample (0.6 ug) was digested with the Elastase protease (5 ug) at 25° C for one hour. The total volume of the digestion reaction was 52.5 ul, and 0.171 ug of the digestion sample was loaded on a protein gel. The Human Neutrophil Elastase protease was purchased
25 from Cedarlane Laboratories Limited, Canada.

Cleavage of pAP-296 protein with calpain

Affinity-purified mutant proricin is treated with individual disease-specific proteases to confirm specific cleavage in the linker region. The pAP-296 protein sample (2.05 ug) was digested with the
30 Calpain protease (10 ug) overnight at 37° C. The total volume of the digestion reaction was 35 ul and 0.761 ug of the reaction sample was

- 79 -

loaded on a protein gel. The Calpain protease was purchased from Sigma Chemical Co., USA

Results

Figures 52, 54, 58 & 66(MMP-2), 60, 64 and 62 show the cleavage of proteases of linkers by HCMV, HAV 3C, MMP-2, t-PA, calpain, and human neutrophil elastase respectively. Without protease digestion, the proricin variants appear as a single band at approximately 60kDA (Lane A in connection with Figure 52; Lane B of Figure 54; Lane A of Figure 58; Lane B of Figure 60; and Lane C of Figure 62; lane B of Figure 64 and lane B of Figure 66). Wild type ricin chain A and B appear as two bands at approximately 30kDA (see for example Lanes C and D of Figure 52) proricin cleavage can clearly be observed with the appearance of 30kDA bands in connection with the protein which has been digested by the respective protease (see Lane B of Figure 52; Lane C of Figure 54; or Lane B of Figure 58 for examples).

EXAMPLE 6

In Vitro Translation Assay (Activation by Cancer Proteases Cathepsin B or MMP-9)

The general protocol for the rabbit reticulocyte lysate reaction to test the cytotoxicity of cancer protease-activated proricin is described briefly in Example 3 and is described in more detail in Example 2.

Results

Activation of pAP 214 and pAP 220 proricin variants by cathepsin B and MMP-9, based on the method of May et al. (EMBO J. 8:301-308, 1989), is illustrated in Figures 50 and 51 respectively. The appearance of the 390 base pair product (positive control) is observed in Lane F of Figure 50 and Lane G of Figure 51. This 390 base pair product is absent in the negative control lanes. Without cathepsin or MMP-9 activation, no or minimal N-glycosidase activity in the pAP 214 variant (Lanes H to L, Figure 50) or the pAP 220 variant (Lanes A to E, Figure 51) was observed. When the pAP 214 variant and the pAP 220 variant were activated by cathepsin or MMP-9 respectively, appearance of the 390 base

- 80 -

pair product was observed in a proricin concentration-dependent manner (Lanes A to E of Figure 50 and Lanes H to L of Figure 51). The present experimental series demonstrated the successful and selective activation of proricin variants by cancer-associated proteases.

5 **EXAMPLE 7**

The general protocol for the rabbit reticulocyte lysate reaction is described briefly in Example 3 and is described in more detail in Example 2, all of which compliments the description below.

10 **Depurination of Rabbit Reticulocyte 28S Ribosomal RNA by Digested and Undigested Ricin Variants**

Affinity-purified mutant proricin mutants which were previously digested with the disease-specific protease, were reduced with 5% 2-mercaptoethanol then diluted to 100ng, 14.2ng, 2.0ng, 291pg, and 41.7pg with 1 X ENDO buffer (25mM Tris pH 7.6, 25mM KCl, 5mM
15 MgCl₂) and incubated with rabbit reticulocyte lysate, untreated (Promega) for 30 minutes at 30°C. To compare the digested with the undigested proricin variant, the proricin in digestion buffer (according to the specific digestion protocol) was treated in the same manner as the digested sample. As a positive and negative control, 10ng of ricin A
20 chain and 1 X ENDO buffer consecutively, was incubated with rabbit reticulocyte lysate, untreated, for 30 min at 30°C.

Aniline Cleavage of rRNA and Gel Fractionation

Total RNA was then extracted from reticulocyte lysate translation mixtures with Trizol reagent (Gibco-BRL) as per
25 manufacturer's instructions. The RNA was incubated with 80ul of 1M aniline (distilled) with 2.8M acetic acid for 3 min at 60°C in the dark. Ethanol-precipitated RNA samples were dissolved in 20ul of 50% formamide, 0.1X E buffer (3.6mM Tris, 3mM NaH₂PO₄, 0.2mM EDTA), and 0.05% xylene cyanol. 10ul of this was heated to 70°C for 2 minutes,
30 loaded and electrophoresed in 1.2% agarose, 0.1X E buffer, and 50% formamide gel with RNA running buffer (0.1 X E buffer, 0.2% SDS).

Results

- 81 -

Activation of pAP-248 proricin variant by HCMV; pAP-256 by HAV3C protease; pAP-270 by MMP-2 protease; pAP-288 by t-PA protease; pAP-294 by human neutrophil elastase; pAP-296 by calpain; and pAP-222 by MMP-2 is illustrated in Figures 52, 55, 59, 61, 63, 65, and 67 respectively. The appearance of the 390 base pair product (deposit of control) is observed in lane L of Figures 53, 55, 61, 63, 65 and 67. The 390 base pair product is observed in lane A of Figures 59 (activation of pAP-270 by MMP-2). This 390 base pair product is absent in the negative control lanes. Without the specific protease activation, no or minimal activity is seen in the lanes which contained only the proricin variant without digestion (see lane A, B, C, D, and E of Figures 53, 55, 61, 63, 65, and 67). The same observation is made in connection with pAP-270 in Figure 59, however, the undigested lanes appear as H, I, J, K and L. When the variant was activated by its respective protease, there is an appearance of the 390 base pair product in a proricin concentration-dependent manner (see Lanes H, I, J, K and L of Figure 53, 55, 61, 63, 65, and 67 and Lanes A, B, C, D, and E of Figure 59). The present experimental series demonstrate the successful and selective activation of the identified proricin variants by selective corresponding proteases.

20 EXAMPLE 8

Procedure for Examining the Cytotoxicity of Ricin and Ricin Variants on the COS-1 Cell Line

Cell Preparation

After washing with 1XPBS (0.137 M NaCl, 2.68 mM KCl, 8.10 mM Na_2HPO_4 , 1.47 mM KH_2PO_4), cells in log phase growth were removed from plates with 1X trypsin/EDTA (Gibco/BRL). The cells were centrifuged at 1100 rpm for 3 min, resuspended in Dulbecco's Modified Eagle Medium containing 10%FBS and 1X pen/strep, and then counted using a haemocytometer. They were adjusted to a concentration of 5×10^4 cells•ml⁻¹. One hundred microliters per well of cells was added to wells 2B - 2G through to wells 9B - 9G of a Falcon 96 well tissue culture plate. A separate 96 well tissue culture plate was

- 82 -

used for each sample of Ricin or Ricin variant. The plates were incubated at 37°C with 5% CO₂ for 24 hours.

Toxin Preparation

The Ricin and Ricin variants were sterile filtered using a 0.22µm
5 filter (Millipore). The concentration of the sterile samples were then
quantified by A₂₈₀ and confirmed by BCA measurements (Pierce). For
the variants digested with the protease in vitro, the digests were carried
out as described in the digestion procedure for each protease. The
digests were then diluted in the 1000 ng•ml⁻¹ dilution and sterile
10 filtered. The Ricin and the undigested pAP214 in the pAP 214
cytotoxicity data were treated in the same manner but without the
Cathepsin B treatment. Ricin and Ricin variants were serially diluted
to the following concentrations: 1000 ng•ml⁻¹, 100 ng•ml⁻¹, 10 ng•ml⁻¹,
1 ng•ml⁻¹, 0.1 ng•ml⁻¹, 0.01 ng•ml⁻¹, 0.001 ng•ml⁻¹ with media
15 containing 10%FBS and 1X pen/strep.

Application of Toxin or Variants to Plates

Columns 2 to 9 were labeled: control, 1000 ng•ml⁻¹, 100 ng•ml⁻¹,
10 ng•ml⁻¹, 1 ng•ml⁻¹, 0.1 ng•ml⁻¹, 0.01 ng•ml⁻¹, 0.001 ng•ml⁻¹
consecutively. The media was removed from all the sample wells with
20 a multichannel pipettor. For each plate of variant and toxin, 50µl of
media was added to wells 2B to 2G as the control, and 50µl of each
sample dilution was added to the corresponding columns. For the
pAP220 + MMP-9 data, the plates were incubated for one hour at 37°C
with 5% CO₂, then washed once and replaced with media, then
25 incubated for 48 hours at 37°C with 5% CO₂. For the pAP 214 +
Cathepsin B data, the toxin was left on the plates and incubated for 24
hours at 37°C with 5% CO₂, then 50 µl of media was added to the wells
with the toxin and incubated for another 24 hours at 37°C with 5% CO₂.

Sample Application

- 83 -

The whole amount of media (and/or toxin) was removed from each well with a multichannel pipettor, and replaced with 100 μ l of the substrate mixture (Promega Cell Titer 96 Aqueous Non-Radioactive Cell Proliferation Assay Kit). The plates were incubated at 37°C with 5% CO₂ for 2 to 4 hours, and subsequently read with a Spectramax 340 96 well plate reader at 490nm. The IC₅₀ values were calculated using the GRAFIT software program.

Results

In experiments with pAP-214 and Cathepsin B incubated with COS-1 cells, it may be seen that cells incubated with pAP-214 alone, pAP-214 was ineffective at causing cell death (see Figure 56). However, the cytotoxicity of pAP-214 digested with Cathepsin B behaves similarly to the ricin control in COS-1 cells. This is also illustrated in Figure 56. Similarly, the cytotoxicity of undigested pAP-220 when incubated with COS-1 cells is lower than the cytotoxicity observed with COS-1 cells incubated with pAP-220 digested with MMP-9. Indeed the results suggest that the toxicity of digested pAP-220 is greater than that of ricin. (See Figure 57).

EXAMPLE 9

20 Procedure for Examining the Cytotoxicity of Ricin and Ricin Variants on Various Tissue Culture Cell Lines

Cell Preparation

After washing with 1XPBS (1.37M NaCl, 26.8mM KCl, 81mM Na₂HPO₄, 14.7mM KH₂PO₄), cells in log phase growth were removed from plates with 1X trypsin/EDTA (Gibco/BRL). The cells were centrifuged at 1100 rpm for 3 min, resuspended in media containing 10%FBS and 1X pen/strep (media used depended on the cell line being tested), and then counted using a haemocytometer. They were adjusted to a concentration of 5 X 10⁴ cells•ml⁻¹ (faster growing cell lines were adjusted to 2 X10⁴ cells•ml⁻¹). One hundred microliters per well of cells was added to wells 2B - 2G through to wells 9B - 9G of a Falcon 96 well

- 84 -

tissue culture plate. A separate 96 well tissue culture plate was used for each sample of Ricin or Ricin variant. The plates were incubated at 37(C with 5% CO₂ for 24 hours.

Toxin Preparation

5 The Ricin and Ricin variants were sterile filtered using a 0.22µm filter (Millipore). The concentration of the sterile samples were then quantified by A₂₈₀ and confirmed by a BCA measurement (Pierce). Ricin and Ricin variants were serially diluted to the following concentrations: 3000 ng•ml⁻¹, 300 ng•ml⁻¹, 30 ng•ml⁻¹, 3 ng•ml⁻¹, 0.3 ng•ml⁻¹,
10 0.03ng•ml⁻¹, 0.003 ng•ml⁻¹ with media containing 10%FBS and 1X pen/strep.

Application of Toxin or Variants to Plates

Columns 2 to 9 were labeled: control, 0.001 ng•ml⁻¹, 0.01 ng•ml⁻¹, 0.1 ng•ml⁻¹, 1ng•ml⁻¹, 10 ng•ml⁻¹, 100 ng•ml⁻¹, 1000 ng•ml⁻¹
15 consecutively. For each plate of variant and toxin, 50µl of media was added to wells 2B to 2G as the control, and 50µl of each sample dilution was added to the corresponding columns containing 100µl per well of cells (i.e. 50 µl of the 3000 ng•ml⁻¹ dilution added to the wells B-G in column 9, labeled 1000 ng•ml⁻¹). The plates were incubated for 48
20 hours at 37(C with 5% CO₂.

Sample Application

An amount of 140µl was removed from each well with a multichannel pipettor, and replaced with 100 µl of the substrate mixture (Promega Cell Titer 96 Aqueous Non-Radioactive Cell Proliferation
25 Assay Kit). The plates were incubated at 37(C with 5% CO₂ for 2 to 4 hours, and subsequently read with a Spectramax 340 96 well plate reader at 490nm. The IC₅₀ values were calculated using the GRAFIT software program.

Results

Referring to Table 2, it may be seen that the survival of cells is correlated with the proricin variant and the cell specific protease produced by the cell type. For example, in the HT1080 cell line, both pAP-214 and pAP-220 required only 2-1/2 times the amount of ricin to achieve the same level of cytotoxicity. On the other hand, pAP-224 required 193 times the amount of ricin to achieve the same level of cell death. As well, it may be seen that in the cells where expression of Cathepsin D is found, pAP-214 and 220 were more effective at causing cell death than ricin and more effective than pAP-224. Details concerning the various cells types used in these experiments are outlined below.

COS-1 (African Green Monkey Kidney Cells)

This is an SV40 transformed cell line which was prepared from established simian cells CV-1. (Reference: Gluzman, Y. (1975) Cell, 23, 175 - 182)(ATCC CRL 1650)

HT-1080 Human Fibrosarcoma

(ATCC CCL 121) This cell line was shown to produce active MMP-9 in tissue culture. References: Moore et al. (1997) Gynecologic Oncology 65, 83-88.

9L Rat Glioblastoma

Glioblastomas are generally associated with cathepsin B expression. Levels of cathepsin B expression correspond to the extent of progression of malignancy i.e. highest levels for glioblastomas over anaplastic astrocytomas over low-grade gliomas and normal brain tissue. The 9L cell line was provided by Dr. William Jia of the B.C. Cancer Agency.

References: Mikkelsen et al. (Aug. 1995) Journal of Neurosurgery 83(2), 285-290. Nakano et al. (1995) J. of Neurosurgery 83(2), 298-307.

MCF-7 Human Breast Cancer Cell Line (Epithelial)

(ATCC CRL 1555) In the absence of estrogen cathepsin B has not been shown to be elevated relative to normal cells. It can be induced with estrogen to produce Cathepsin D. Production of MMP-9 is unknown.

- 86 -

Having illustrated and described the principles of the invention in a preferred embodiment, it should be appreciated to those skilled in the art that the invention can be modified in arrangement and detail without departure from such principles. We claim all modifications
5 coming within the scope of the following claims.

All publications, patents and patent applications referred to herein are incorporated by reference in their entirety to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference in
10 its entirety.

**FULL CITATIONS FOR CERTAIN REFERENCES REFERRED TO IN
THE SPECIFICATION**

Bever Jr., C.T., Panitch, H.S., and Johnson, K.P. (1994) *Neurology* 44(4), 745-8. Increased cathepsin B activity in peripheral blood mononuclear
5 cells of multiple sclerosis patients.

Cohen, P., Graves, H.C., Peehl, D.M., Kamarei, M., Giudice, L.C., and Rosenfeld, R.G. (1992) *Journal of Clinical Endocrinology and Metabolism* 75(4), 1046-53. Prostate-specific antigen (PSA) is an insulin-like growth factor binding protein-3 protease found in seminal plasma.

10 Conover, C.A. and De Leon, D.D. (1994) *J. Biol. Chem.* 269(10), 7076-80. Acid activated insulin-like growth factor-binding protein-3 proteolysis in normal and transformed cells. Role of cathepsin D.

Hansen, G., Schuster, A., Zubrod, C., and Wahn, V. (1995) *Respiration* 62(3), 117-24. Alpha 1-proteinase inhibitor abrogates proteolytic and
15 secretagogue activity of cystic fibrosis sputum.

Muller, H.L., Oh, Y., Gargosky, S.E., Lehrnbecher, T., Hintz, R.L., and Rosenfeld, R.G. (1993) *Journal of Clinical Endocrinology and Metabolism* 77(5), 1113-9. Concentrations of insulin-like growth factor (IGF)-binding protein-3 (IGFBP-3), IGF, and IGFBP-3 protease activity in
20 cerebrospinal fluid of children with leukemia, central nervous system tumor, or meningitis.

TABLE 1

Table 1 - Sequence and Location of Oligonucleotide Primers

Name of Primer	Primer Sequence †	Corresponds to preprorin nucleotide numbers: (see Figures 8-10)
Ricin-109	5' - GGAGATGAAACCGGGAGGAAATACTATTGTAAT-3'	27 to 59
Ricin-99Eco	5' - <u>GCGGAATT</u> CCGGGAGGAAATACTATTGTAAT -3'	37 to 59
Ricin267	5' - ACGGTTTATTTTAGTTGA-3'	300 to 317
Ricin486	5' - ACTTGCTGGTAATCTGAG -3'	519 to 536
Ricin725	5' - AGAATAGTTGGGGGAGAC -3'	758 to 775
Ricin937	5' - AATGCTGATGTTTGTATG -3'	970 to 987
Ricin1151	5' - CGGGAGTCTATGTGATGA -3'	1184 to 1201
Ricin1399	5' -GCAAATAGTGGACAAGTA -3'	1432 to 1449
Ricin 1627	5' - GGATTGGTGTAGATGTG -3'	1660 to 1677
Ricin1729C	5' - ATAAGTCTGCTGTCCTTTCA -3'	1864 to 1846
Ricin1729C Xba	5' - <u>CGCTCTAGATA</u> ACTTGCTGTCCTTTCA	1864 to 1846

† Underlined sequences inserted for subcloning purposes and not included in final preprorin sequences

Table 2: **Comparative Toxicities to Selected Cell Lines of Ricin and Ricin Provariants**

<u>Cell Line</u>	<u>IC₅₀_{Ricin}</u> <u>(ng/ml)</u>	<u>IC₅₀_{pAP214}</u> <u>IC₅₀_{Ricin}</u>	<u>IC₅₀_{pAP220}</u> <u>IC₅₀_{Ricin}</u>	<u>IC₅₀_{pAP224}</u> <u>IC₅₀_{Ricin}</u>
COS-1	0.1	17	22	150
HT1080	0.5	2.46	2.14	193
9L	10.8	1.3	1.7	32.3
MCF-7 (without estrogen)	0.09	27.8	40	742

I CLAIM:

1. A purified and isolated nucleic acid having a nucleotide sequence encoding an A chain of a ricin-like toxin, a B chain of a ricin-like toxin and a heterologous linker amino acid sequence linking
5 the A and B chains, the heterologous linker sequence containing a cleavage recognition site for a disease-specific protease.
2. The nucleic acid sequence of claim 1 wherein the linker sequence contains a cleavage recognition site recognized by a protease selected from the group consisting of: a cancer associated protease, a
10 viral protease, a fungal protease, and a parasite protease.
3. A nucleic acid sequence of claim 2 wherein the A chain is ricin A chain, abrin toxin A chain, diphtheria toxin A chain, or Domain I of Pseudomonas endotoxin.
4. A nucleic acid sequence of claim 2 wherein the A chain is
15 volkensin toxin A chain, cholera toxin A chain, modeccin toxin A chain or shiga toxin A chain.
5. A nucleic acid sequence of claim 2 wherein the B chain is ricin B chain, abrin toxin A chain, diphtheria toxin B chain, or Domain II of Pseudomonas endotoxin.
- 20 6. A nucleic acid sequence of claim 2 wherein the B chain is volkensin toxin B chain, cholera toxin B chain, modeccin toxin B chain or shiga toxin B chain.
7. A nucleic acid sequence of claim 2 wherein the cleavage recognition site is recognized by a cancer-associated protease which is
25 selected from the group consisting of: cathepsin B, an Epstein-Barr

- 91 -

virus-specific protease, a matrix metalloproteinase, cathepsin L, cathepsin D, urokinase-type plasminogen activator, tissue-type plasminogen activator, human prostate-specific antigen, kallikrein, neutrophil elastase, and calpain.

5 8. A nucleic acid sequence of claim 2 wherein the cleavage recognition site is recognized by a parasitic protease which is a *Plasmodium falciparum* protease.

9. A nucleic acid sequence of claim 2 wherein the cleavage recognition site is recognized by viral protease which is selected from
10 the group consisting of: human cytomegalovirus, human herpes virus, varicella zoster virus, hepatitis A virus, hepatitis C virus, and infectious laryngotracheitis virus.

10. A nucleic acid sequence of claim 2 wherein the cleavage recognition site is recognized by fungal protease which is a *Candida* acid
15 protease.

11. A nucleic acid sequence of claim 2 having the nucleotide sequence according to SEQ ID No. 3; SEQ ID No 5; SEQ ID No 7; SEQ ID No 9; SEQ ID No 11; SEQ ID No 13; SEQ ID No 15; SEQ ID No 17; SEQ ID No 19; SEQ ID No 21; SEQ ID No 23; SEQ ID No 25; SEQ ID No 27;
20 SEQ ID No 29; SEQ ID No 31; SEQ ID No 33; SEQ ID No 35; SEQ ID No 37; SEQ ID No 39; SEQ ID No 48; SEQ ID No 50; SEQ ID No 52; SEQ ID No 54; SEQ ID No 74; SEQ ID No 77; SEQ ID No 80; SEQ ID No 83; SEQ ID No 86; SEQ ID No 89; SEQ ID No 92; SEQ ID No 95; SEQ ID No 98; SEQ ID No 101; SEQ ID No 104; SEQ ID No 107; SEQ ID No 110; SEQ ID
25 No 113; SEQ ID No 116; SEQ ID No 119; SEQ ID No 122; or SEQ ID No 125.

12. A plasmid incorporating the nucleic acid of claim 1 to 11.

- 92 -

13. A baculovirus transfer vector incorporating the nucleic acid of claim 1 to 11.
14. A recombinant protein comprising an A chain of a ricin-like toxin, a B chain of a ricin-like toxin and a heterologous linker amino acid sequence, linking the A and B chains, wherein the linker sequence contains a cleavage recognition site for a disease-specific protease.
15. The recombinant protein of claim 14 wherein the linker sequence contains a cleavage recognition site which is recognized by a protease selected from the group consisting of: a cancer, viral, fungal, and a parasitic protease.
16. A recombinant protein of claim 14 wherein the A chain is ricin A chain, abrin toxin B chain, diphtheria toxin A chain, or Domain I of Pseudomonas endotoxin.
17. A recombinant protein of claim 14 wherein the A chain is volkensin toxin A chain, cholera toxin A chain, modeccin toxin A chain or shiga toxin A chain.
18. A recombinant protein of claim 14 wherein the B chain is ricin B chain, abrin toxin B chain, diphtheria toxin B chain, or Domain II of Pseudomonas endotoxin.
19. A recombinant protein of claim 14 wherein the B chain is volkensin toxin B chain, cholera toxin B chain, modeccin toxin B chain or shiga toxin B chain.
20. A recombinant protein of claim 14 wherein the cleavage recognition site is recognized by a cancer-associated protease selected

from the group consisting of: cathepsin B, an Epstein-Barr virus-specific protease, a matrix metalloproteinase, cathepsin L, cathepsin D, urokinase-type plasminogen activator, tissue-type plasminogen activator, human prostate-specific antigen, kallikrein, neutrophil elastase, and calpain.

21. A recombinant protein of claim 14 wherein the cleavage recognition site is recognized by a parasitic protease which is a *Plasmodium falciparum* protease.
22. A recombinant protein of claim 14 wherein the cleavage recognition site is recognized by a viral protease which is selected from the group consisting of: human cytomegalovirus, human herpes virus, varicella zoster virus, hepatitis A virus, hepatitis C virus and infectious laryngotracheitis virus.
23. A recombinant protein of claim 14 wherein the cleavage recognition site is recognized by a fungal protease which is a *Candida* acid protease.
24. A recombinant protein of claim 14 having the linker amino acid sequence according to SEQ ID No. 40; SEQ ID No. 41; SEQ ID No. 42; SEQ ID No. 43; SEQ ID No. 44; SEQ ID No. 45; SEQ ID No. 46; SEQ ID No. 55; SEQ ID No. 56; SEQ ID No. 57; SEQ ID No. 58; SEQ ID No. 59; SEQ ID No. 60; SEQ ID No. 61; SEQ ID No. 62; SEQ ID No. 63; SEQ ID No. 64; SEQ ID No. 65; SEQ ID No. 66; SEQ ID No. 67; SEQ ID No. 68; SEQ ID No. 69; SEQ ID No. 70; SEQ ID No. 71; SEQ ID No. 72; SEQ ID No. 75; SEQ ID No. 78; SEQ ID No. 81; SEQ ID No. 84; SEQ ID No. 87; SEQ ID No. 90; SEQ ID No. 93; SEQ ID No. 96; SEQ ID No. 99; SEQ ID No. 102; SEQ ID No. 105; SEQ ID No. 108; SEQ ID No. 111; SEQ ID No. 114; SEQ ID No. 117; SEQ ID No. 120; SEQ ID No. 123; or SEQ ID No. 126.

- 94 -

25. A method of inhibiting or destroying cells affected by a disease, which cells are associated with a protease specific to the disease comprising the steps of:

- 5 (a) preparing a purified and isolated nucleic acid having a nucleotide sequence encoding an A chain of a ricin-like toxin, a B chain of a ricin-like toxin, and a heterologous linker amino acid sequence, linking the A and B chains, wherein the linker sequence contains a cleavage recognition site for the protease;
- 10 (b) introducing the nucleic acid into a host cell and expressing the nucleic acid in the host cell to obtain a recombinant protein comprising an A chain of a ricin-like toxin, a B chain of a ricin-like toxin and a linker amino acid sequence;
- (c) suspending the protein in a pharmaceutically acceptable carrier, diluent or excipient, and
- 15 (d) contacting the cells with the recombinant protein.

26. The method of claim 25 where the disease is one of cancer or cells infected with a fungus, virus or parasite.

27. A method of inhibiting or destroying cells affected by a disease, which cells are associated with a protease specific to the disease
20 comprising the step of contacting the cells with a recombinant protein according to any one of claims 14 to 24.

28. A method of treating a disease comprising administering a recombinant protein according to any one of claims 14 to 24 to an animal in need thereof.

25 29. A method of treating a disease comprising administering a nucleic acid molecule according to any one of claims 2 to 11 to an animal in need thereof.

- 95 -

30. A method of treating a mammal with cancer or infected with a fungus, virus or parasite, comprising the steps of preparing a recombinant protein of claim 14 wherein the linker sequence contains a cleavage recognition site for a cancer, fungal, viral or parasitic protease
5 and administering the protein to the mammal.

31. A process for preparing a pharmaceutical for treating a mammal with cancer, fungal infection, viral infection or parasitic infection, comprising the steps of :

(a) preparing a purified and isolated nucleic acid having a
10 nucleotide sequence encoding an A chain of a ricin-like toxin, a B chain of a ricin-like toxin, and a heterologous linker amino acid sequence, linking the A and B chains, wherein the linker sequence contains a cleavage recognition site for a cancer, viral or parasitic protease;

(b) introducing the nucleic acid into a host cell and expressing
15 the nucleic acid in the host cell to obtain a recombinant protein comprising an A chain of a ricin-like toxin, a B chain of a ricin-like toxin and a linker amino acid sequence;

(c) suspending the protein in a pharmaceutically acceptable carrier, diluent or excipient.

20 32. A use of a recombinant protein according to any one of claims 14 to 24 to treat a disease.

33 A use of a nucleic acid molecule according to any one of claims 1 to 11 to treat a disease.

34. A pharmaceutical composition for treating cancer or a fungal, or
25 viral, or parasitic infection in an animal comprising the recombinant protein of claim 14 and a pharmaceutically acceptable carrier, diluent or excipient.

- 96 -

35. A pharmaceutical composition for treating cancer or a fungal, or viral, or parasitic infection in an animal comprising the nucleic acid molecule of claim 2 and a pharmaceutically acceptable carrier, diluent or excipient.

1/254

FIGURE 1**Complete Sequence of Baculovirus
Transfer Vector, pVL1393**

ID PVL1393 preliminary; circular DNA; SYN;
 9632 BP.
 XX
 AC IG1137;
 XX
 DT 01-FEB-1993 (Rel. 7, Created)
 DT 01-JUL-1995 (Rel. 12, Last updated, Version
 1)
 XX
 DE E. coli plasmid vector pVL1393 - complete.
 XX
 KW cloning vector.
 XX
 OS Cloning vector
 OC Artificial sequences; Cloning vehicles.
 XX
 RN [1]
 RC p2Bac from baculovirus
 RC p2Blue from p2Bac
 RC pBlueBac from AcNPV
 RC pBlueBac2 from AcNPV
 RC pBlueBacIII from AcNPV
 RC pBlueBacHisA from AcNPV
 RC pBlueBacHisB from AcNPV
 RC pBlueBacHisC from AcNPV
 RC pVL1392, pVL1393 from pAc360
 RA ;
 RT ;
 RL The Digest 5:2-2(1992).
 XX
 CC NM (pVL1393)
 CC CM (yes)
 CC NA (ds-DNA)
 CC TP (circular)
 CC ST ()
 CC TY (plasmid)
 CC SP (British
 Biotechnology) (Invitrogen)
 CC HO (E.coli NM522) (E.coli
 INValphaF') (insect)
 CC CP ()
 CC FN (expression) (transfer)
 CC SE ()
 CC PA (pAC360)
 CC BR (pVL1392)
 CC OF ()
 CC OR ()
 XX
 FH Key Location/Qualifiers
 FH

2/254

FIGURE 1 (Cont'd)

```

FT    misc_feature      0..0
FT                                /note="1. pAc360, ori/amp/AcMNPV
polyhedrin gene
FT                                -> pVL1393 9632bp"
FT    transposon        0..0
FT                                /note="TRN AcMNPV"
FT    misc_binding      868..868
FT                                /note="SIT SacII"
FT    misc_binding      1395..1395
FT                                /note="SIT ApaI"
FT    misc_binding      1901..1901
FT                                /note="SIT XhoI"
FT    promoter          0..0
FT                                /note="PRO AcMNPV polyhedrin gene"
FT    misc_binding      0..0
FT                                /note="MCS
FT                                BamHI-SmaI-XbaI-EcoRI-NotI-XmaIII-PstI-
BglII"
FT    rep_origin        0..0
FT                                /note="ORI E. coli pMB1 (ColE1 and
pBR322)"
FT    CDS               complement(0..0)
FT                                /note="ANT E. coli beta-lactamase gene
(bla)
FT                                ampicillin resistance gene (apr/amp)"
XX
SQ    Sequence 9632 BP; 2602 A; 2122 C; 2176 G; 2732 T; 0
other;

```

```

aagctttact cgtaaagcga gttgaaggat catatttagt tgcgtttatg
agataagatt gaaagcacgt gtaaaatggt tcccgcgcgt tggcacaact
atttacaatg cggccaagtt ataaaagatt ctaatctgat atgttttaaa
acacctttgc ggcccgagtt gtttgcgtag gtgactagcg aagaagatgt
gtggaccgca gaacagatag taaaacaaaa ccctagtatt ggagcaataa
tcgatttaac caacacgtct aaatattatg atggtgtgca tttttgctgg
gcgggcctgt tatacaaaaa aattcaagta cctggccaga ctttgccgcc
tgaaagcata gttcaagaat ttattgacac ggtaaaagaa tttacagaaa
agtgtcccgg catgttggtg ggcgtgcact gcacacacgg tattaatcgc
accggttaca tgggtgtgcag atattttaatg cacaccctgg gtattgcgcc
gcaggaagcc atagatagat tcgaaaaagc cagagggtcac aaaattgaaa
gacaaaatta cgttcaagat ttattaattt aattaatatt atttgcattc
tttaacaaat actttatcct attttcaaat tgttgcgctt cttccagcga
acaaaaacta tgcttcgctt gctccggtta gcttgtagcc gatcagtggc
gttggtccaa tcgacggtag gattaggccg gatattctcc accacaatgt
tggcaacggt gatgttacgt ttatgctttt ggttttccac gtacgtcttt
tggccggtaa tagccgtaaa cgtagtgccg tcgcgcgtca cgcacaacac
cggatgtttg cgcttgctcg cggggtattg aaccgcgcga tccgacaaat
ccaccacttt ggcaactaaa tcggtgacct gcgcgtcttt tttctgcatt
atttcgtctt tcttttgcat ggtttcctgg aagccggtgt acatgcggtt
tagatcagtc atgacgcgcg tgacctgcaa atctttggcc tcgatctgct
tgtccttgat ggcaacgatg cgttcaataa actcttggtt tttacaagat
tcctcggttt tttgcgccac caccgcttgc agcgcgtttg tgtgctcggt
gaatgtcgca atcagcttag tcaccaactg tttgctctcc tcctcccgtt
gtttgatcgc gggatcgtag ttgccggtgc agagcacttg aggaattact
tcttctaaaa gccattcttg taattctatg gcgtaaggca atttgactt

```


3/254

FIGURE 1 (Cont'd)

cataatcagc	tgaatcacgc	cggatttagt	aatgagcact	gtatgcggct
gcaaatacag	cgggtcgcgc	cttttcacga	cgctgttaga	ggtaggggccc
ccatttttga	tgggtctgctc	aaataacgat	ttgtatttat	tgtctacatg
aacacgtata	gctttatcac	aaactgtata	ttttaaactg	ttagcgacgt
ccttggccac	gaaccggacc	tgttggtcgc	gctctagcac	gtaccgcagg
ttgaacgtat	cttctccaaa	tttaaatctt	ccaattttaa	cgcgagccat
tttgatacac	gtgtgtcgat	tttgcaacaa	ctattgtttt	ttaacgcaaa
ctaaacttat	tgtggtaagc	aataattaaa	tatgggggaa	catgcgccgc
tacaacactc	gtcgttatga	acgcagacgg	cgccgggtctc	ggcgcaagcg
gctaaaacgt	gttgcgcggt	caacgcggca	aacatcgcaa	aagccaatag
tacagttttg	atttgcataat	taacggcgat	tttttaaatt	atcttattta
ataaatagtt	atgacgccta	caactccccg	cccgcgttga	ctcgctgcac
ctcgagcagt	tgcgttgacgc	cttcctccgt	gtggccgaac	acgtcgagcg
ggtgggtcgat	gaccagcggc	gtgccgcacg	cgacgcacaa	gtatctgtac
accgaatgat	cgtcgggcga	aggcacgtcg	gcctccaagt	ggcaatatgt
gcaaattcga	aaatatatac	agttgggttg	tttgccgata	tctatcgtgg
cgttgggcat	gtacgtccga	acgttgattt	gcatgcaagc	cgaaattaaa
tcattgcgat	tagtgcgatt	aaaacgttgt	acatcctcgc	ttttaatcat
gccgtcgatt	aaatcgcgca	atcgagtcaa	gtgatcaaag	tgtggaataa
tgttttcttt	gtattcccga	gtcaagcgca	gcgcgtattt	taacaaacta
gccatcttgt	aagttagttt	catttaatgc	aactttatcc	aataatatat
tatgtatcgc	acgtcaagaa	ttaacaatgc	gcccgttgtc	gcatctcaac
acgactatga	tagagatcaa	ataaagcgcg	aattaaatag	cttgcgacgc
aacgtgcacg	atctgtgcac	gcgttccggc	acgagctttg	attgtaataa
gtttttacga	agcgatgaca	tgacccccgt	agtgacaacg	atcacgccca
aaagaactgc	cgactacaaa	attaccgagt	atgtcgggtga	cgttaaaact
attaagccat	ccaatcgacc	gttagtcgaa	tcaggaccgc	tgggtgcgaga
agccgcgaag	tatggcgaat	gcatcgtata	acgtgtggag	tccgctcatt
agagcgtcat	gttttagacaa	gaaagctaca	tatttaattg	atcccgatga
ttttattgat	aaattgaccc	taactccata	cacggtattc	tacaatggcg
gggttttggg	caaaatttcc	ggagtgcgat	tgtacatgct	gttaacggct
ccgcccacta	ttaatgaaat	taaaaattcc	aatttttaaaa	aacgcagcaa
gagaaacatt	tgtatgaaag	aatgcgtaga	aggaaagaaa	aatgtcgtcg
acatgctgaa	caacaagatt	aatatgcctc	cgtgtataaa	aaaaatattg
aacgatttga	aagaaaacaa	tgtaccgcgc	ggcgggtatgt	acaggaagag
gtttatacta	aactgttaca	ttgcaaacgt	ggtttctgtgt	gccaaagtgtg
aaaaccgatg	tttaatcaag	gctctgacgc	atttctacaa	ccacgactcc
aagtgtgtgg	gtgaagtcac	gcatctttta	atcaaattccc	aagatgtgta
taaaccacca	aactgccaaa	aaatgaaaac	tgtcgacaag	ctctgtccgt
ttgctggcaa	ctgcaagggg	ctcaatccta	tttgtaatta	ttgaataata
aaacaattat	aaatgctaaa	tttgtttttt	attaacgata	caaaccaaac
gcaacaagaa	catttgtagt	attatctata	attgaaaacg	cgtagttata
atcgctgagg	taatatttaa	aatcattttc	aaatgattca	cagttaattt
gcgacaatat	aatttttattt	tcacataaac	tagacgcctt	gtcgtcttct
tcttcgtatt	ccttctcttt	ttcatatttc	tcctcataaa	aattaacata
gttattatcg	tatccatata	tgtatctatc	gtatagagta	aattttttgt
tgtcataaat	atatatgtct	tttttaaatg	gggtgtatagt	accgctgcgc
atagtttttc	tgtaatttac	aacagtgtca	ttttctggta	gttcttcgga
gtgtgttgct	ttaattatta	aatttatata	atcaatgaat	ttgggatcgt
cggttttgta	caatatgttg	ccggcatagt		
acgcagcttc	ttctagtcca	attacaccat	tttttagcag	caccggatta
acataacttt	ccaaaatggt	gtacgaaccg	ttaaacaata	acagttcacc
tcccttttct	atactattgt	ctgcgagcag	ttgtttggtg	ttaaaaataa
cagccattgt	aatgagacgc	acaaactaat	atcacaaact	ggaaatgtct

4/254

FIGURE 1 (Cont'd)

ctgtcccgat	ttatttgaaa	cactacaaat	taaaggcgag	ctttcgtacc
aacttggttag	caatattatt	agacagctgt	gtgaagcgct	caacgatttg
cacaagcaca	atttcataca	caacgacata	aaactcgaaa	atgtcttata
tttcgaagca	cttgatcgcg	tgtatgtttg	cgattacgga	ttgtgcaaac
acgaaaactc	acttagcggtg	cacgacggca	cgttggagta	ttttagtcgg
gaaaaaattc	gacacacaac	tatgcacgtt	tcgtttgact	ggtacgcggc
gtgttaacat	acaagttgct	aacgtaatca	tggtcatagc	tgtttcctgt
gtgaaattgt	tatccgctca	caattccaca	caacatacga	gccggaagca
taaagtgtaa	agcctgggggt	gcctaattgag	tgagctaact	cacattaatt
gcgttgcgct	cactgcccgc	tttccagtcg	ggaaacctgt	cgtgcccagct
gcattaatga	atcggcccaac	gcgcggggag	aggcggtttg	cgtattgggc
gctcttccgc	ttcctcgctc	actgactcgc	tgcgctcggt	cgttcggctg
cggcgagcgg	tatcagctca	ctcaaaggcg	gtaatacggt	tatccacaga
atcaggggat	aacgcaggaa	agaacatgtg	agcaaaaggc	cagcaaaagg
ccaggaaccg	taaaaaggcc	gcgttgctgg	cgtttttcca	taggctccgc
ccccctgacg	agcatcacaa	aaatcgacgc	tcaagtcaga	ggtggcgaaa
cccgacagga	ctataaagat	accaggcggt	tccccctgga	agctccctcg
tgcgctctcc	tggtccgacc	ctgccgctta	ccggatacct	gtccgccttt
ctcccttcgg	gaagcgtggc	gctttctcat	agctcacgct	gtaggtatct
cagttcggtg	taggtcggtc	gtcccaagct	gggctgtgtg	cacgaacccc
ccgttcagcc	cgaccgctgc	gccttatccg	gtaactatcg	tcttgagtc
aacccggtaa				
gacacgactt	atcgccactg	gcagcagcca	ctggtaacag	gattagcaga
gcgaggtatg	taggcgggtgc	tacagagttc	ttgaagtggg	ggcctaacta
cggctacact	agaaggacag	tatttggtat	ctgcgctctg	ctgaagccag
ttaccttcgg	aaaaagagtt	ggtagctctt	gatccggcaa	acaaaccacc
gctggtagcg	gtggtttttt	tgtttgcaag	cagcagatta	cgcgcagaaa
aaaaggatct	caagaagatc	ctttgatctt	ttctacgggg	tctgacgctc
agtggaaacga	aaactcacgt	taagggattt	tggtcatgag	attatcaaaa
aggatcttca	cctagatcct	tttaaattaa	aatgaagt	ttaaatcaat
ctaaagtata	tatgagtaaa	cttgggtctga	cagttacca	tgcttaatca
gtgaggcacc	tatctcagcg	atctgtctat	ttcgttcatc	catagttgcc
tgtgcccccg	tcgtgtagat	aactacgata	cgggaggggt	taccatctgg
ccccagtgct	gcaatgatac	cgcgagaccc	acgctcacccg	gctccagatt
tatcagcaat	aaaccagcca	gccggaaggg	ccgagcgcag	aagtggctct
gcaactttat	ccgcctccat	ccagttctat	aattggttgc	gggaagctag
agtaagtagt	tcgccagtta	atagtttgcg	caacgttggt	gccattgcta
caggcatcgt	ggtgtcacgc	tcgtcgtttg	gtatggcttc	attcagctcc
ggttcccaac	gatcaaggcg	agttacatga	tcccccatgt	tgtgcaaaaa
agcgggttagc	tccttcgggtc	ctccgategt	tgtcagaagt	aagtggcccg
cagtgttatc	actcatgggt	atggcagcac	tgcataattc	tcttactgtc
atgccatccg	taagatgctt	ttctgtgact	ggtgagtact	caaccaagtc
attctgagaa	tagtgtatgc	ggcgaccgag	ttgctcttgc	ccggcgctcaa
tacgggataa	taccgcgcca	catagcagaa	ctttaaaagt	gctcatcatt
ggaaaacggt	cttcggggcg	aaaactctca	aggatcttac	cgctgttgag
atccagttcg	atgtaaccca	ctcgtgcacc	caactgatct	tcagcatctt
ttactttcac	cagcgtttct	gggtgagcaa	aaacaggaag	gcaaaatgcc
gcaaaaaaag	gaataaaggc	gacacggaaa	tggtgaatac	tcatactctt
ccttttttcaa	tattattgaa	gcatttatca	gggttattgt	ctcatgagcg
gatacatatt	tgaatgtatt	tagaaaaata	aacaaatagg	ggttccgcgc
acatttcccc	gaaaagtgcc	acctgacgtc	taagaaacca	ttattatcat
gacattaacc	tataaaaata	ggcgtatcac	gaggcccttt	cgtctcgcgc
gttttcgggtga	tgacgggtgaa	aacctctgac	acatgcagct	cccgagagacg
gtcacagctt	gtctgtaagc	ggatgccggg	agcagacaag	cccgtcaggg

5/254

FIGURE 1 (Cont'd)


atcaatatat	agttgctgat	atcatggaga	taattaaaat	gataaccatc
tcgcaaataa	ataagtattt	tactgttttc	gtaacagttt	tgtaataaaa
aaacctataa	atattccgga	ttattcatac	cgtcccacca	tcgggcgcg
atccccggta	ccttctagaa	ttccggagcg	gccgctgcag	atctgatcct
ttcctgggac	ccggcaagaa	ccaaaaactc	actctcttca	aggaaatccg
taatgttaaa	cccgaacgga	tgaagcttgt	cgttggtatg	aaaggaaaag
agttctacag	ggaaacttgg	acccgcttca	tggaagacag	cttccccatt
gttaacgacc	aagaagtgat	ggatgttttc	cttggtgtca	acatgcgtcc
cactagaccc	aaccgttggt	acaaattcct	ggcccaacac	gctctgcgtt
gcgacccccg	ctatgtacct	catgacgtga	ttaggatcgt	cgagccttca
tgggtggggc	gcaacaacga	gtaccgcac	agcctggcta	agaagggcgg
cggctgcccc	ataatgaacc	ttcactctga	gtacaccaac	tcgttcgaac
agttcatcga	tcgtgtcatc	tgggagaact	tctacaagcc	catcgtttac
atcgggtaccg	actctgctga	agaggaggaa	attctccttg	aagtttccct
ggtgttcaaa	gtaaaggagt	ttgcaccaga	cgcacctctg	ttcactggtc
cggcggtatta	aaacacgata	cattgttatt	agtacattta	ttaagcgcta
gattctgtgc	gttggtgatt	tacagacaat	tggtgtacgt	attttaataa
ttcattaaat	ttataatctt	taggggtgga	tgtagagcgg	aaaatcaa
gattttcagc	gtctttatat	ctgaatttaa	atattaaatc	ctcaatagat
ttgtaaaata	ggtttcgatt	agtttcaa	aagggttgtt	tttccgaacc
gatggctgga	ctatctaatt	gattttcgct	caacgccaca	aaacttgcca
aatctttag	cagcaatcta	gctttgtcga	tattcgtttg	tggtttgttt
tgtaataaag	gttcgacgtc	gttcaaaata	ttatgcgctt	ttgtatttct
ttcatcactg	tcgttagtgt	acaattgact	cgacgtaaac	acgttaaata
aagcttggac	atatttaaca	tcgggcgtgt	tagctttatt	aggccgatta
tcgtcgtcgt	cccaaccctc	gtcgttagaa	gttgcttccg	aagacgattt
tgccatagcc	acacgacgcc	tattaattgt	gtcggctaac	acgtccgcga
tcaaatttgt	agttgagctt	tttggaatta	tttctgattg	cgggcgtttt
tgggcggggt	tcaatcta	tgtgcccgat	tttaattcag	acaacacggt
agaaagcgat	ggtgcaggcg	gtggtaacat	ttcagacggc	aaatctacta
atggcgggcg	tgggtggagct	gatgataaat	ctaccatcgg	tggaggcgca
ggcggggctg	gcggcgagg	cggaggcgga	ggtggtggcg	gtgatgcaga
cggcggttta	ggctcaa	tctctttagg	caacacagtc	ggcacctcaa
ctattgtact	ggtttcgggc	gccgtttttg	gttgaccgg	tctgagacga
gtgcgatttt	tttcgtttct	aatagcttcc	aacaattgtt	gtctgtcgtc
taaaggtgca	gcgggttgag	gttcgctcgg	cattggtgga	gcgggcggca
attcagacat	cgatggtggt	ggtggtggtg	gaggcgctgg	aatgttaggc
acgggagaag	gtggtggcgg	cgggtgccgc	ggtataattt	gttctggttt
agtttggttcg	cgcacgattg	tgggcacggg	cgcaggcgcc	gctggctgca
caacggaagg	tcgtctgctt	cgaggcagcg	cttggggtgg	tggcaattca
atattataat	tgggaataca	atcgtaaaaa	tctgctataa	gcattgta
ttcgctatcg	tttaccgtgc	cgatatttaa	caaccgctca	atgtaagcaa
ttgtattgta	aagagattgt	ctcaagctcg	ccgcacgcgg	ataacaagcc
ttttcatttt	tactacagca	ttgtagtggc	gagacacttc	gctgtcgtcg
acgtacatgt	atgctttgtt	gtcaaaaacg	tcgttggtgca	gctttaaa
atttaaaaga	acatctctgt	tcagcaccac	tgtgtgtgtc	taaatgttgt
ttttgataat	ttgcgcttcc	gcagtatcga	cacgttcaaa	aaattgatgc
gcatcaattt	tggtgttcc	attattgaat	aaataagatt	gtacagattc
atatctacga	ttcgtcatgg	ccaccacaaa	tgctacgctg	caaacgctgg
tacaatttta	cgaaaactgc	aaaaacgtca	aaactcggtg	taaaataatc
aacggcgctg	ttggcaaaat	atctatttta	tcgcacaagc	ccactagcaa
attgtatttg	cagaaaacaa	tttcggcgca	caattttaac	gctgacgaaa
taaaagttca	ccagttaatg	agcgaccacc	caaattttat	aaaaatctat
tttaatcacg	gttccatcaa	caaccaagtg	atcgtgatgg	actacattga

cgctcagcg ggtgtgagcg ggtgtcggcg ctggtctaac tatgagcgcat
cagagcgagat tgtactgaga gtgcacacata tgcggtgtga aataccgca
agatgcgtaa gtagaaataa cggcatcagcg cggcatcgc catcaggt
gagcaacgtc tgggaaggcg gatcgtgagcg ggcctcgcg ctaatacgcc
agctgagcaa aggggagatgt gctgcagagcg gatgaagtgt ggtaaagcca
gggtttccc agtcagagcg tgtaaaaagc agggcgagtg cc

FIGURE 1 (Cont'd)

6/254

7/254

FIGURE 2A

SUBSTITUTE SHEET (RULE 26)

8/254

FIGURE 2B**WT preprorin linker****primer CATH-B1**

5' - ATGGTGCCAAATTTTAAT-3'

TCTTTGCTTATAAGGCCAGTGGTGCCAAATTTTAAT
 AGAAACGAATATTCCGGTACACCACGGTTTAAATTA

3' - TCTCGATTAAAGCAAAGAAACTG-5'

primer CATH-B2


PCR mutagenesis

ligate with pBluescript SK

pAP 213 linker
(Cathepsin-B variant)

TCTTTGCTTATAATCGAGAAATGGTGCCAAATTTTAAT
 AGAAACGAATTTAGCTCTTACCACGGTTTAAATTA

9/254

FIGURE 2C**SUBSTITUTE SHEET (RULE 26)**

10/254

FIGURE 2D

10 20 30 40 50

1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTAAATC

101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTTAATATTTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTTCGCGG
CGCCACGGTGACACGTTTCGATGTGTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAACAACCTGGAGCTGATGTGAGACATGATATACCAGTGTTCGCAA
AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGGTTTGCTTATAAACCAACGGTTTATTTTAGTTGAACTCTCA
TGTCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT

301 AATCATGCAGAGCTTCTGTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT

351 TGTGGTTCGGCTACCGTGTGGAATAGCGCATATTTCTTTCATCCTGACA
ACACCAGCCGATGGCAGGACCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT
TAGTCCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGAAGTACAAGTTTTA

451 CGATATACATTTCGCTTTGGTGGTAATTATGATAGACTTGAACAACCTGCG
GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACCTGTTGAACG

501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCTTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA

601 CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTTCAAGAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTACTAAAGTCTTCGTCGTTTC

651 ATTCCAATATATTGAGGGAGAAAATGCGCACGAGAATTAGGTACAACCGGA
TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAAATGTGAACCTTATCAACCCCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACCGGTGGAGGTGGT

901 TCGTCACAGTTTTCTTTGCTTAAATCGAGAATGGTGCCAAATTTTAATGC
AGCAGTGTCAAAGAAACGAATTTAGCTCTTACCACGGTTTAAATACG

SUBSTITUTE SHEET (RULE 26)

11/254

FIGURE 2D (CONT'D)

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCCTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACCTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTCAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTACGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCTCAG
TCCGACTTGTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAATTTAAACATATCACCTAACCAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCCTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT


1701 TGGTGACCCAAACCAATATGGTTACCATTTTGTATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

12/254

FIGURE 3A

SUBSTITUTE SHEET (RULE 26)

13/254

FIGURE 3B

WT preproreicin linker

primer MMP3-1

5'- TTTTGGACTTATGAATGCTGATGTTGT -3'

TCTTGTCTTATAAGGCCAGTGGTGCCCAATTTTAAT
AGAAACGAATATTCGGTCACCCACGGTTAAAAATTA

3'- GGTAGCAGTGTCAAAGCAGGCTTCGGTGTCGTT -5'

primer MMP3-2


PCR mutagenesis

Ligate with pBluescript SK

pAP 215 linker
(MMP-3 variant)

CGTCCGAAGCCACAGCAATTTTGGACTTATGAAT
GCAGGCTTCGGTGTCTGTTAAAAAACCTGAATACTTA

14/254

FIGURE 3C

SUBSTITUTE SHEET (RULE 26)

15/254

FIGURE 3D

10 20 30 40 50

1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTATC

101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTTAATATTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTCCGCG
CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAACAACCTGGAGCTGATGTGAGACATGATATACCAGTGTGCCAA
AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGGTTTGCCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA
TGTCTCAACCAACGGATATTGGTTGCCAAATAAAATCAACTTGAGAGT

301 AATCATGCAGAGCTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT

351 TGTGGTCGGCTACCGTGCTGGAATAGCGCATATTTCTTTTCATCCTGACA
ACACCAGCCGATGGCAGACCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT
TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTTA

451 CGATATACATTTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACCTGC
GCTATATGTAAGCGGAAACCACCATTAATACTATCTGAACTTGTTGAACG

501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCTTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA

601 CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTCAGAAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT

651 ATTCCAATATATTGAGGGAGAAATGCGCAGGAGAAATAGGTACAACCGGA
TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAATGTGAACTCTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTCGTCCGAAGCCACAGCAATTTTTTGGACTTATGAATGC
AGCAGTGTCAAAGCAGGCTTCGGTGTGTTAAAAACCTGAATACTTACG

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCATCGTAGGTGCAAAATG

16/254

FIGURE 3D (CONT'D)

ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACCTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTCAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTTACAACCATTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACCTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTGTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAATTTAAACATATCACCTAACCAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCCTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT


1701 TGGTGACCCAAACCAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

17/254

FIGURE 4A

SUBSTITUTE SHEET (RULE 26)

18/254

FIGURE 4B**WT preprotricin linker****primer MMP7-1**

5' - TTGTGGCGAAGTTTAAATGCTGATGTT-3'

TCTTTGCTTATAAGGCCAGTGGTGCCAAATTTTAAT
 AGAACGGAATATTCCGGTCACACGGTTAAAATTA

3' - AGTGTCAAAAGAACGCGGTGACCGT-5'

primer MMP7-2


PCR mutagenesis

ligate with pBluescript SK

**pAP 217 linker
(MMP-7 variant)**

TCTTTGCGTCCACTGGCATGTGGCGAAGTTTAAAT
 AGAACGCGAGGTGACCGTAACACCGCTTCAAATTA

19/254

FIGURE 4C

SUBSTITUTE SHEET (RULE 26)

20/254

FIGURE 4D

10 20 30 40 50

1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTATC

101 AGGATAACAACATATTTCCCAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTAAATATTTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTGCGCGG
CGCCACGGTGACACGTTTCGATGTGTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAACAACTGGAGCTGATGTGAGACATGATATACCAGTGTGCCAA
AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGGTTTGCTATAAAACCAACGGTTTATTTTAGTTGAACCTCTCA
TGCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT

301 AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT

351 TGTGGTTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA
ACACCAGCCGATGGCAGACCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT
TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTT

451 CGATATACATTTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACCTGCG
GCTATATGTAAGCGGAAACCACCATTAATACTATCTGAACCTGTTGAACG

501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCCCTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA

601 CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTTCAAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTT

651 ATTCCAATATATTGAGGGAGAAATGCGCAGGAGAATTAGGTACAACCGGA
TAAGGTTATATACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAATGTGAACTCTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCTTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTTCTTTGCGTCCACTGGCATTGTGGCGAAGTTTAAATGC
AGCAGTGTCAAAGAAACGAGGTGACCGTAACACCGCTTCAAATACG

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTGGAATG


SUBSTITUTE SHEET (RULE 26)

21/254

FIGURE 4D (CONT'D)

ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC
1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACCTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT
1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTCAGATTATGTCTACGTTTAGTCGAGACCTGAAA
1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC
1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT
1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG
1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCACCATGGTGTG
1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA
1351 AATAATACACAACCTTTTTGTTACAACCATGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC
1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT
1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC
1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA
1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA
1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA
1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT
1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA
1751 CTCTTGCAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT
1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG
1851 TGCAG
ACGTC

22/254

FIGURE 5A

SUBSTITUTE SHEET (RULE 26)

23/254

FIGURE 5B**WT preproricin linker**

primer MMP9-1

5'- GGGCAGCGAAATTTTAATGCTGAT -3'

TCTTTGCTTATAAGGCCAGTGGTGCCAAATTTTAAT
 AGAAACGAAATATTCGGT CACCACGGTTTAAATTA

** **** ***
 3'- AGCAGTGTCAAAAGAGGCGTTCCTTAACGT-5'

primer MMP9-1


PCR mutagenesis

ligate with pBluescript SK

pAP 219 linker
 (MMP-9 variant)

TCTCCGCAAGGAATTGCAGGCGAGCGAAATTTTAAT
 AGAGGCGTTCCTTAACGTCGCCGTCGCTTTAAATTA

24/254

FIGURE 5C

SUBSTITUTE SHEET (RULE 26)

25/254

FIGURE 5D

10 20 30 40 50

1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC

101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTAAATATTTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTCCGGG
CGCCACGGTGACACGTTTCGATGTGTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAACAACTGGAGCTGATGTGAGACATGATATACCAGTGTGCCAA
AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA
TGTCTCAACCAAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT

301 AATCATGCAGAGCTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT

351 TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTTCATCCTGACA
ACACCAGCCGATGGCAGACCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAAT
TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGAATAAGTTTGA

451 CGATATACATTTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACTTGC
GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACTTGTGTAACG

501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA

601 CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTTCAAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT

651 ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA
TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAAATGTGAACTCTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTTCTCCGCAAGGAATTCAGGGCAGCGAAATTTTAATGC
AGCAGTGTCAAAAGAGGCGTTCCTTAACGTCCCGTCGCTTTAAATACG

SUBSTITUTE SHEET (RULE 26)

26/254

FIGURE 5D (CONT'D)

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAACGCAATA
CAGATACACAAC TACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTT CAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGT TAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCAACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGT TACAACCATTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTCTGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTGTCAACCGAGAAATACGTC TACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAAACGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGT TAGAT
AGTTCTTACTACCTTGGTAAATTTAAACATATCACCTAACCACAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT

1701 TGGTGACCCAAACCAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA


1751 CTCTTGCAGTGTGTGTCTCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

SUBSTITUTE SHEET (RULE 26)

27/254

FIGURE 6A

SUBSTITUTE SHEET (RULE 26)

28/254

FIGURE 6B

WT preproreicin linker

primer THERM-1

5' - AGGGAATTTCCTCTTTTATGCTGATGTTGTATG - 3'

TCTTTGCTTATAAGGCCAGTGTGCCCAAATTTTAAT
AGAAACGAATATTCCGGTCACCGGTTAAATTA

GGTGGTAGCAGTGTCAAACTACACCTACTTTCCCTACAC - 5'

3' -

primer THERM-2


↓ PCR mutagenesis

↓ ligate with pBluescript SK

New Cleavage Site

pAP 221 linker
(THERM-MMP variant)GATGTGGATGAAAGGGATGTGAGGGAATTTGCTCTTTTAA
CTACACCTACTTTCCCTACACTCCCTTAAACGAAGAAAAT

29/254

FIGURE 6C

SUBSTITUTE SHEET (RULE 26)

30/254

FIGURE 6D

10 20 30 40 50
1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA
51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTAAATC
101 AGGATAACAACATATTCCTCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTTAATATTGAAATGGTGT
151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTGCGGG
CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC
201 TCGTTTAACTGAGCTGATGTGAGACATGATATACCAGTGTGCGCAA
AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT
251 ACAGAGTTGGTTTGCTATAAAACCAACGGTTTATTTTAGTTGAACCTCTCA
TGTCTCAACCAACGGATATTGTTGCGCAAATAAAATCAACTTGAGAGT
301 AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT
351 TGTGGTTCGGCTACCGTGTGGAATAGCGCATATTTCTTTCATCCTGACA
ACACCAGCCGATGGCAGACCTTTATCGCGTATAAAGAAAGTAGGACTGT
401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAAT
TAGTCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGAATAAAGTTTAA
451 CGATATACATTGCGCTTTGGTGGTAATTATGATAGACTTGAACAACCTGCG
GCTATATGTAAGCGGAAACCACCATTAATACTATCTGAACCTGTTGAACG
501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC
551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAACCT
GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA
601 CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTGAGAAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTC
651 ATTCCAATATATTGAGGGAGAAATGCGCAGCAGAAATTAGGTACAACCGGA
TAAGGTTATATACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT
701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAAATGTGAACCTCTTATCAACCCCTCT
751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCTCGGAAACGATCAGGTTA
801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAGTCACACATGCTACACTCAT
851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT
901 TCGTCACAGTTTGATGTGGATGAAAGGGATGTGAGGGAATTTGCTTCTTT
AGCAGTGTCAAACCTACACCTACTTTCCCTACACTCCCTTAAACGAAGAAA
951 TTTAGCTGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTC


SUBSTITUTE SHEET (RULE 26)

31/254

FIGURE 6D (CONT'D)

AAATCGACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAG
1001 GAAATGGTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAAC
CTTACCAGATACACAAC TACAATCCCTACCTTCTAAGGTGTTGCCTTG
1051 GCAATACAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTG
CGTTATGTCAACACCGGTACGTT CAGATTATGTCTACGTTTAGTCGAGAC
1101 GACTTTGAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTAACTA
CTGAAACTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGAT
1151 CTTACGGGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACT
GAATGCCCCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGA
1201 GCTGCAACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCAT
CGACGTTGACTACGGTGGGCGACCGTTTATACCTATTACCTTGGTAGTA
1251 AAATCCCAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGA
TTTAGGGTCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCACCAT
1301 CCACACTTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTT
GGTGTAATGTACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAA
1351 CCTACTAATAATACACAACCTTTTGTACAACCATTTGTTGGGCTATATGG
GGATGATTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACC
1401 TCTGTGCTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCA
AGACACGAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGT
1451 GTGAAAAGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGT
CACTTTTCCGACTTGTTGTACCCGAGAAATACGTCTACCAAGTTATGCA
1501 CCTCAGCAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGA
GGAGTCGTTTGGCTCTATTACGGAATGTTCACTAAGATTATATGCCCT
1551 AACAGTTGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGAT
TTGTCAACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTA
1601 GGATGTTCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGAATGGTG
CCTACAAGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCAC
1651 TTAGATGTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCC
AATCTACACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGG
1701 TCTCCATGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAG
AGAGGTACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTC
1751 ATTACTCTCTTGCAGTGTGTGTCTGTCCTGCCATGAAAAATAGATGGCTTAAA
TAATGAGAGAACGTACACACACAGGACGGTACTTTTATCTACCGAATTT
1801 TAAAAAGGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCG
ATTTTTCCTGTAAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGC
1851 AATTCCTGCAG
TTAAGGACGTC

32/254

FIGURE 7A

SUBSTITUTE SHEET (RULE 26)

33/254

FIGURE 7B

WT preprorocin linker

primer MAL-A1

5' - AATTATGATGAAGAGGATGCTGATGTTGTATG -3'

TCTTTGCTTATAAGGCCAGTGGTGCCCAATTTTAAT
 AGAAACGAATATTCGGTCACCCACGGTTAAAAATTA

3' - GGTAGCAGTGTCAAAAGTCCACCAAGTTAACGTC -5'

primer MAL-A2


PCR mutagenesis

ligate with pBluescript SK

pAP 223 linker
(MAL-A variant)

CAGGTGGTTCAATTGCAGAATTATGATGAAGAGGAT
 GTCCACCAAGTTAACGTCCTAATACTACTTCTCCTA

34/254

FIGURE 7C

SUBSTITUTE SHEET (RULE 26)

35/254

FIGURE 7D

10 20 30 40 50

1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51 GGCAACATGGCTTTGTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTAACT

101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTTAATATTTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTTCGCGG
CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAACAACCTGGAGCTGATGTGAGACATGATATACCAGTGTGGCAA
AGCAAATGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGCTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA
TGTCTCAACCAAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT

301 AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT

351 TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA
ACACCAGCCGATGGCAGCACCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAAT
TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAACCTTTTA

451 CGATATACATTGCGCTTTGGTGGTAATTATGATAGACTTGAACAACCTTGC
GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACTTGTGTAACG

501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCTTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA

601 CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTTCAGAAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT

651 ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA
TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAATGTGAACCTTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTCAGGTGGTTCAATTGCAGAATTATGATGAAGAGGATGC
AGCAGTGTCAAAGTCCACCAAGTTAAGTCTTAATACTACTTCTCCTACG

36/254

FIGURE 7D (CONT'D)

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACACTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTACAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTCGATCTAATGGAAAGTGTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCCTCAG
TCCGACTTGTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAACGGAATGTTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTAGAT
AGTTCTTACTACCTTGGTAAATTTAAACATATCACCTAACCAACAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT


1701 TGGTGACCCAAACCAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCCTGCAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

37/254

FIGURE 8A

SUBSTITUTE SHEET (RULE 26)

38/254

FIGURE 8B**WT preprorcin linker****primer MAL-B1**

5' - TCGGAGGACAATGATGAAGCTGATGTTGTATG - 3'

TCTTTGCTTATAAGGCCACGTGGTGCCAAATTTTAAT
 AGAAACGAATATTCCGGTCACCCACGGTTTAAATTA

3' - GGTAGCAGTGTCAAAAACGGCTAAAAGCCCCCTT - 5'


primer MAL-B2

↓ PCR mutagenesis
 ↓ ligate with pBluescript SK

pAP 225 linker
(MAL-B variant)

TTGCCGATTTTCGGGGAATCGGAGGACAATGATGAA
 ACGGCTAAAAGCCCCCTTAGCCTCCTGTTACTACTT

39/254

FIGURE 8C

SUBSTITUTE SHEET (RULE 26)

40/254

FIGURE 8D

10 20 30 40 50
1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA
51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTAAATC
101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTAAATATTTGAAATGGTGT
151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTGCGGG
CGCCACGGTGACACGTTTCGATGTGTTGAAATAGTCTCGACAAGCGCC
201 TCGTTTAACTGAGCTGATGTGAGACATGATATACCAGTGTGTCGCAA
AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT
251 ACAGAGTTGGTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA
TGTCTCAACCAAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT
301 AATCATGCAGAGCTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAAATGTAATCGCGACCTACAGTGGTTACGTAT
351 TGTGGTTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA
ACACCAGCCGATGGCACGACCTTTATCGCGTATAAAGAAAGTAGGACTGT
401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAAT
TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTTA
451 CGATATACATTTCGCCTTTGGTGGAATTATGATAGACTTGAACAACCTGCG
GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACTTGTGTAACG
501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC
551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA
601 CTGGCTCGTTCTTTATAATTTGCATCCAAATGATTTCAGAAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT
651 ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA
TAAGGTTATATACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT
701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAATGTGAACCTTATCAACCCCTCT
751 CTTTCACTGCAATTCAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCTCGGAAACGATCAGGTTA
801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTAAGTCACACATGCTACACTCAT
851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT
901 TCGTCACAGTTTTTGGCGATTTTCGGGGAATCGGAGGACAATGATGAAGC
AGCAGTGTCAAAAACGGCTAAAAGCCCCCTAGCCTCCTGTTACTACTTCG

SUBSTITUTE SHEET (RULE 26)

41/254

FIGURE 8D (CONT'D)

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACCTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTACAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCAACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACAAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTGTACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT


1701 TGGTGACCCAAACCAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

42/254

FIGURE 9A

SUBSTITUTE SHEET (RULE 26)

43/254

FIGURE 9B

WT preprorin linker

primer MAL-C1

5' - GCGATATCAGTTACTATGGCTGATGTTTGTATG -3'

TCTTTGCTTATAAGGCCAGTGGTGCCAAATTTTAAT
 AGAAACGAATATTCCGGTCACCCACGGTTTAAATTA

3' - GGTAGCAGTGTCAAAGTCCACCAATGTCCCCCTT -5'

primer MAL-C2


PCR mutagenesis

ligate with pBluescript SK

pAP 227 linker
(MAL-C variant)

CAGGTGGTTACAGGGGAAGCGATATCAGTTACTATG
 GTCCACCAATGTCCCCCTTCGCTATAGTCAATGATAC

44/254

FIGURE 9C

SUBSTITUTE SHEET (RULE 26)

45/254

FIGURE 9D

10 20 30 40 50

1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGAATC

101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTTAATATTTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTCCGCG
CGCCACGGTGACACGTTTCGATGTGTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAACTGAGCTGATGTGAGACATGATATACCAGTGTTCGCAA
AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA
TGTCTCAACCAAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT

301 AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT

351 TGTGGTTCGGCTACCGTGCTGGAATAGCGCATATTTCTTTCATCCTGACA
ACACCAGCCGATGGCAGGACCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT
TAGTCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGAATAAAGTTT

451 CGATATACATTTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACCTGC
GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACTTGTGAAACG

501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTGA

601 CTGGCTCGTTTCCTTTATAATTTGCATCCAAATGATTTGAGAAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTT

651 ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA
TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAATGTGAACTCTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTCAGGTGGTTACAGGGGAAGCGATATCAGTTACTATGGC
AGCAGTGTCAAAGTCCACCAATGTCCCTTCGCTATAGTCAATGATACCG

46/254

FIGURE 9D (CONT'D)

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACATAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTACGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCTTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCAACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACAACCATGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTCAACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTAGAT
AGTTCTTACTACCTTGGTAAATTTAAACATATCACCTAACCACAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT


1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAGTGTGTGTCTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCTGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

47/254

FIGURE 10A

SUBSTITUTE SHEET (RULE 26)

48/254

FIGURE 10B

WT preprorin linker

primer MAL-D1

5' - CTGTCGTTCCCTACTAATGCTGATGTTGT -3'

TCTTTGCTTATAAGGCCA
 AGAAACGAATATTCCGGT

3' - GGTAGCAGTGTCAAACGAAACCTCTCTTGCAAG -5'

primer MAL-D2

PCR mutagenesis


ligate with pBluescript SK

pAP 229 linker

(MAL-D variant)

GCTTTGGAGAGAACGTTCCCTGCTTCCCTACTAAT
 CGAAACCTCTCTTGCAAGGACAGCAAGGGATGATTA

49/254

FIGURE 10C

SUBSTITUTE SHEET (RULE 26)

50/254

FIGURE 10D

10 20 30 40 50

1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC

101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTTAATATTTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTGCGGG
CGCCACGGTGACACGTTTCGATGTGTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAACAACCTGGAGCTGATGTGAGACATGATATACCAGTGTGCGCA
AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTAGTTGAACTCTCA
TGTCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT

301 AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT

351 TGTGGTTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA
ACACCAGCCGATGGCACGACCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT
TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTGA

451 CGATATACATTTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACCTGC
GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACTTGTGTAACG

501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCTTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA

601 CTGGCTCGTTCCCTTTATAATTTGCATCCAAATGATTTTCAAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT

651 ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA
TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAAATGTGAACTCTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGAGGTTGGT

901 TCGTCACAGTTTGTCTTGGAGAGAACGTTTCCTGTCGTTCCCTACTAATGC
AGCAGTGTCAAACGAAACCTCTCTTGCAAGGACAGCAAGGGATGATTACG

SUBSTITUTE SHEET (RULE 26)

51/254

FIGURE 10D (CONT'D)

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCCTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACATAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTCAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCACCATGGTGTG

1301 TTACAGTGCAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTTCACAACCATTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTACCCCGAGAAATACGTTTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTGGCTCTATTAAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTAGAT
AGTTCTTACTACCTTGGTAAATTTAAACATATCACCTAACCACAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT


1701 TGGTGACCCAAACCAAATATGTTTACCATTATTTTGTATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAGTGTGTGTCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCTGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

52/254

FIGURE 11A

SUBSTITUTE SHEET (RULE 26)

53/254

FIGURE 11B

WT preproridin linker

primer MAL-E1

5'- AATAATTCACAGCATCAGGCTGATGTTGTATG -3'

TCTTTGCTTATAAGGCCA
 AGAACGAATATTCGGT

3'- GGTAGCAGTGTCAAAATTAAAGGTTCTATACGAT -5'

primer MAL-E2


PCR mutagenesis

ligate with pBluescript SK

pAP 231 linker
(MAL-E variant)

AAATTCCAAGATATGCTAAATAATTACAGCATCAG
 TTTAAGGTTCTATACGATTTTAAAGTGTCTAGTC

54/254

FIGURE 11C

SUBSTITUTE SHEET (RULE 26)

55/254

FIGURE 11D

10 20 30 40 50

1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTAAATC

101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTAAATATTTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTCCGGG
CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAACAACCTGGAGCTGATGTGAGACATGATATACCAGTGTGCCAA
AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGGSTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA
TGTCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT

301 AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT

351 TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA
ACACCAGCCGATGGCAGGACCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAAT
TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTTA

451 CGATATACATTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACCTGCG
GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACCTGTTGAACG

501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAACCT
GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA

601 CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTCAAGAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT

651 ATTCCAATATATTGAGGGAGAAAATGCGCACGAGAATTAGGTACAACCGGA
TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAATGTGAACTCTTATCAACCCCTCT

751 CTTTCCAATGCAATCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTTCCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTAAATTCCAAGATATGCTAAATAATTCACAGCATCAGGC
AGCAGTGTCAAATTTAAGGTTCTATACGATTTATTAAGTGTGCTAGTCCG

56/254

FIGURE 11D (CONT'D)

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACATAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTACAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTGAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCAACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACAAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAACCAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTACCCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTAGAT
AGTTCTTACTACCTTGGTAAATTTAAACATATCACCTAACCACAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT


1701 TGGTGACCCAAACCAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTTCTGTGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

57/254

FIGURE 12A

SUBSTITUTE SHEET (RULE 26)

58/254

FIGURE 12B**WT preprorcin linker**

primer HSV1-A

5' - TCGTCGGCACATGTTAATGCTGATGTTTGT -3'

TCTTTGCTTATAAGGCCAGTGGTGCCAAATTTTAAT
 AGAAACGGAATATTCCGGT CACCACGGTTTAAATTA

3' - AGCAGTGTCAAAAGACGCGAACATTTCGGT -5'

primer HSV1-A


PCR mutagenesis

ligate with pBluescript SK

pAP 233 linker
 (HSV1-A variant)

TCTGCGCTTGTAACGCATCGTCGGCACATGTTAAT
 AGACCGGAACATTTGCGTAGCAGCCGTGTACAATTA

59/254

FIGURE 12C

SUBSTITUTE SHEET (RULE 26)

60/254

FIGURE 12D

10 20 30 40 50

1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51 GGCAACATGGCTTTGTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC

101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTAAATATTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTGCGCGG
CGCCACGGTGACACGTTTCGATGTGTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAACAACTGGAGCTGATGTGAGACATGATATACCAAGTGTGCCAA
AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA
TGTCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT

301 AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT

351 TGTGGTGGCTACCGTGCTGGAAATAGCGCATATTTCTTTTCATCCTGACA
ACACCAGCCGATGGCAGCAGCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAAT
TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTTA

451 CGATATACATTGCGCTTTGGTGGTAATTATGATAGACTTGAACAACCTGCG
GCTATATGTAAGCGGAAACCACCATTAATACTATCTGAACCTGTTGAACG

501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCTTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA

601 CTGGCTCGTTTCTTTATAATTTGCATCCAAATGATTTTCAAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT

651 ATTCCAATATATTGAGGGAGAAATGCGCAGGAGAATTAGGTACAACCGGA
TAAGGTTATATACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAAATGTGAACCTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTTCTGCGCTTGTAACGCGATCGTCGGCACATGTTAATGC
AGCAGTGTCAAAGACGCGAACATTTGCGTAGCAGCCGTGTACAATTACG

SUBSTITUTE SHEET (RULE 26)

61/254

FIGURE 12D (CONT'D)

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACACTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTCAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAATCGTCGCTGTAGTCCCTTGTCAACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTTACAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTACATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTCAACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT


1701 TGGTGACCCAAACCAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAAGTGTGTGTCTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAAGTAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

62/254

FIGURE 13A

SUBSTITUTE SHEET (RULE 26)

63/254

FIGURE 13B

WT preproridin linker

primer HSV1-B

5' - TCGGAGAAATTTAAGAAATGCTGATGTTTGT -3'

TCTTTGCTTATAAGGCCAGTGGTGCCCAATTTTAAT
 AGAAACGAATATTCCCGGT CACCACGGTTTAAATTA

3' - AGCAGTGTCAAAAGATGCATAAATGTCCGT -5'

primer HSV1-B


PCR mutagenesis

ligate with pBluescript SK

pAP 235 linker
(HSV1-B variant)

TCTACGTATTTACAGGCATCGGAGAAATTTAAGAAT
 AGATGCATAAATGTCCGTAGCCTCTTTAAATCTTA

64/254

FIGURE 13C

SUBSTITUTE SHEET (RULE 26)

65/254

FIGURE 13D

10 20 30 40 50

1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC

101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTTAATATTTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTTCGCGG
CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAACAACCTGGAGCTGATGTGAGACATGATATACCAGTGTGCCAA
AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA
TGTCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT

301 AATCATGCAGAGCTTCTGTGTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT

351 TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTTCATCCTGACA
ACACCAGCCGATGGCAGCACCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAAT
TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTTA

451 CGATATACATTTCGCCCTTTGGTGGTAATTATGATAGACTTGAACAACCTGC
GCTATATGTAAGCGGAAACCACCATTAATACTATCTGAACTTGTTGAACG

501 TGGTAATCTGAGAGAAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA

601 CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTCAGAAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCTGTTT

651 ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA
TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAATGTGAACCTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGAGGTTGGT

901 TCGTCACAGTTTCTACGTATTTACAGGCATCGGAGAAATTTAAGAATGC
AGCAGTGTCAAAGATGCATAAATGTCCGTAGCCTCTTTAAATTCCTTACG

66/254

FIGURE 13D (CONT'D)

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAACGCAATA
CAGATACACAACCTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTACGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCTTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCAACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTGTTACAACCATTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTACATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTGGCTCTATTAAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT


1701 TGGTGACCCAAACCAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCTCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

67/254

FIGURE 14A

SUBSTITUTE SHEET (RULE 26)

68/254

FIGURE 14B**WT preprorin linker**

primer VZV-A1

5' - GTGGAGGCAAGTTCTTAATGCTGATGTTTGT -3'

TCTTTGCTTATAAGGCCAGTGTGCGCCAAATTTTAAT
 AGAACGAAATATTCGGT CACCACGGTTAAATTA
 3' - AGCAGTGTCAAAAGAGTCCTACATTTGCGT -5'

primer VZV-A2


PCR mutagenesis

ligate with pBluescript SK

pAP 237 linker
 (VZV-A variant)

TCTCAGGATGTAAACGCCAGTGGAGGCAAGTTCTAAT
 AGAGTCCCTACATTTGCGTCACCTCCGTTCAAGATTA

69/254

FIGURE 14C**SUBSTITUTE SHEET (RULE 26)**

70/254

FIGURE 14D

10 20 30 40 50

1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCACT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC

101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTTAATATTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTGCGGG
CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAACTGAGCTGATGTGAGACATGATATACCAGTGTGCGCAA
AGCAAATGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGGTTTGCTATAAAACCAACGGTTTATTTTAGTTGAACTCTCA
TGTCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT

301 AATCATGCAGAGCTTCTGTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT

351 TGTGGTTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA
ACACCAGCCGATGGCAGACCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAAT
TAGTCCCTTACGTCTTCGTTAGTGAGTAGAAAAGTGAAGTACAAGTTT

451 CGATATACATTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACCTGCG
GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACTTGTGAAACG

501 TGGTAATCTGAGAGAAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTCGCGAAATAAATGTCATGACCACCGTGAGTCGAAGGTTGA

601 CTGGCTCGTTCCCTTTATAATTTGCATCCAAATGATTTTCAAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTT

651 ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA
TAAGGTTATATACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAATGTGAAGTCTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTTCTCAGGATGTAAACGCAGTGGAGGCAAGTTCTAATGC
AGCAGTGTCAAAAGAGTCCCTACATTTGCGTCACCTCCGTTCAAGATTACG

SUBSTITUTE SHEET (RULE 26)

71/254

FIGURE 14D (CONT'D)

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAACGCAATA
CAGATACACAACCTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTACAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCAACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTGTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTAAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT


1701 TGGTGACCCAAACCAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCACTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

72/254

FIGURE 15A

SUBSTITUTE SHEET (RULE 26)

73/254

FIGURE 15B

WT preprorin linker

primer VZV-B1

5' - TCGACGGGATATGGTAATGCTGATGTTGT -3'

TCTTTGCTTATAAGGCCAGTGGTGCCAAATTTTAAT
 AGAAACGAATATTCCCGT CACCACGGTTAAATTA

3' - AGCAGTGTCAAAAAGACACATAAAATGTCCGT -5'

primer VZV-A2


PCR mutagenesis

ligate with pBluescript SK

pAP 239 linker
(VZV-B variant)

TCTGTGTATTTACAGGCATCGACGGGATATGGTAAT
 AGACACATAAATGTCCGTAGCTGCCCTATACCATTA

74/254

FIGURE 15C

SUBSTITUTE SHEET (RULE 26)

75/254

FIGURE 15D

10 20 30 40 50
1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA
51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTAAATC
101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTTAATATTTGAAATGGTGT
151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTTCGCGG
CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC
201 TCGTTTAACAACTGGAGCTGATGTGAGACATGATATACCAGTGTGCGCAA
AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT
251 ACAGAGTTGGTTTGCTATAAACCACGGTTTATTTTAGTTGAACTCTCA
TGTCTCAACCAAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT
301 AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT
351 TGTGGTTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTTCATCCTGACA
ACACCAGCCGATGGCACGACCTTTATCGCGTATAAAGAAAGTAGGACTGT
401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAAT
TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTTA
451 CGATATACATTGCGCTTTGGTGGTAATTATGATAGACTTGAACAACCTGCG
GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACCTGTTGAACG
501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC
551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA
601 CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTTCAAGAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT
651 ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA
TAAGGTTATATACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT
701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAATGTGAACCTTATCAACCCCTCT
751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA
801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTAAGTCACACATGCTACACTCAT
851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGAGGTGGT
901 TCGTCACAGTTTCTGTGTATTTACAGGCATCGACGGGATATGGTAATGC
AGCAGTGTCAAAAGACACATAAATGTCCGTAGCTGCCCTATACCATTACG

SUBSTITUTE SHEET (RULE 26)

76/254

FIGURE 15D (CONT'D)

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACCTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTACAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTGAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACAACCATTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTGTGTCACCCGAGAAATACGCTTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAATTTAAACATATCACCTAACCACAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT

1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAGTGTGTGTGTCTCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT


1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

77/254

FIGURE 16A

PCR Mutagenesis of Preproricin Gene to Create an EBV-A Variant Gene
a) Cloning Strategy



SUBSTITUTE SHEET (RULE 26)

78/254

FIGURE 16B

WT preprorinclin linker

primer EBV-A1

5' - TCGGCGTCAGGTGTTAATGCTGATGTTTGT -3'

TCTTTGCTTATAAGGCCAGTGGTCCCAAATTTTAAT

AGAAACGAATATTCCTGGTCACCCACGGTTTAAATTA

3' - AGCAGTGTCAAAAGATTTCGAACATGTCCGT -5'

primer EBV-A2

PCR mutagenesis


ligate with pBluescript SK

pAP 241 linker
(EBV-A variant)

TCTAAGCTTGACAGGCATCGGCGTCAGGTGTTAAT

AGATTGGAACATGTCCGTAGCCGCGAGTCCACAATTA

79/254

FIGURE 16C**SUBSTITUTE SHEET (RULE 26)**

80/254

FIGURE 16D

10 20 30 40 50

1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCACT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTAAATC

101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTTAATATTTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTTCGCGG
CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAACTGAGCTGATGTGAGACATGATATACCAAGTGTGCGCA
AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGGTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA
TGTCTCAACCAAACGGATATTGGTTGCCAAATAAAATCAACTTGAGAGT

301 AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT

351 TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTTCATCCTGACA
ACACCAGCCGATGGCAGACCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT
TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTT

451 CGATATACATTTCGCTTTTGGTGGTAATTATGATAGACTTGAACAACCTGC
GCTATATGTAAGCGGAAACCACCATTAATACTATCTGAACCTGTTGAACG

501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCTTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA

601 CTGGCTCGTTCCCTTTATAATTTGCATCCAAATGATTTCAGAAGCAGCAAG
GACCGAGCAAGGAAATATTAACGTAGGTTTACTAAAGTCTTCGTCGTT

651 ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA
TAAGGTTATATACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAATGTGAACTCTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTTCTAAGCTTGACAGGCATCGGCGTCAGGTGTTAATGC
AGCAGTGTCAAAGATTGGAACATGTCCGTAGCCGCAGTCCACAATTACG

81/254

FIGURE 16D (CONT'D)

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACACTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTCAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCAACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTGTTACAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTGTCACCCGAGAAATACGTTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTGTGGCTCTATTAAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAATTTAAACATATCACCTAACCACAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT


1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

82/254

FIGURE 17A

SUBSTITUTE SHEET (RULE 26)

83/254

FIGURE 17B**WT preprorinclin linker****primer EBV-B1**

5'- TCGGACGCACCTGATAATGCTGATGTTGT -3'

TCTTTCCTTATAAGGCCAGTGGTGCCAAATTTAAT
 AGAAACGAATATTCCGGTCACCCACGGTTAAATTA

3'- AGCAGTGTCAAAAAGCATAGATTTCCT -5'

primer EBV-B2


PCR mutagenesis

ligate with pBluescript SK

**pAP 243 linker
(EBV-B variant)**

TCTTCGTATCTAAAGGCATCGGACGCACCTGATAAT
 AGAAGCATAGATTTCCTAGCCTGCGTGGACTATTA

84/254

FIGURE 17C

SUBSTITUTE SHEET (RULE 26)

85/254

FIGURE 17D

10 20 30 40 50

1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCACT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTAAATC

101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTAAATATTTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTGCGGG
CGCCACGGTGACACGTTTCGATGTGTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAACTGAGCTGATGTGAGACATGATATACCAGTGTGCCAA
AGCAAAATGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGGTTTGCCCTATAAACCACGGTTTATTTTAGTTGAACCTCTCA
TGCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT

301 AATCATGCAGAGCTTCTGTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAAATGTAATCGCGACCTACAGTGGTTACGTAT

351 TGTGGTCGGCTACCGTGCTGGAATAGCGCATATTTCTTTCATCCTGACA
ACACCAGCCGATGGCACGACCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAAT
TAGTCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGAAGTACAAGTTTCA

451 CGATATACATTGCGCTTTGGTGGTAATTATGATAGACTTGAACAACCTGCG
GCTATATGTAAGCGGAAACCACCATTAATACTATCTGAACCTGTTGAACG

501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCTTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA

601 CTGGCTCGTTCCCTTTATAATTTGCATCCAAATGATTTCAGAAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTCA

651 ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA
TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAAATGTGAACCTTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTTCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTCTTCGTATCTAAAGGCATCGGACGCACCTGATAATGC
AGCAGTGTCAAAAGAAGCATAGATTTCCGTAGCCTGCGTGGAAGTATTACG

86/254

FIGURE 17D (CONT'D)

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCATATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAACGCAATA
CAGATACACAACCTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTACGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAATCGTCGCTGTAGTCCCTTGTCAACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCAAGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACAAACATTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT


1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAAGTGTGTGTCTGTCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAG

1851 TGCAG
ACGTC

87/254

FIGURE 18A

SUBSTITUTE SHEET (RULE 26)

88/254

FIGURE 18B

WT preprorin linker

primer HCMV-A1

5' - TCGGTAGACTTGCCTAATGCTGATGTTGT -3'

TCTTTGCTTATAAGGCCA
AGAAACGAATATTCGGT

3' - AGCAGTGTCAAAAGACCCCAACATTTACGT-5'

primer HCMV-A2


PCR mutagenesis

ligate with pBluescript SK

pAP 245 linker
(HCMV-A variant)

TCTGGGGTGTAAATGCATCGGTAGACTTGCTAAT
AGACCCCAACATTTACGTAGCACATCTGAACGATTA

89/254

FIGURE 18C**SUBSTITUTE SHEET (RULE 26)**

90/254

FIGURE 18D

10 20 30 40 50

1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC

101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTAAATATTTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTTCGCGG
CGCCACGGTGACACGTTTCGATGTGTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAACAACCTGGAGCTGATGTGAGACATGATATACCAGTGTGCCAA
AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGGTTTGCCTATAAAACCAACGGTTTATTTTAGTTGAACTCTCA
TGTCTCAACCAACGGATATTTGGTTGCCAAATAAATCAACTTGAGAGT

301 AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT

351 TGTGGTGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA
ACACCAGCCGATGGCACGACCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAAT
TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGAATAAGTTTTA

451 CGATATACATTGCGCTTTGGTGGTAATTATGATAGACTTGAACAACCTTGC
GCTATATGTAAGCGGAAACCACCATTAATACTATCTGAACCTTGTGTAACG

501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTCCGGAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA

601 CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTGAGAAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT

651 ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA
TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAAATGTGAACCTTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTTCTGGGGTTGTAAATGCATCGTGTAGACTTGCTAATGC
AGCAGTGTCAAAGACCCCAACATTTACGTAGCACATCTGAACGATTACG

SUBSTITUTE SHEET (RULE 26)

91/254

FIGURE 18D (CONT'D)

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCCTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAACGCAATA
CAGATACACAACACTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTACAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACAAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAATTTAAACATATCACCTAACCACAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT


1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAAGTGTGTGTCTCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

92/254

FIGURE 19A

SUBSTITUTE SHEET (RULE 26)

93/254

FIGURE 19B

WT preprotrichin linker

primer HCMV-B1

5' - TCGGTGTCACCTGAAAAATGCTGATGTTTGT - 3'

TCTTTGCTTATAAGGCCAGTGGTGCCAAATTTTAAT
 AGAAACGGAATATTCCGGTCACCCACGGTTTAAATTA

3' - AGCAGTGTCAAAAGAAGCATACATTTCGGT - 5'

primer HCMV-B2


PCR mutagenesis

ligate with pBluescript SK

pAP 247 linker
(HCMV-B variant)

TCTTCGTATGTAAGGCATCGGTGTCACCTGAAAAAT
 AGAAGCATACATTTCGGTAGCCACAGTGGACTTTTA

94/254

FIGURE 19C

SUBSTITUTE SHEET (RULE 26)

95/254

FIGURE 19D

10 20 30 40 50

1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC

101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTTAATATTTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTGCGGG
CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAACAACCTGGAGCTGATGTGAGACATGATATACCAGTGTGCCAA
AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA
TGTCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT

301 AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT

351 TGTGGTGGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA
ACACCAGCCGATGGCAGGACCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAAT
TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTTA

451 CGATATACATTGCGCTTTGGTGGTAATTATGATAGACTTGAACAACTTGC
GCTATATGTAAGCGGAAACCACCATTAATACTATCTGAACTTGTGTAACG

501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA

601 CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTTCAAGAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT

651 ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA
TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAAATGTGAACTCTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTTCTTCGTATGTAAAGGCATCGGTGTCACCTGAAAATGC
AGCAGTGTCAAAGAAGCATACATTTCCGTAGCCACAGTGGACTTTTACG

SUBSTITUTE SHEET (RULE 26)

96/254

FIGURE 19D (CONT'D)

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACCTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTACGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTACGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAATCGTCGCTGTAGTCCCTTGTACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTGTTACAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTGGCTCTATTAAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT


1701 TGGTGACCCAAACCAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCTGTGCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

97/254

FIGURE 20A

SUBSTITUTE SHEET (RULE 26)

98/254

FIGURE 20B

WT prepronicin linker

primer HHV6-A1

5' - TCGGTGCCAAATTTTAAT - 3'

TCTTTGCTTATAAGGCCAGTGGTGCCAAATTTTAAT
AGAAACGAAATATCCGGTCACCCACGGTTAAATTA
3' - AGCAGTGTCAAAAGCTAAAATTTACGT - 5'

primer HHV6-A2


PCR mutagenesis

ligate with pBluescript SK

pAP 249 linker
(HHV-6 variant)

TCTTCGATTTTAAATGCATCGGTGCCAAATTTTAAT
AGAAGCTAAAATTTACGTAGCCACGGTTAAATTA

99/254

FIGURE 20C

SUBSTITUTE SHEET (RULE 26)

100/254

FIGURE 20D

10 20 30 40 50
1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA
51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGAATC
101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTTAATATTTGAAATGGTGT
151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTTCGCGG
CGCCACGGTGACACGTTTCGATGTGTTTGAATAGTCTCGACAAGCGCC
201 TCGTTTAACAACCTGGAGCTGATGTGAGACATGATATACCAGTGTGCGAA
AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT
251 ACAGAGTTGGTTTGCCTATAAACCACGGTTTATTTTAGTTGAACTCTCA
TGTCTCAACCAAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT
301 AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT
351 TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA
ACACCAGCCGATGGCACGACCTTTATCGCGTATAAAGAAAGTAGGACTGT
401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT
TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTTA
451 CGATATACATTTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACCTTGC
GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACTTGTGTAACG
501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCTTTACCAGGTGATCTCCTCC
551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA
601 CTGGCTCGTTCTTTTATAATTTGCATCCAAATGATTTCAGAAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT
651 ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA
TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT
701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAAATGTGAACTCTTATCAACCCCTCT
751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCTCGGAAACGATCAGGTTA
801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAGTCACACATGCTACACTCAT
851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGAGGTTGGT
901 TCGTCACAGTTTTCTTCGATTTTAAATGCATCGGTGCCAAATTTTAATGC
AGCAGTGTCAAAGAAGCTAAAATTTACGTAGCCACGGTTTAAAATTACG

SUBSTITUTE SHEET (RULE 26)

101/254

FIGURE 20D (CONT'D)

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACCTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTACAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAATCGTTCGCTGTAGTCCCTTGTCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACAACCATTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCCTCAG
TCCGACTTGTGTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAATTTAAACATATCACCTAACCAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCCTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT

1701 TGGTGACCCAAACCAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCCTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

SUBSTITUTE SHEET (RULE 26)

102/254

FIGURE 21

Ricin linker (wild type):

A chain- S L L I R P V V P N F N -B chain

pAP-213/pAP-214 linker (Cathepsin B):

A chain- S L L K S R M V P N F N -B chain

pAP-215/pAP-216 linker (MMP-3):

A chain- R P K P Q Q F F G L M N -B chain

pAP-217/pAP-218 linker (MMP-7):

A chain- S L R P L A L W R S F N -B chain

pAP-219/pAP-220 linker (MMP-9):

A chain- S P Q G I A G Q R N F N -B chain

pAP-221/pAP-222 linker (THERMOLYSIN-LIKE MMP):

A chain- D V D E R D V R G F A S F L -B chain

pAP-241/pAP-242 linker (EBV-A):


A chain- S K L V Q A S A S G V N -B chain

pAP-243/pAP-244 linker (EBV-B):

A chain- S S Y L K A S D A P D N -B chain

SUBSTITUTE SHEET (RULE 26)

103/254

FIGURE 22A

SUBSTITUTE SHEET (RULE 26)

104/254

FIGURE 22B

WT preprorin linker

primer ILTV-B1

5' - AATGAGGTAATTACTAATGCTGATGTTTGT - 3'

TCTTTGCTTATAAGGCCAGTGGTGCCCAATTTTAAT
AGAAACGAATATTCCGGTCACCCAGGTTAAATTA

3' - AGCAGTGTCAAAAGATTTCATAGATGTCCGT - 5'

primer ILTV-B2


PCR mutagenesis

ligate with pBluescript SK

pAP 253 linker
(ILTV-B variant)

TCTAAGTATCTACAGGCAATGAGGTAATTACTAAT
AGATTCATAGATGTCCGGTTACTCCATTAAATGATTA

105/254

FIGURE 22C

SUBSTITUTE SHEET (RULE 26)

106/254

FIGURE 22D

10 20 30 40 50

1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAACCTAGGTGGAGTCCCACCAGAAAGTGTAACT

101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTAAATATTTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTGCGCG
GCCCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAACAACCTGGAGCTGATGTGAGACATGATATACCAGTGTGCGCAA
AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACCTCTCA
TGTCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT

301 AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT

351 TGTGGTGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA
ACACCAGCCGATGGCACGACCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAAT
TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGAATACAAGTTTTA

451 CGATATACATTGCGCTTTGGTGGTAATTATGATAGACTTGAACAACCTTGC
GCTATATGTAAGCGGAAACCACCATTAACTACTATCTGAACTTGTGAAACG

501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTTCGGAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA

601 CTGGCTCGTTCCCTTTATAATTTGCATCCAAATGATTTGAGAAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT

651 ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA
TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAATGTGAACTCTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTTCTAAGTATCTACAGGCAAATGAGGTAATTACTAATGC
AGCAGTGTCAAAGATTCATAGATGTCCGTTTACTCCATTAATGATTACG

SUBSTITUTE SHEET (RULE 26)

107/254

FIGURE 22D (CONT'D)

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACCTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCCGGTACGTTTACAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCTTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCCTCAG
TCCGACTTGTTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT

1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA


1751 CTCTTGCAGTGTGTGTGTCTCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

108/254

FIGURE 23A



SUBSTITUTE SHEET (RULE 26)

109/254

FIGURE 23B

WT preprorin linker

primer HAV-A1

5' - TCGTTCTCAAAATGGAATGCTGATGTTTGT -3'

TCTTTGCTTATAAGGCCAGTGGTGCCAAATTTTAAT
 AGAACGAATATTCCGGTCACCCACGGTTTAAATTA

3' - AGCAGTGTCAAAAGACTCGAATCTTTGCGTT -5'

primer HAV-A2


PCR mutagenesis

ligate with pBluescript SK

 pAP 257 linker
 (HAV-A variant)

TCTGAGCTTAGAACGCAATCGTTCTCAAAATGGAAT
 AGACTCGAATCTTGGTTAGCAAGAGTTTAACTTA

110/254

FIGURE 23C

SUBSTITUTE SHEET (RULE 26)

111/254

FIGURE 23D

10 20 30 40 50

1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC

101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTTAATATTTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTGCGGG
CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAAACAACTGGAGCTGATGTGAGACATGATATACCAGTGTGCCAA
AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA
TGCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTGAGAGT

301 AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTACGTAT

351 TGTGGTGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA
ACACCAGCCGATGGCACGACCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAAT
TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGAAGTACAAGTTT

451 CGATATACATTGCGCTTTGGTGGTAATTATGATAGACTTGAACAACCTTGC
GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACTTGTGAAACG

501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCTTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA

601 CTGGCTCGTTTCCTTTATAATTTGCATCCAAATGATTTTCAAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT

651 ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA
TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAAATGTGAACTCTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTTCTGAGCTTAGAACGCAATCGTTCTCAAATTGGAATGC
AGCAGTGTCAAAAGACTCGAATCTTGCGTTAGCAAGAGTTTAACCTTACG

SUBSTITUTE SHEET (RULE 26)

112/254

FIGURE 23D (CONT'D)

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACACTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTCAATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAATCGTCGCTGTAGTCCCTTGTCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTGTTACAACCATTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT


1701 TGGTGACCCAAACCAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

113/254

FIGURE 24A

SUBSTITUTE SHEET (RULE 26)

114/254

FIGURE 24B

WT preprorin linker

primer HAV-B1

5' - GGGATCGATGATGATAATGCTGATGTTTGT -3'

TCTTTGCTTATAAGGCCAGTGGTGCCAAATTTTAAT
AGAAACGAATATTCCGGTCACCCACGGTTAAATTA

3' - AGCAGTGTCAAAAGACTCGAAACCCAGCGTT -5'


primer HAV-B2

PCR mutagenesis
ligate with pBluescript SK

pAP 255 linker
(HAV-B variant)

TCTGAGCTTTGGTCGCAAGGATCGATGATGATAAT
AGACTCGAAACCCAGCGTTCCCTAGCTACTACTATTA

115/254

FIGURE 24C

SUBSTITUTE SHEET (RULE 26)

116/254

FIGURE 24D

10 20 30 40 50

1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTAAATC

101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTTAATATTTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTTCGCGG
CGCCACGGTGACACGTTTCGATGTGTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAACAACCTGGAGCTGATGTGAGACATGATATACCAGTGTTCGCAA
AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGGTTTGCCTATAAAACCAACGGTTTATTTTAGTTGAACTCTCA
TGTCTCAACCAAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT

301 AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT

351 TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA
ACACCAGCCGATGGCAGACCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAAT
TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGAAGTACAAGTTT

451 CGATATACATTTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACTTGC
GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACTTGTGTAACG

501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA

601 CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTGAGAAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTT

651 ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA
TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAAATGTGAACCTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTTCCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTTCTGAGCTTTGGTCGCAAGGGATCGATGATGATAATGC
AGCAGTGTCAAAAGACTCGAAACCAGCGTTCCCTAGCTACTACTATTACG

SUBSTITUTE SHEET (RULE 26)

117/254

FIGURE 24D (CONT'D)

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAACGCAATA
CAGATACACAACACTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTACAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTTCAAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCAACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACAACCATGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTGTACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAATTTAAACATATCACCTAACCACAATCTA

1651 GTGAGGCGATCCGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT


1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

118/254

FIGURE 25A

SUBSTITUTE SHEET (RULE 26)

119/254

FIGURE 25B**WT preprorin linker**

primer CAN-1

5' - TTCAGGCTAAATTTAATGCTGAT -3'

TCTTTGCTTATAAGGCCAGTGGTGCCAAATTTTAAT
 AGAACGAATATTCCGGTCACCCACGGTTAAATTA
 3' - AGCAGTGTCAAAAGATTCCGACGTTTCAAG -5'

primer CAN-2


PCR mutagenesis

ligate with pBluescript SK

pAP 259 linker
(CAN variant)

TCTAAGCCTGCAAAAGTTCTTCAGGCTAAATTTTAAT
 AGATTCCGACGTTTCAAGAAGTCCGATTAAATTA

120/254

FIGURE 25C

SUBSTITUTE SHEET (RULE 26)

121/254

FIGURE 25D

10 20 30 40 50

1 GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51 GGCAACATGGCTTTGTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC

101 AGGATAACAACATATTCCCAAACAATACCCAATTATAAACTTTACCACA
TCCTATTGTTGTATAAGGGGTTTGTATGGGTAAATATTTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTTCGCGG
CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAACAACCTGGAGCTGATGTGAGACATGATATACCAGTGTTCGCAA
AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA
TGTCTCAACCAAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT

301 AATCATGCAGAGCTTCTGTACATTAGCGCTGGATGTCACCAATGCATA
TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT

351 TGTGGTTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA
ACACCAGCCGATGGCACGACCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT
TAGTCCCTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTTA

451 CGATATACATTTCGCCCTTTGGTGGTAATTATGATAGACTTGAACAACCTGC
GCTATATGTAAGCGGAAACCACCATTAATACTATCTGAACCTGTTGAACG

501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
ACCATTAGACTCTCTTTTATAGCTCAACCCCTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA

601 CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTTCAAGAGCAGCAAG
GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT

651 ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA
TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAATGTGAACCTTTATCAACCCCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGAGGTTGGT

901 TCGTCACAGTTTTCTAAGCCTGCAAAGTTCTTCAGGCTAAATTTTAATGC
AGCAGTGTCAAAAGATTCGGACGTTTCAAGAAGTCCGATTTAAAATTACG

SUBSTITUTE SHEET (RULE 26)

122/254

FIGURE 25D (CONT'D)

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACACTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTACAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAATCGTCGCTGTAGTCCCTTGTCCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTTACAACCATTTGTTGGGCTATATGGTCTGTG
TTATATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTACATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCCTCAG
TCCGACTTGTTGTACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTGGCTCTATTAAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCAACATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT

1701 TGGTGACCCAAACCAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

123/254

FIGURE 26

Ricin linker (wild type):

A chain- S L L I R P V V P N F N -B chain

pAP-223/224 linker (MAL-A):

A chain- Q V V Q L Q N Y D E E D -B chain

pAP-225/226 linker (MAL-B):

A chain- L P I F G E S E D N D E -B chain

pAP-227/228 linker (MAL-C):

A chain- Q V V T G E A I S V T M -B chain

pAP-229/230 linker (MAL-D):

A chain- A L E R T F L S F P T N -B chain

pAP-231/pAP-232 linker (MAL-E):

A chain- K F Q D M L N I S Q H Q -B chain

124/254

FIGURE 27

Ricin linker (wild type):

A chain- S L L I R P V V P N F N -B chain

pAP-245/246 linker (CMV-A):

A chain- S G V V N A S C R L A N -B chain

pAP-247/248 linker (CMV-B):

A chain- S S Y V K A S V S P E N -B chain

pAP-233/234 linker (HERPES SIMPLEX-1 A):

A chain- S A L V N A S S A H V N -B chain

pAP-235/236 linker (HERPES SIMPLEX-1 B):

A chain- S T Y L Q A S E K F K N -B chain

pAP-249/250 linker (HUMAN HERPES VIRUS-6):

A chain- S S I L N A S V P N F N -B chain

pAP-237/pAP-238 linker (VZV-A):

A chain- S Q D V N A V E A S S N -B chain

pAP-239/pAP-240 linker (VZV-B):

A chain- S V Y L Q A S T G Y G N -B chain

pAP-253/pAP-254 linker (ILV):

A chain- S K Y L Q A N E V I T N -B chain

pAP-255/pAP-256 linker (HAV-A):

A chain- S E L R T Q S F S N W N -B chain

pAP-257/pAP-258 linker (HAV-B):

A chain- S E L W S Q G I D D D N -B chain

SUBSTITUTE SHEET (RULE 26)

125/254

FIGURE 28

Ricin linker (wild type):

A chain- S L L I R P V V P N F N -B chain

pAP-259/260 linker (CAP-A):

A chain- S K P A K F F R L N F N -B chain


pAP-261/262 linker (CAP-B):

A chain- S K P I E F F R L N F N -B chain


pAP-263/264 linker (CAP-C):

A chain- S K P A E F F A L N F N -B chain

126/254

FIGURE 29**PCR Mutagenesis of Preproricin Gene to Create A Variant Gene in Baculovirus Transfer Vector, pVL 1393****a) Cloning Strategy**

127/254

FIGURE 30A**PCR Mutagenesis of Preprorincin Gene to Create An HCV-A Variant Gene in Baculovirus Transfer Vector, pVL 1393****a) Cloning Strategy**

SUBSTITUTE SHEET (RULE 26)

128/254

FIGURE 30B**Sequence of HCV-A Linker Region****WT preprorincin linker****primer HCV-A1**

5' - TCGACATGGGTTTAAATGCTGATGTT -3'

TCTTGCTTATAAGGCCAGTGGTGCCAAATTTTAAT

AGAAACGAATATTCGGGTCAACACGGTTTAAATTA

3' - GGTAGCAGTGTCAAACTAAACCTCCATCACTGT 5'

5' primer HCV-A2↓
PCR mutagenesis↓
ligate with pVL1393**pAP 262 linker
(HCV-A variant)**

GATTTGGAGGTAGTGACATCGACATGGGTTTAAAT

CTAAACCTCCATCACTAGCTGTACCCAAAATTA

129/254

FIGURE 30C (P1)

Sequence of pAP262 insert

	10	20	30	40	50
1	GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT				
	CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA				
51	GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG				
	CCGTTGTACCGAAACAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC				
101	AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA				
	TCCTATTGTTGTATAAGGGGTTTGTATGGGTTAATATTTGAAATGGTGT				
151	GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTTCGCGG				
	CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC				
201	TCGTTTAAACAACTGGAGCTGATGTGAGACATGATATACCAAGTGTGCCAA				
	AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT				
251	ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA				
	TGTCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT				
301	AATCATGCAGAGCTTCTGTACATTAGCGCTGGATGTCACCAATGCATA				
	TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT				
351	TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA				
	ACACCAGCCGATGGCAGACCTTTATCGCGTATAAAGAAAGTAGGACTGT				
401	ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT				
	TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGAATAAGTTT				
451	CGATATACATTTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACCTGC				
	GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACCTGTTGAACG				
501	TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG				
	ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC				
551	CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC				
	GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA				
601	CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTCAGAAGCAGCAAG				
	GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTT				
651	ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA				
	TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT				
701	GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA				
	CTAGACGTGGTCTAGGATCGCATTAAATGTGAACCTTTATCAACCCCTCT				
751	CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT				
	GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA				

SUBSTITUTE SHEET (RULE 26)

130/254

FIGURE 30C (P2)

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTGATTTGGAGGTAGTGACATCGACATGGGTTTTTAATGC
AGCAGTGTCAAACCTAAACCTCCATCACTGTAGCTGTACCCAAAAATTACG

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACCTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTCAAGTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTTACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAATCGTCGCTGTAGTCCCTTGTCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTTACAACCATGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTACATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA

131/254

FIGURE 30C (P3)

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT

1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCA GTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAAGTAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

Total number of bases is: 1855.

Sequence name: PAP262

132/254


FIGURE 30D

**-Amino Acid Sequence Comparison of Mutant
Preproricin Linker Region of HCV-A to Wild Type**

Wild type Ricin linker: A chain- S L L I R P V V P N F N -B chain

pAP-262 linker: A chain- D L E V V T S T W V F N -B chain
(HCV-A linker)

133/254

FIGURE 31A**PCR Mutagenesis of Preprorincin Gene to Create An HCV-B Variant Gene in Baculovirus Transfer Vector, pVL 1393****a) Cloning Strategy**

SUBSTITUTE SHEET (RULE 26)

134/254

FIGURE 31B**Sequence of HCV-B Linker Region****WT preproridin linker****primer HCV-B1**

5' - GCGTCACACCTTTTAAATGCTGATGTT -3'

TCTTTGCTTATAAGGCCAGTGGTGCCAAATTTTAAT
 AGAAACGAATATTCGGGTCACCCACGGTTAAAATTA

3' - GGTAGCAGTGTCAAACACTCTACCTTCTCACA-5'

5' primer HCV-B2

PCR mutagenesis

ligate with pVL1393

**pAP 264 linker
(HCV-B variant)**

GATGAGATGGAAGAGTGTGGTGCACACCTTTTAAAT
 CTACTCTACCTTCTCACAACGCGAGTGTGGAAAATTA

135/254

FIGURE 31C (P1)

Sequence of pAP264 insert

	10	20	30	40	50
1	GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT				
	CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA				
51	GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG				
	CCGTTGTACCGAAACAAACCTAGGTGGAGTCCCACCAGAAAGTGAATC				
101	AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA				
	TCCTATTGTTGTATAAGGGGTTTGTTATGGGTAAATATTTGAAATGGTGT				
151	GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTGCGGG				
	CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC				
201	TCGTTTAACAACTGGAGCTGATGTGAGACATGATATACCACTGTTGCCAA				
	AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT				
251	ACAGAGTTGGTTTGCCTATAAAACCAACGGTTTATTTTAGTTGAACTCTCA				
	TGTCTCAACCAAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT				
301	AATCATGCAGAGCTTTCTGTTCATTAGCGCTGGATGTCACCAATGCATA				
	TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT				
351	TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA				
	ACACCAGCCGATGGCACGACCTTTATCGCGTATAAAGAAAGTAGGACTGT				
401	ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT				
	TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTTA				
451	CGATATACATTTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACCTTGC				
	GCTATATGTAAGCGGAAACCACCATTAATACTATCTGAACCTTGTGAACG				
501	TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG				
	ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC				
551	CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC				
	GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA				
601	CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTCAGAAGCAGCAAG				
	GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT				
651	ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA				
	TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT				
701	GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA				
	CTAGACGTGGTCTAGGATCGCATTAAATGTGAACCTTTATCAACCCCTCT				
751	CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT				
	GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA				

SUBSTITUTE SHEET (RULE 26)

136/254

FIGURE 31C (P2)

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTGATGAGATGGAAGAGTGTGCGTCACACCTTTTTAATGC
AGCAGTGTCAAACCTACTCTACCTTCTCACACGCAGTGTGGAAAATTACG

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTGCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACCTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTACGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTACAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAATCGTCGCTGTAGTCCCTTGTACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTTACAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTACATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA

137/254

FIGURE 31C (P3)

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT

1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

Total number of bases is: 1855.

Sequence name: PAP264

138/254

FIGURE 31D

**-Amino Acid Sequence Comparison of Mutant
Preproricin Linker Region of HCV-B to Wild Type**


Wild type Ricin linker: A chain- S L L I R P V V P N F N -B chain

pAP-264 linker: A chain- D E M E E C A S H L F N -B chain
(HCV-B linker)

139/254

FIGURE 32A

- PCR Mutagenesis of Preprorincin Gene to Create An HCV-C Variant Gene in Baculovirus Transfer Vector, pVL 1393

a) Cloning Strategy

SUBSTITUTE SHEET (RULE 26)

140/254

FIGURE 32B**Sequence of HCV-C Linker Region****WT preproreicin linker**

primer HCV-C1

5' - TCGATGTCATATTTTAATGCTGATGTT - 3'

TCTTTGCTTATAAGGCCAGTGGTGCCAAATTTTAAT
 AGAAACGAAATATTCGGT CACCACGGTTAAATA

3' - GGTAGCAGTGTCAAACTCCTGCAACATACAACA - 5'

5' primer HCV-C2

PCR mutagenesis

ligate with pVL1393

pAP 266 linker
 (HCV-C variant)

GAGGACGTTGTATGTTGTTTCGATGTCATATTTTAAT
 CTCCTGCAACATACAACAAGCTACGTACGTATAAAATA

141/254

FIGURE 32C (P1)

Sequence of pAP266 insert

	10	20	30	40	50
1	GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT				
	CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA				
51	GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG				
	CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC				
101	AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA				
	TCCTATTGTTGTATAAGGGGTTTGTATGGGTAAATATTTGAAATGGTGT				
151	GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTTCGCGG				
	CGCCACGGTGACACGTTTCGATGTGTTGAAATAGTCTCGACAAGCGCC				
201	TCGTTTAACAACTGGAGCTGATGTGAGACATGATATACCAGTGTTGCCAA				
	AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT				
251	ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA				
	TGTCTCAACCAAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT				
301	AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA				
	TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT				
351	TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA				
	ACACCAGCCGATGGCAGCACCTTTATCGCGTATAAAGAAAGTAGGACTGT				
401	ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT				
	TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTTA				
451	CGATATACATTTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACTTGC				
	GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACCTTGTTGAACG				
501	TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG				
	ACCATTAGACTCTCTTTTATAGCTCAACCCTTACCAGGTGATCTCCTCC				
551	CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAT				
	GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA				
601	CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTCAGAAGCAGCAAG				
	GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT				
651	ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA				
	TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT				
701	GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA				
	CTAGACGTGGTCTAGGATCGCATTAAATGTGAACTCTTATCAACCCCTCT				
751	CTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT				
	GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA				

142/254

FIGURE 32C (P2)

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTGAGGACGTTGTATGTTGTTTCGATGTCATATTTTAATGC
AGCAGTGTCAAACCTCCTGCAACATACAACAAGCTACAGTATAAAATTACG

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACATAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTCAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAATCGTCGCTGTAGTCCCTTGTCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTTACAACCATTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTGGCTCTATTAAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCAACATCTA

143/254

FIGURE 32C (P3)

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT

1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCA GTGTGTGTCTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

Total number of bases is: 1855.

Sequence name: PAP266

144/254


FIGURE 32D

**-Amino Acid Sequence Comparison of Mutant
Preproricin Linker Region of HCV-C to Wild Type**

Wild type Ricin linker: A chain- S L L I R P V V P N F N -B chain

pAP-266 linker: A chain- E D V V C C S M S Y F N -B chain
(HCV-C linker)

145/254

FIGURE 33A**PCR Mutagenesis of Preproricin Gene to Create An HCV-D Variant Gene in Baculovirus Transfer Vector, pVL 1393****a) Cloning Strategy**

SUBSTITUTE SHEET (RULE 26)

146/254

FIGURE 33B**Sequence of HCV-D Linker Region****WT preproridin linker**

primer HCV-D1

5' - GGGCCCAATAACTGCTTATGCTGATGTTGTATG -3'

TCTTTGCTTATAAGGCCAGTGGTGCCAAATTTTAAT
 AGAAACGAAATATTCCGGTACACCACGGTTTAAATTA

3' - GGTAGCAGTGTCAAATTCCCCACCTCTAACGAT -5'

5' primer HCV-D2

PCR mutagenesis

ligate with pVL1393

**pAP 268 linker
(HCV-D variant)**

AAGGGTGGAGATTGCTAGGCCCAATAACTGCTTAT
 TTCCCCACCTCTAACGATCGCGGTATTGACGAATA

147/254

FIGURE 33C (P1)

Sequence of pAP268 insert

	10	20	30	40	50
1	GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT				
	CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA				
51	GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG				
	CCGTTGTACCGAAACAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC				
101	AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA				
	TCCTATTGTTGTATAAGGGGTTTGTATGGGTAAATATTTGAAATGGTGT				
151	GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTTCGCGG				
	CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC				
201	TCGTTTAACAACTGGAGCTGATGTGAGACATGATATACCAGTGTTGCCAA				
	AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT				
251	ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA				
	TGTCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT				
301	AATCATGCAGAGCTTCTGTACATTAGCGCTGGATGTCACCAATGCATA				
	TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT				
351	TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA				
	ACACCAGCCGATGGCAGACCTTTATCGCGTATAAAGAAAGTAGGACTGT				
401	ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT				
	TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTTA				
451	CGATATACATTTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACCTGC				
	GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACTTGTTGAACG				
501	TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG				
	ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC				
551	CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC				
	GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA				
601	CTGGCTCGTTTCCTTTATAATTTGCATCCAAATGATTTCAGAAGCAGCAAG				
	GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT				
651	ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA				
	TAAGGTTATATACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT				
701	GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA				
	CTAGACGTGGTCTAGGATCGCATTAATGTGAACTCTTATCAACCCCTCT				
751	CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT				
	GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA				

148/254

FIGURE 33C (P2)

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTAAGGGGTGGAGATTGCTAGCGCCAATAACTGCTTATGC
AGCAGTGTCAAATCCCCACCTCTAACGATCGCGGTTATTGACGAATACG

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCCTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACATAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTCAATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAATCGTCGCTGTAGTCCCTTGTCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA

149/254

FIGURE 33C (P3)

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT
1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA
1751 CTCTGCAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT
1801 GGACATTGTAAATTTTGTAAGTAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG
1851 TGCAG
ACGTC

Total number of bases is: 1855.

Sequence name: PAP268

150/254


FIGURE 33D

**-Amino Acid Sequence Comparison of Mutant
Preproricin Linker Region of HCV-D to Wild Type**

Wild type Ricin linker: A chain- S L L I R P V V P N F N -B chain

pAP-268 linker:
(HCV-D linker) A chain- K G W R L L A P I T A Y -B chain

151/254

FIGURE 34A**PCR Mutagenesis of Preproricin Gene to Create A Variant Gene in Baculovirus Transfer Vector, pVL 1393****a) Cloning Strategy**

SUBSTITUTE SHEET (RULE 26)

152/254

FIGURE 34B**Sequence of MMP-2 Linker Region****WT preprocin linker**

```
primer 270-3'
5' - TGGGCTCCTAATTTTAATGCTGATGTTTGT -3'
      | **  **  *
-----TCTTTGCTTATAAGGCCA|GTGGTACCAAATTTTAAT-----
-----AGAAACGAATATTCCGGT|CACCATGGTTTAAAATTA-----
      *** ** ***
3' -AGCAGTGTCAAAAGAAACGGGGACCCAAAT -5'
primer 270-5'
```

1) PCR mutagenesis

2) Ligate with pVL1393

**pAP 270 linker
(MMP-2 variant)**

```
-----TCTTTGCCCCCTGGGTTTA|TGGGCTCCTAATTTTAAT-----
-----AGAAACGGGGACCCAAAT|ACCCGAGGATTAAAATTA-----
```

153/254

FIGURE 34C (P1)

Sequence of pAP270 insert

	10	20	30	40	50
1	GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT				
	CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA				
51	GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG				
	CCGTTGTACCGAAACAAACCTAGGTGGAGTCCCACCAGAAAGTGTAAATC				
101	AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA				
	TCCTATTGTTGTATAAGGGGTTTGTATGGGTAAATATTTGAAATGGTGT				
151	GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTGCGCG				
	CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC				
201	TCGTTTAACAACTGGAGCTGATGTGAGACATGATATAACAGTGTTGCCAA				
	AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT				
251	ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA				
	TGTCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT				
301	AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA				
	TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT				
351	TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA				
	ACACCAGCCGATGGCACGACCTTTATCGCGTATAAAGAAAGTAGGACTGT				
401	ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAAT				
	TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGAATAAAGTTTGA				
451	CGATATACATTTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACTTGC				
	GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACTTGTTGAACG				
501	TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG				
	ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC				
551	CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC				
	GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA				
601	CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTTCAAGCAGCAAG				
	GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT				
651	ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA				
	TAAGGTTATATACTCCCTCTTTACCGCGTCTCTTAATGCAATTTGGCCT				

SUBSTITUTE SHEET (RULE 26)

154/254

FIGURE 34C (P2)

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAAATGTGAACTCTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTTCTTTGCCCTGGGTTTATGGGCTCCTAATTTTAATGC
AGCAGTGTCAAAGAAACGGGGACCCAAATACCCGAGGATTAAAATTACG

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACACTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTCAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTACAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAATCGTCGCTGTAGTCCCTTGTCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACAAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT

155/254

FIGURE 34C (P3)

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT

1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAAGTAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

Total number of bases is: 1855.

Sequence name: PAP270


156/254

FIGURE 34D

**Amino acid sequence Comparison of Mutant Preproricin Linker
region of MMP-2 to Wild Type**

Wild type ricin linker:	A chain- S L L I R P V V P N F N -B chain
pAP-270 (MMP-2) linker:	A chain- S L P L G L W A P N F N -B chain

157/254

FIGURE 35A**PCR Mutagenesis of Preproricin Gene to Create A Variant Gene in Baculovirus Transfer Vector, pVL 1393****a) Cloning Strategy**

SUBSTITUTE SHEET (RULE 26)

158/254

FIGURE 35B**Sequence of Cathepsin B (Site 2) Linker Region****WT preprocin linker**

```
primer 272-3'
5' - AGGATGCCAAATTTTAATGCTGATGTTTGT -3'
      | * * * *
-----TCTTTGCTTATAAGGCCA|GTGGTACCAAATTTTAAT-----
-----AGAAACGAATATTCCGGT|CACCATGGTTTAAAATTA-----
      | * * * *
3' - AGCAGTGTCAAAGAAACGAATATCGATCT -5'
      primer 272-5'
```

1) PCR mutagenesis

2) Ligate with pVL1393

pAP 272 linker**(Cathepsin B Site 2 variant)**

```
-----TCTTTGCTTATAGCTAGA|AGGATGCCTAATTTTAAT-----
-----AGAAACGAATATCGATCT|TCCTACGGATTAAAATTA-----
```

159/254

FIGURE 35C (P1)

Sequence of pAP272 insert

	10	20	30	40	50
1	GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT				
	CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA				
51	GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG				
	CCGTTGTACCGAAACAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC				
101	AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA				
	TCCTATTGTTGTATAAGGGGTTTGTATGGGTAAATATTTGAAATGGTGT				
151	GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTGCGGG				
	CGCCACGGTGACACGTTTCGATGTGTTGAAATAGTCTCGACAAGCGCC				
201	TCGTTTAACAACTGGAGCTGATGTGAGACATGATATACCAGTGTTGCCAA				
	AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT				
251	ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA				
	TGTCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT				
301	AATCATGCAGAGCTTTCTGTACATTAGCGCTGGATGTCACCAATGCATA				
	TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT				
351	TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA				
	ACACCAGCCGATGGCACGACCTTTATCGCGTATAAAGAAAGTAGGACTGT				
401	ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT				
	TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTIONAAGTTT				
451	CGATATACATTGCGCTTTGGTGGTAATTATGATAGACTTGAACAACCTGC				
	GCTATATGTAAGCGGAAACCACCATTAATACTATCTGAACTTGTTGAACG				
501	TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG				
	ACCATTAGACTCTCTTTTATAGCTCAACCCTTACCAGGTGATCTCCTCC				
551	CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC				
	GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA				
601	CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTTCAAGAGCAGCAAG				
	GACCGAGCAAGGAAATATTAACGTAGGTTTACTAAAGTCTTCGTCGTTT				
651	ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA				
	TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT				

SUBSTITUTE SHEET (RULE 26)

160/254
FIGURE 35C (P2)

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAAATGTGAACTCTTATCAACCCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTTCTTTGCTTATAGCTAGAAGGATGCCTAATTTTAATGC
AGCAGTGTCAAAGAAAGGAATATCGATCTTCCTACGGATTAAAATTACG

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACCTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTCAAGATTATGTCTACGTTTGTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTGAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAATCGTCGCTGTAGTCCCTTGTCAACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTGTTACAACCATTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATAACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTGTTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT

161/254

FIGURE 35C (P3)

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT

1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAAGTAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAAACATTTAAACATTGACTTTCCTGTGCTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

Total number of bases is: 1855.

Sequence name: PAP272

162/254

FIGURE 35D


**Amino acid sequence Comparison of Mutant Preproricin Linker
region of Cathepsin B Site 2 to Wild Type**

Wild type ricin linker: A chain- S L L I R P V V P N F N -B chain
pAP-272 (Cathepsin B 2) linker: A chain- S L L I A R R M P N F N -B chain

163/254

FIGURE 36A

PCR Mutagenesis of Preproricin Gene to Create A Variant Gene in Baculovirus Transfer Vector, pVL 1393

a) Cloning Strategy

SUBSTITUTE SHEET (RULE 26)

164/254

FIGURE 36B**Sequence of Cathepsin L Linker Region****WT preprocin linker**

```

      primer 274-3'
      5' - TCATGGGCTAATTTTAATGCTGATGTTTGT -3'
            |***** *
-----TCTTTGCTTATAAGGCCA|GTGGTACCAAATTTTAAT-----
-----AGAAACGAATATTCCGGT|CACCATGGTTTAAAATTA-----
            *** **
3' -AGCAGTGTCAAAAGAAACGAATATAAGGCC -5'
      primer 274-5'

```

1) PCR mutagenesis

2) Ligate with pVL1393

pAP 274 linker**(Cathepsin L variant)**

```

-----TCTTTGCTTATATTCCGG|TCATGGGCTAATTTTAAT-----
-----AGAAACGAATATAAGGCC|AGTACCCGATTAAAATTA-----

```

165/254

FIGURE 36C (P1)

Sequence of pAP274 insert

```
      10      20      30      40      50
      |      |      |      |      |
1  GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
   CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51  GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
   CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC

101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
   TCCTATTGTTGTATAAGGGGTTGTTATGGGTTAATATTTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTTCGCGG
   CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAACTGAGCTGATGTGAGACATGATATACCAAGTGTGCCAA
   AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA
   TGTCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT

301 AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
   TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT

351 TGTGGTTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA
   ACACCAGCCGATGGCACGACCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAAT
   TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGAATAAAGTTTGA

451 CGATATACATTTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACTTGC
   GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACTTGTGTAACG

501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
   ACCATTAGACTCTCTTTTATAGCTCAACCCTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
   TATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA

601 CTGGCTCGTTCCTTTTATAATTTGCATCCAAATGATTTTCAAGAGCAGCAAG
   GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT

651 ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA
   TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT
```

166/254

FIGURE 36C (P2)

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAAATGTGAACTCTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTTCTTTGCTTATATTCCGGTCATGGGCTAATTTTAATGC
AGCAGTGTCAAAGAAAGGAATATAAGGCCAGTACCCGATTAAATACG

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACATAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTGAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAATCGTCGCTGTAGTCCCTTGTCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATAACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTGTTTATCACCTGTTACATACCTATCTCCTGACATCGTCACTTT

167/254

FIGURE 36C (P3)

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT

1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAAACTATCTGTCTAATGA

1751 CTCTTGCAAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAAGTAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

Total number of bases is: 1855.

Sequence name: PAP274


168/254

FIGURE 36D

**Amino acid sequence Comparison of Mutant Preproricin Linker
region of Cathepsin L to Wild Type**

Wild type ricin linker: A chain- S L L I R P V V P N F N -B chain
pAP-274 (Cathepsin L) linker: A chain- S L L I F R S W A N F N -B chain

169/254

FIGURE 37A**PCR Mutagenesis of Preproricin Gene to Create A Variant Gene in Baculovirus Transfer Vector, pVL 1393****a) Cloning Strategy**

170/254

FIGURE 37B**Sequence of Cathepsin D Linker Region****WT preprocin linker**

primer 276-3'

5' - ACTGTTATTGTTATCACCGCTGATGTTTGT -3'

| *** * * * * *

```

-----TCTTTGCTTATAAGGCCA|GTGGTACCAAATTTTAAT-----
-----AGATAACGAATATTCCGG|CACCATGGTTTAAATTA-----
          * * * * *

```

3' -AGCAGTGTCAAAAGACCACAACAGTAGCGA -5'

primer 276-5'

1) PCR mutagenesis

2) Ligate with pVL1393

pAP 276 linker**(Cathepsin D variant)**

```

-----TCTGGTGTGTCATCGCT|ACTGTTATTGTTATCACC-----
-----AGACCACAACAGTAGCGA|TGACAATAACAATAGTGG-----

```


171/254

FIGURE 37C (P1)

Sequence of pAP276 insert

	10	20	30	40	50
1	GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT				
	CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA				
51	GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG				
	CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTATC				
101	AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA				
	TCCTATTGTTGTATAAGGGGTTTGTATGGGTAAATATTTGAAATGGTGT				
151	GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTGCGGG				
	CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC				
201	TCGTTTAAACAACTGGAGCTGATGTGAGACATGATATACCACTGTTGCCAA				
	AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT				
251	ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA				
	TGTCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTGAGAGT				
301	AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA				
	TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT				
351	TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA				
	ACACCAGCCGATGGCACGACCTTTATCGCGTATAAAGAAAGTAGGACTGT				
401	ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAAT				
	TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTIONAAGTTT				
451	CGATATACATTGCGCTTTGGTGGTAATTATGATAGACTTGAACAACTTGC				
	GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACTTGTTGAACG				
501	TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG				
	ACCATTAGACTCTCTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC				
551	CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC				
	GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA				
601	CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTGAGAAGCAGCAAG				
	GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTT				
651	ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA				
	TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT				

172/254

FIGURE 37C (P2)

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAAATGTGAACTCTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTTCTGGTGTGTGTCATCGCTACTGTTATTGTTATCACCGC
AGCAGTGTCAAAGACCACAACAGTAGCGATGACAATAACAATAGTGGCG

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACACTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTCAAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAATCGTCGCTGTAGTCCCTTGTCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTTACAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTACATCTCCTGACATCGTCACTTT

173/254

FIGURE 37C (P3)

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTCAACCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT

1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAAGTGTGTGTCTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

Total number of bases is: 1855.

Sequence name: PAP276

174/254


FIGURE 37D

**Amino acid sequence Comparison of Mutant Preproricin Linker
region of Cathepsin D to Wild Type**

Wild type ricin linker: A chain- S L L I R P V V P N F N -B chain

pAP-276 (Cathepsin D) linker: A chain- S G V V I A T V I V I T -B chain

175/254

FIGURE 38A**PCR Mutagenesis of Preproricin Gene to Create A Variant Gene in Baculovirus Transfer Vector, pVL 1393****a) Cloning Strategy**

176/254

FIGURE 38B**Sequence of MMP-1 (Interstitial collagenase) Linker Region****WT preprocin linker**

primer 278-3'

5' - ATTTGGGGACAGTTTAATGCTGATGTTTGT -3'

* * * * *

```

-----TCTTTGCTTATAAGGCCA|GTGGTACCAAATTTTAAT-----
-----AGAAACGAATATTCCGGT|CACCATGGTTTAAAATTA-----

```

** * * * * *

3' -AGCAGTGTCAAAGAAACCCAGGAGTTCCG -5'

primer 278-5'

1) PCR mutagenesis

2) Ligate with pVL1393

pAP 278 linker**(MMP-1 variant)**

```

-----TCTTTGGGTCCTCAAGGC|ATTTGGGGACAGTTTAAT-----
-----AGAAACCCAGGAGTTCCG|TAAACCCCTGTCAAATTA-----

```

177/254

FIGURE 38C (P1)

Sequence of pAP278 insert

	10	20	30	40	50
1	GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT				
	CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA				
51	GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG				
	CCGTTGTACCGAAACAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC				
101	AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA				
	TCCTATTGTTGTATAAGGGGTTTGTATGGGTAAATATTTGAAATGGTGT				
151	GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTTCGCGG				
	CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC				
201	TCGTTTAACAACTGGAGCTGATGTGAGACATGATATAACCAGTGTTGCCAA				
	AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT				
251	ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA				
	TGTCTCAACCAAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT				
301	AATCATGCAGAGCTTTCTGTACATTAGCGCTGGATGTCACCAATGCATA				
	TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT				
351	TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA				
	ACACCAGCCGATGGCACGACCTTTATCGCGTATAAAGAAAGTAGGACTGT				
401	ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAAT				
	TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTTA				
451	CGATATACATTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACTTGC				
	GCTATATGTAAGCGGAAACCACCATTAATACTATCTGAACTTGTTGAACG				
501	TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG				
	ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC				
551	CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC				
	GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA				
601	CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTCAGAAGCAGCAAG				
	GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCTGTT				
651	ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA				
	TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT				

SUBSTITUTE SHEET (RULE 26)

178/254

FIGURE 38C (P2)

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAATGTGAACTCTTATCAACCCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTTCTTTGGGTCCTCAAGGCATTTGGGGACAGTTTAATGC
AGCAGTGTCAAAGAAACGCAGGAGTTCCGTAAACCCCTGTCAAATTACG

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTGCAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACCTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTACGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAATCGTCGCTGTAGTCCCTTGTCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACCAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTACATACCTATCTCCTGACATCGTCACTTT

179/254

FIGURE 38C (P3)

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT

1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAAGTGAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

Total number of bases is: 1855.

Sequence name: PAP278


180/254

FIGURE 38D

Figure 38. d) Amino acid sequence Comparison of Mutant Preproricin Linker region of MMP-1 (Interstitial collagenase) to Wild Type

Wild type ricin linker:	A chain- S L L I R P V V P N F N -B chain
pAP-278 (MMP-1) linker:	A chain- S L G P Q G I W G Q F N -B chain

181/254

FIGURE 39A**PCR Mutagenesis of Preproricin Gene to Create A Variant Gene in Baculovirus Transfer Vector, pVL 1393****a) Cloning Strategy**

182/254

FIGURE 39B**Sequence of Urokinase-Type Plasminogen Activator Linker Region****WT preprocin linker**

primer 280-3'

5' - GTTGTCTGGTGGCTCTGTAGCTGATGTTTGT -3'

* * * * *

```

-----TCTTTGCTTATAAGGCCA|GTGGTACCAAATTTTAAT-----
-----AGAAACGAATATTCCGGT|CACCATGGTTTAAAATTA-----
*****

```

3' - AGCAGTGTCAAATTTTTTAGGGGACCTTCT -5'

primer 280-5'

1) PCR mutagenesis

2) Ligate with pVL1393

pAP 280 linker**(uPA variant)**

```

-----AAAAAATCCCCTGGAAGA|GTTGTCTGGTGGCTCTGTA-----
-----TTTTTTAGGGGACCTTCT|CAACAGCCACCGAGACAT-----

```

183/254

FIGURE 39C (P1)

Sequence of pAP280 insert

```

      10      20      30      40      50
      |      |      |      |      |
1  GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
   CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA

51  GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
   CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC

101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
   TCCTATTGTTGTATAAGGGGTTTGTATGGGTTAATATTTGAAATGGTGT

151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTTCGCGG
   CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC

201 TCGTTTAACAACCTGGAGCTGATGTGAGACATGATATAACCAGTGTTGCCAA
   AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT

251 ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA
   TGTCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT

301 AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
   TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT

351 TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA
   ACACCAGCCGATGGCACGACCTTTATCGCGTATAAAGAAAGTAGGACTGT

401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT
   TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTTA

451 CGATATACATTTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACTTGC
   GCTATATGTAAGCGGAAACCACCATTAATACTATCTGAACTTGTTGAACG

501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
   ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC

551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
   TATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA

601 CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTCAGAAGCAGCAAG
   GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCTGTTT

651 ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA
   TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT

```

SUBSTITUTE SHEET (RULE 26)

184/254

FIGURE 39C (P2)

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAAATGTGAACTCTTATCAACCCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTAAAAAATCCCCTGGAAGAGTTGTCGGTGGCTCTGTAGC
AGCAGTGTCAAATTTTTTAGGGGACCTTCTCAACAGCCACCGAGACATCG

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACACTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTCAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCAACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTTACAACCATTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT

185/254

FIGURE 39C (P3)

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT

1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCACTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

Total number of bases is: 1855.

Sequence name: PAP280


186/254

FIGURE 39D

Figure 39. d) Amino acid sequence Comparison of Mutant Preproricin Linker region of Urokinase-Type Plasminogen Activator to Wild Type

Wild type ricin linker:	A chain- S L L I R P V V P N F N -B chain
pAP-280 (uPA) linker:	A chain- K K S P G R V V G G S V-B chain

187/254

FIGURE 40A**PCR Mutagenesis of Preproricin Gene to Create A Variant Gene in Baculovirus Transfer Vector, pVL 1393****a) Cloning Strategy**

SUBSTITUTE SHEET (RULE 26)

188/254

FIGURE 40B**Sequence of MT-MMP Linker Region****WT preprocin linker**

primer 282-3'

5' - GCTCCTGGTATTCTTGGCGCTGATGTTTGT -3'

***** * * * *

-----TCTTTGCTTATAAGGCCA|GTGGTACCAAATTTTAAT-----
 -----AGAAACGAATATTCCGGT|CACCATGGTTTAAAATTA-----
 * ***** * *

3' - AGCAGTGTCAAAGGGGTTCTGAGGATCCC -5'

primer 282-5'

1) PCR mutagenesis

2) Ligate with pVL1393

pAP 282 linker**(MT-MMP variant)**

-----CCCCAAGGACTCCTAGGG|GCTCCTGGTATTCTTGGC-----
 -----GGGGTTCCTGAGGATCCC|CGAGGACCATAAGAACCG-----

189/254

FIGURE 40C (P1)

Sequence of pAP282 insert

	10	20	30	40	50
1	GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT				
	CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA				
51	GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG				
	CCGTTGTACCGAAACAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC				
101	AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA				
	TCCTATTGTTGTATAAGGGGTTTGTATGGGTAAATATTTGAAATGGTGT				
151	GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTGCGGG				
	CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC				
201	TCGTTTAACTGGAGCTGATGTGAGACATGATATAACCAGTGTGCCAA				
	AGCAAATTGTTGACCTCGACTACACTCTGTAATATATGGTCACAACGGTT				
251	ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA				
	TGTCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT				
301	AATCATGCAGAGCTTTCTGTACATTAGCGCTGGATGTCACCAATGCATA				
	TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT				
351	TGTGGTTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA				
	ACACCAGCCGATGGCAGCAGCTTTATCGCGTATAAAGAAAGTAGGACTGT				
401	ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT				
	TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTGA				
451	CGATATACATTTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACCTTGC				
	GCTATATGTAAGCGGAAACCACCATTAATACTATCTGAACTTGTTGAACG				
501	TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG				
	ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC				
551	CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC				
	GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA				
601	CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTCAGAAGCAGCAAG				
	GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT				
651	ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA				
	TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT				

SUBSTITUTE SHEET (RULE 26)

190/254

FIGURE 40C (P2)

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAAATGTGAACTCTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTCCCCAAGGACTCCTAGGGGCTCCTGGTATTCTTGCGCG
AGCAGTGTCAAAGGGGTTCTGAGGATCCCCGAGGACCATAAGAACCGCG

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACACTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTGAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTTACAACCATTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTACACCTATCTCCTGACATCGTCACTTT

SUBSTITUTE SHEET (RULE 26)

191/254

FIGURE 40C (P3)

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT

1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAAGTAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

Total number of bases is: 1855.

Sequence name: PAP282


192/254

FIGURE 40D

**Amino acid sequence Comparison of Mutant Preproricin Linker
region of MT-MMP to Wild Type**

Wild type ricin linker:	A chain- S L L I R P V V P N F N -B chain
pAP-282 (MT-MMP) linker:	A chain- P Q G L L G A P G I L G-B chain

193/254

FIGURE 41A**PCR Mutagenesis of Preproricin Gene to Create A Variant Gene in Baculovirus Transfer Vector, pVL 1393****a) Cloning Strategy**

SUBSTITUTE SHEET (RULE 26)

FIGURE 41B**Sequence of MMP-11 (Stromelysin-3) Linker Region****WT preprocin linker**

primer 284-3'
 5'- ATGGAAGAGAGCCCATGCTCGTTTAGTTTCATGTGGAAGAGCCTCACACTGCTGATGTTTGTATGGAT-3'
 -----TCTTTGCTTATAAGGCCA|GTGGTACCAAAATTTTAAT-----
 -----AGAAACGAATAATCCGGT|CACCATGGTTTAAAAATTA-----
 3'-GGTGGTAGCAGTGTCAAAGTGCCGGGGCTCCCAAATTTCTCACCCTAAAATACTTAGACTGCAG -5'
 primer 284-5'

194/254

1) PCR mutagenesis

2) Ligate with pVL1393

pAP 284 linker

(MMP-11 variant)

---CACGGCCCCGAGGGTTTAAGAGTGGGATTTTATGAATCTGACGTC|ATGGAAGAGAGGCCATGCTCGTTTAGTTTCATGTGGAAGAGCCTCACACT---
 ---GTGCCGGGGCTCCCAAATTTCTCACCCTAAAATACTTAGACTGCAG|TACCCCTTCTCCGGTACGAGCAAAATCAAGTACAGCAAACTCGGAGTGTGA---

195/254

FIGURE 41C (P1)

Sequence of pAP284 insert

	10	20	30	40	50
1	GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT				
	CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA				
51	GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG				
	CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTATC				
101	AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA				
	TCCTATTGTTGTATAAGGGGTTTGTATGGGTAAATATTTGAAATGGTGT				
151	GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTTCGCGG				
	CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC				
201	TCGTTTAACAACCTGGAGCTGATGTGAGACATGATATAACAGTGTTGCCAA				
	AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT				
251	ACAGAGTTGGTTTGCCTATAAAACCAACGGTTTATTTTAGTTGAACTCTCA				
	TGTCTCAACCAAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT				
301	AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA				
	TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTACGTAT				
351	TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA				
	ACACCAGCCGATGGCACGACCTTTATCGCGTATAAAGAAAGTAGGACTGT				
401	ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT				
	TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGA CTACAAGTTT				
451	CGATATACATTTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACCTTGC				
	GCTATATGTAAGCGGAAACCACCATTAATACTATCTGAACTTGTTGAACG				
501	TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG				
	ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC				
551	CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC				
	GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA				
601	CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTCAGAAGCAGCAAG				
	GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTT				
651	ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA				
	TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT				

SUBSTITUTE SHEET (RULE 26)

196/254
FIGURE 41C (P2)

701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
CTAGACGTGGTCTAGGATCGCATTAAATGTGAACTCTTATCAACCCCCTCT
751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA
801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGTTTAAAGTCACACATGCTACACTCAT
851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT
901 TCGTCACAGTTT
AGCAGTGTCAA

Linker Sequence:

CACGGCCCCGAGGGTTTAAAGAGTGGGATTTTATGAATCTGACGTCATGGG
GTGCCGGGGCTCCCAAATTCTCACCCATAAATACTTAGACTGCAGTACCC

AAGAGGCCATGCTCGTTTAGTTCATGTCTGAAGAGCCTCACACT
TTCTCCGGTACGAGCAAATCAAGTACAGCAACTCGGAGTGTGA

949 GC
CG
951 TGATGTTTGATGGATCCTGAGCCCATAGTGCGTATCGTAGGTGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC
1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACAACTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT
1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTACGATTATGTCTACGTTTAGTCGAGACCTGAAA
1101 GAAAAGAGACAATACTATTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC
1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT
1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCATTACCTTGGTAGTATTTAGG
1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAATCGTCGCTGTAGTCCCTTGTCAACCATGGTGTG

SUBSTITUTE SHEET (RULE 26)

197/254

FIGURE 41C (P3)

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACCAACCATTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT
GTTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT

1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC

198/254

FIGURE 41D


**Amino acid sequence Comparison of Mutant Preproricin Linker
region of MMP-11 (Stromelysin-3) to Wild Type**

Wild type ricin linker: A chain- S L L I R P V V P N F N -B chain

pAP-284 (MMP-11) linker:

A chain- H G P E G L R V G F Y E S D V M G R G H A R L V H V E E P H T -B chain

199/254

FIGURE 42A**PCR Mutagenesis of Prepronricin Gene to Create A Variant Gene in Baculovirus Transfer Vector, pVL 1393****a) Cloning Strategy**

SUBSTITUTE SHEET (RULE 26)

200/254

FIGURE 42B**Sequence of MMP-13 = Collagenase-3 Linker Region****WT preprocin linker**

primer 286-3'

5' - GGTCAACGAGGCATTGTCGCTGATGTTTGT -3'
*****-----TCTTTGCTTATAAGGCCA | GTGGTACCAAATTTTAAT-----
-----AGAAACGAATATTCCGGT | CACCATGGTTTAAAATTA-----

3' -AGCAGTGTCAAACCTGGAGTCCCCGAACGA -5'

primer 286-5'

1) PCR mutagenesis

2) Ligate with pVL1393

**pAP 286 linker
(MMP-13 variant)**-----GGACCTCAGGGGCTTGCT | GGTCAACGAGGCATTGTC-----
-----CCTGGAGTCCCCGAACGA | CCAGTTGCTCCGTAACAG-----

201/254

FIGURE 42C (P1)

Sequence of pAP286 insert

	10	20	30	40	50
1	GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT				
	CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACTACATACGTCA				
51	GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG				
	CCGTTGTACCGAAACAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC				
101	AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA				
	TCCTATTGTTGTATAAGGGGTTTGTATGGGTAAATATTTGAAATGGTGT				
151	GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTGCGGG				
	CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC				
201	TCGTTTAACAACCTGGAGCTGATGTGAGACATGATATAACCAGTGTTGCCAA				
	AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT				
251	ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA				
	TGTCTCAACCAAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT				
301	AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA				
	TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT				
351	TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA				
	ACACCAGCCGATGGCAGGACCTTTATCGCGTATAAAGAAAGTAGGACTGT				
401	ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT				
	TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTTA				
451	CGATATACATTGCGCTTTGGTGGTAATTATGATAGACTTGAACAACCTTGC				
	GCTATATGTAAGCGGAACCAACCATTAATACTATCTGAACTTGTTGAACG				
501	TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG				
	ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC				
551	CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC				
	GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA				
601	CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTCAGAAGCAGCAAG				
	GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTT				
651	ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA				
	TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT				
701	GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA				

SUBSTITUTE SHEET (RULE 26)

202/254

FIGURE 42C (P2)

CTAGACGTGGTCTAGGATCGCATTAAATGTGAACTCTTATCAACCCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTGGACCTCAGGGGCTTGCTGGTCAACGAGGCATTGTTCG
AGCAGTGTCAAACCTGGAGTCCCCGAACGACCAGTTGCTCCGTAACAGCG

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTGCAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACACTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTCAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTACAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAATCGTCGCTGTAGTCCCTTGTCCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACAAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAACAGT

203/254

FIGURE 42C (P3)

GTTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA

1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA

1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA

1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT

1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA

1751 CTCTTGCAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT

1801 GGACATTGTAAATTTTGTAACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG

1851 TGCAG
ACGTC


204/254

FIGURE 42D

**Amino acid sequence Comparison of Mutant Preproricin Linker
region of MMP-13 (Collagenase-3) to Wild Type**

Wild type ricin linker:	A chain- S L L I R P V V P N F N -B chain
pAP-286 (MMP-13) linker:	A chain- G P Q G L A G Q R G I V -B chain

205/254

FIGURE 43A**PCR Mutagenesis of Preproricin Gene to Create A Variant Gene in Baculovirus Transfer Vector, pVL 1393****a) Cloning Strategy**

SUBSTITUTE SHEET (RULE 26)

206/254

FIGURE 43B**Sequence of Tissue-type Plasminogen Activator (tPA) Linker Region****WT preprocin linker**

primer 288-3'

5' - GGTCGTAAAGCTCTTGAAGCTGATGTTTGT -3'

***** ** *

-----TCTTTGCTTATAAGGCCA|GTGGTACCAAATTTTAAT-----
 -----AGAAACGAATATTCCGGT|CACCATGGTTTAAAATTA-----

3' -AGCAGTGTCAAACCGCCTAGACCCGTTTCC -5'

primer 288-5'

1) PCR mutagenesis

2) Ligate with pVL1393

**pAP 288 linker
(tPA variant)**

-----GGCGGATCTGGGCAAAGG|GGTCGTAAAGCTCTTGAA-----
 -----CCGCCTAGACCCGTTTCC|CCAGCATTTCGAGAACTT-----

207/254

FIGURE 43C (P1)

Sequence of pAP288 insert

	10	20	30	40	50
1	GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT				
	CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA				
51	GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG				
	CCGTTGTACCGAAACAAACCTAGGTGGAGTCCCACCAGAAAGTGTAAATC				
101	AGGATAACAACATATTCCTCCAAACAATACCCAATTATAAACTTTACCACA				
	TCCTATTGTTGTATAAGGGGTTTGTATGGGTAAATATTTGAAATGGTGT				
151	GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTTCGCGG				
	CGCCACGGTGACACGTTTCGATGTGTTTGAATAGTCTCGACAAGCGCC				
201	TCGTTTAACAACCTGGAGCTGATGTGAGACATGATATACCAGTGTTCGCAA				
	AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT				
251	ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA				
	TGTCTCAACCAAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT				
301	AATCATGCAGAGCTTCTGTACATTAGCGCTGGATGTCACCAATGCATA				
	TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT				
351	TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA				
	ACACCAGCCGATGGCACGACCTTTATCGCGTATAAAGAAAGTAGGACTGT				
401	ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT				
	TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTTA				
451	CGATATACATTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACCTTGC				
	GCTATATGTAAGCGGAAACCACCATTAATACTATCTGAAGTTGTTGAACG				
501	TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG				
	ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC				
551	CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC				
	GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA				
601	CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTTCAAGAGCAGCAAG				
	GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT				
651	ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA				
	TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT				
701	GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA				

208/254

FIGURE 43C (P2)

CTAGACGTGGTCTAGGATCGCATTAAATGTGAACTCTTATCAACCCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTGGCGGATCTGGGCAAAGGGGTCTGTAAGCTCTTGAAGC
AGCAGTGTCAAACCGCCTAGACCCGTTTCCCCAGCATTTTCGAGAACTTCG

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTCGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACCTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTACGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAATCGTCGCTGTAGTCCCTTGTCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTTACAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTACATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT

SUBSTITUTE SHEET (RULE 26)

209/254

FIGURE 43C (P3)

GTTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA
1551 TGTAAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA
1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA
1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT
1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAAACTATCTGTCTAATGA
1751 CTCTTGCAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT
1801 GGACATTGTAAATTTTGTAAGTAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG
1851 TGCAG
ACGTC

Total number of bases is: 1855.

Sequence name: PAP288


210/254

FIGURE 43D

**Amino acid sequence Comparison of Mutant Preproricin Linker
region of Tissue-type Plasminogen Activator (tPA) to Wild Type**

Wild type ricin linker:	A chain- S L L I R P V V P N F N -B chain
pAP-288 (tPA) linker:	A chain- G G S G Q R G R K A L E-B chain

211/254

FIGURE 44A**PCR Mutagenesis of Preproricin Gene to Create A Variant Gene in Baculovirus Transfer Vector, pVL 1393****a) Cloning Strategy**

SUBSTITUTE SHEET (RULE 26)

212/254

FIGURE 44B**Sequence of human Prostate-Specific Antigen (PSA) Linker Region****WT preprocin linker**

primer 290-3'
 5' - TCTTCCGATATTTTAAATGCTGATGTTTGT - 3'
 ***** *

-----TCTTTGCTTATAAGGCCA|GTGGTACCAAATTTTAAT-----
 -----AGAAACGAATATTCCGGT|CACCATGGTTTAAAATTA-----
 ***** *

3' -AGCAGTGTCAAAGAAACAGTCGAGAAGAG -5'
 primer 290-5'

1) PCR mutagenesis

2) Ligate with pVL1393

**pAP 290 linker
(PSA variant)**

-----TCTTTGTCAGCTCTTCTC|TCTTCCGATATTTTAAAT-----
 -----AGAAACAGTCGAGAAGAG|AGAAGGCTATAAAAATTA-----

213/254

FIGURE 44C (P1)

Sequence of pAP290 insert

	10	20	30	40	50
1	GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT				
	CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA				
51	GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTTCACATTAG				
	CCGTTGTACCGAAACAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC				
101	AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA				
	TCCTATTGTTGTATAAGGGGTTTGTATGGGTTAATATTTGAAATGGTGT				
151	GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTTCGCGG				
	CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC				
201	TCGTTTAACAACCTGGAGCTGATGTGAGACATGATATACCAGTGTTGCCAA				
	AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT				
251	ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA				
	TGTCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT				
301	AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA				
	TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT				
351	TGTGGTTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA				
	ACACCAGCCGATGGCAGGACCTTTATCGCGTATAAAGAAAGTAGGACTGT				
401	ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT				
	TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTAA				
451	CGATATACATTTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACTTGC				
	GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACTTGTTGAACG				
501	TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG				
	ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC				
551	CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC				
	GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA				
601	CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTCAGAAGCAGCAAG				
	GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTTT				
651	ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA				
	TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT				
701	GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA				

214/254

FIGURE 44C (P2)

CTAGACGTGGTCTAGGATCGCATTAAATGTGAACTCTTATCAACCCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTTCTTTGTCAGCTCTTCTCTCTTCCGATATTTTAAATGC
AGCAGTGTCAAAAGAAACAGTCGAGAAGAGAGAAGGCTATAAAAATTACG

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTGCAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACCTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTCAAGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCAACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTGTACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT

215/254

FIGURE 44C (P3)

GTTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA
1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA
1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA
1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT
1701 TGGTGACCCAAACCAAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA
1751 CTCTTGCAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT
1801 GGACATTGTAAATTTTGTAAGTAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG
1851 TGCAG
ACGTC

Total number of bases is: 1855.

Sequence name: PAP290


216/254

FIGURE 44D

**Amino acid sequence Comparison of Mutant Preproricin Linker
region of human Prostate-Specific Antigen (PSA) to Wild Type**

Wild type ricin linker:	A chain- S L L I R P V V P N F N -B chain
pAP-290 (PSA) linker:	A chain- S L S A L L S S D I F N -B chain

217/254

FIGURE 45A**PCR Mutagenesis of Preproricin Gene to Create A Variant Gene in Baculovirus Transfer Vector, pVL 1393****a) Cloning Strategy**

SUBSTITUTE SHEET (RULE 26)

218/254

FIGURE 45B

Sequence of Kallikrein (hK3) Linker Region

WT preprocin linker

```

      primer 292-3'
      5' - ATTATCGGTGGCTTTAATGCTGATGTTTGT -3'
            * * * * *
-----TCTTTGCTTATAAGGCCA|GTGGTACCAAATTTTAAT-----
-----AGAAACGAATATTCCGGT|CACCATGGTTTAAATTA-----
            * * * * *
      3' - AGCAGTGTCAAAAGAAACGGATCTAAATTT -5'
      primer 292-5'

```

1) PCR mutagenesis

2) Ligate with pVL1393

```

      pAP 292 linker
      (Kallikrein variant)
-----TCTTTGCCTAGATTTAAA|ATTATCGGTGGCTTTAAT-----
-----AGAAACGGATCTAAATTT|TAATAGCCACCGAAATTA-----

```


219/254

FIGURE 45C (P1)

Sequence of PAP292 insert

	10	20	30	40	50
1	GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT				
	CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA				
51	GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG				
	CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTAAATC				
101	AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA				
	TCCTATTGTTGTATAAGGGGTTTGTATGGGTTAATATTTGAAATGGTGT				
151	GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTTCGCGG				
	CGCCACGGTGACACGTTTCGATGTGTTGAAATAGTCTCGACAAGCGCC				
201	TCGTTTAACTGAGCTGATGTGAGACATGATATACCAGTGTGCCAA				
	AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT				
251	ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA				
	TGTCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT				
301	AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA				
	TTAGTACGTCCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT				
351	TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA				
	ACACCAGCCGATGGCAGGACCTTTATCGCGTATAAAGAAAGTAGGACTGT				
401	ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT				
	TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTTA				
451	CGATATACATTTCGCTTTGGTGGTAATTATGATAGACTTGAACAACTTGC				
	GCTATATGTAAGCGGAAACCACCATTAATACTATCTGAACTTGTTGAACG				
501	TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG				
	ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC				
551	CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC				
	GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA				
601	CTGGCTCGTTCCTTTATAATTTGCATCCAAATGATTTCAGAAGCAGCAAG				
	GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTT				
651	ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA				
	TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT				
701	GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA				

SUBSTITUTE SHEET (RULE 26)

220/254

FIGURE 45C (P2)

CTAGACGTGGTCTAGGATCGCATTAAATGTGAACTCTTATCAACCCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTTCTTTGCCTAGATTTAAAATTATCGGTGGCTTTAATGC
AGCAGTGTCAAAAGAAACGGATCTAAATTTAATAGCCACCGAAATTACG

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTGAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACACTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTCAAGTTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAAATCGTCGCTGTAGTCCCTTGTCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACAAACCATTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT

221/254

FIGURE 45C (P3)

GTTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA
1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA
1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA
1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT
1701 TGGTGACCCAAACCAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAAACTATCTGTCTAATGA
1751 CTCTTGCAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT
1801 GGACATTGTAAATTTTGTAAGTGAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG
1851 TGCAG
ACGTC

Total number of bases is: 1855.

Sequence name: PAP292


222/254

FIGURE 45D

Amino acid sequence Comparison of Mutant Preproricin Linker
region of Kallikrein (hK3) to Wild Type

Wild type ricin linker:	A chain- S L L I R P V V P N F N -B chain
pAP-292 (hK3) linker:	A chain- S L P R F K I I G G F N -B chain

223/254

FIGURE 46A**PCR Mutagenesis of Preproricin Gene to Create A Variant Gene in Baculovirus Transfer Vector, pVL 1393****a) Cloning Strategy**

SUBSTITUTE SHEET (RULE 26)

224/254

FIGURE 46B

Sequence of Neutrophil Elastase Linker Region

WT preprocin linker

```

          primer 294-3'
          5' - GTTCCTGGTAATTTTAATGCTGATGTTTGT -3'
                ** *****
-----TCTTTGCTTATAAGGCCA|GTGGTACCAATTTTAAT-----
-----AGAAACGAATATTCCGGT|CACCATGCTTTAAAATTA-----
                *** ***
3' -AGCAGTGTCAAAGAAACGAACCGTAACGA -5'
          primer 294-5'

```

1) PCR mutagenesis

2) Ligate with pVL1393

```

          pAP 294 linker
          (Neutrophil elastase variant)
-----TCTTTGCTTGGCATTGCT|GTTCCTGGTAATTTTAAT-----
-----AGAAACGAACCGTAACGA|CAAGGACCATTAAAATTA-----

```

225/254

FIGURE 46C (P1)

Sequence of pAP294 insert

	10	20	30	40	50
1	GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT				
	CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA				
51	GGCAACATGGCCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG				
	CCGTTGTACCGAAACAAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC				
101	AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA				
	TCCTATTGTTGTATAAGGGGTTTGTATGGGTTAATATTTGAAATGGTGT				
151	GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTGCGGG				
	CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC				
201	TCGTTTAAACAACCTGGAGCTGATGTGAGACATGATATACCAGTGTTGCCAA				
	AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT				
251	ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA				
	TGTCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT				
301	AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA				
	TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT				
351	TGTGGTGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA				
	ACACCAGCCGATGGCAGGACCTTTATCGCGTATAAAGAAAGTAGGACTGT				
401	ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT				
	TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTTA				
451	CGATATACATTGCGCTTTGGTGGTAATTATGATAGACTTGAACAACCTTGC				
	GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACTTGTGTAACG				
501	TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG				
	ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC				
551	CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC				
	GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA				
601	CTGGCTCGTTCCTTTATAAATTGTCATCCAAATGATTTCAGAAGCAGCAAG				
	GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTT				
651	ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA				
	TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT				
701	GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA				

226/254

FIGURE 46C (P2)

CTAGACGTGGTCTAGGATCGCATTAAATGTGAACCTTTATCAACCCCTCT

751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA

801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTTAAGTCACACATGCTACACTCAT

851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT

901 TCGTCACAGTTTTCTTTGCTTGGCATTGCTGTTCCCTGGTAATTTAATGC
AGCAGTGTCAAAGAAACGAACCGTAACGACAAGGACCATTAAAATTACG

951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTGCAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC

1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACCTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT

1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTACGATTATGTCTACGTTTAGTCGAGACCTGAAA

1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC

1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTCAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT

1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCTATTACCTTGGTAGTATTTAGG

1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAATCGTCGCTGTAGTCCCTTGTCCACCATGGTGTG

1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA

1351 AATAATACACAACCTTTTGTACAAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAAACAATGTTGGTAACAACCCGATATACCAGACAC

1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTACATACCTATCTCCTGACATCGTCACTTT

1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC

1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT

227/254

FIGURE 46C (P3)

GT TTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA
1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA
1601 TCAAGAATGATGGAACCATT TTTAAATTTGTATAGTGGATTGGTGT TAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA
1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT
1701 TGGTGACCCAAACCAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA
1751 CTCTTGCA GTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGT CACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT
1801 GGACATTGTAAATTTTGTA ACTGAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG
1851 TGCAG
ACGTC

Total number of bases is: 1855.

Sequence name: PAP294

228/254


FIGURE 46D

Amino acid sequence Comparison of Mutant Preproricin Linker
region of Neutrophil elastase to Wild Type

Wild type ricin linker: A chain- S L L I R P V V P N F N -B chain

pAP-294 (Neutrophil elastase) linker:
 A chain- S L L G I A V P G N F N -B chain

229/254

FIGURE 47A**PCR Mutagenesis of Preproricin Gene to Create A Variant Gene in Baculovirus Transfer Vector, pVL 1393****a) Cloning Strategy**

SUBSTITUTE SHEET (RULE 26)

230/254

FIGURE 47B**Sequence of Calpain Linker Region****WT preprocin linker**

primer 296-3'

5'- ACTCCTAGAACCCCCCAGCTGATGTTTGT -3'

-----TCTTTGCTTATAAGGCCA|GTGGTACCAAATTTTAAT-----
 -----AGAAACGAATATTCCGGT|CACCATGGTTTAAATTA-----
 * *** ** *****

3'-AGCAGTGTCAAAAAAAGTTTTTATAACAA -5'

primer 296-5'

1) PCR mutagenesis

2) Ligate with pVL1393

pAP 296 linker

(Calpain variant)

-----TTTTTCAAAAATATTGTT|ACTCCTAGAACCCCCCA-----
 -----AAAAAGTTTTTATAACAA|TGAGGATCTTGGGGGGGT-----

231/254

FIGURE 47C (P1)

Sequence of pAP296 insert

```
      10      20      30      40      50
      |      |      |      |      |
1  GAATTCATGAAACCGGGAGGAAATACTATTGTAATATGGATGTATGCAGT
   CTTAAGTACTTTGGCCCTCCTTTATGATAACATTATACCTACATACGTCA
51 GGCAACATGGCTTTGTTTTGGATCCACCTCAGGGTGGTCTTTCACATTAG
   CCGTTGTACCGAAACAAACCTAGGTGGAGTCCCACCAGAAAGTGTAATC
101 AGGATAACAACATATTCCCCAAACAATACCCAATTATAAACTTTACCACA
   TCCTATTGTTGTATAAGGGGTTTGTTATGGGTTAATATTTGAAATGGTGT
151 GCGGGTGCCACTGTGCAAAGCTACACAACTTTATCAGAGCTGTTTCGCGG
   CGCCACGGTGACACGTTTCGATGTGTTTGAAATAGTCTCGACAAGCGCC
201 TCGTTTAACAACTGGAGCTGATGTGAGACATGATATACCAGTGTTGCCAA
   AGCAAATTGTTGACCTCGACTACACTCTGTACTATATGGTCACAACGGTT
251 ACAGAGTTGGTTTGCCTATAAACCAACGGTTTATTTTAGTTGAACTCTCA
   TGTCTCAACCAACGGATATTTGGTTGCCAAATAAAATCAACTTGAGAGT
301 AATCATGCAGAGCTTTCTGTTACATTAGCGCTGGATGTCACCAATGCATA
   TTAGTACGTCTCGAAAGACAATGTAATCGCGACCTACAGTGGTTACGTAT
351 TGTGGTCGGCTACCGTGCTGGAAATAGCGCATATTTCTTTCATCCTGACA
   ACACCAGCCGATGGCAGCACCTTTATCGCGTATAAAGAAAGTAGGACTGT
401 ATCAGGAAGATGCAGAAGCAATCACTCATCTTTTCACTGATGTTCAAAT
   TAGTCCTTCTACGTCTTCGTTAGTGAGTAGAAAAGTGACTACAAGTTTTA
451 CGATATACATTTCGCCTTTGGTGGTAATTATGATAGACTTGAACAACTTGC
   GCTATATGTAAGCGGAAACCACCATTAACTATCTGAACTTGTTGAACG
501 TGGTAATCTGAGAGAAAATATCGAGTTGGGAAATGGTCCACTAGAGGAGG
   ACCATTAGACTCTCTTTTATAGCTCAACCCTTTACCAGGTGATCTCCTCC
551 CTATCTCAGCGCTTTATTATTACAGTACTGGTGGCACTCAGCTTCCAAC
   GATAGAGTCGCGAAATAATAATGTCATGACCACCGTGAGTCGAAGGTTGA
601 CTGGCTCGTTCCTTTATAATTGTCATCCAAATGATTTCAGAAGCAGCAAG
   GACCGAGCAAGGAAATATTAAACGTAGGTTTACTAAAGTCTTCGTCGTT
651 ATTCCAATATATTGAGGGAGAAATGCGCACGAGAATTAGGTACAACCGGA
   TAAGGTTATATAACTCCCTCTTTACGCGTGCTCTTAATCCATGTTGGCCT
701 GATCTGCACCAGATCCTAGCGTAATTACACTTGAGAATAGTTGGGGGAGA
```

232/254

FIGURE 47C (P2)

CTAGACGTGGTCTAGGATCGCATTAAATGTGAACTCTTATCAACCCCCTCT
751 CTTTCCACTGCAATTCAAGAGTCTAACCAAGGAGCCTTTGCTAGTCCAAT
GAAAGGTGACGTTAAGTTCTCAGATTGGTTCCTCGGAAACGATCAGGTTA
801 TCAACTGCAAAGACGTAATGGTTCCAAATTCAGTGTGTACGATGTGAGTA
AGTTGACGTTTCTGCATTACCAAGGTTAAGTCACACATGCTACACTCAT
851 TATTAATCCCTATCATAGCTCTCATGGTGTATAGATGCGCACCTCCACCA
ATAATTAGGGATAGTATCGAGAGTACCACATATCTACGCGTGGAGGTGGT
901 TCGTCACAGTTTTTTTTTCAAAAATATTGTTACTCCTAGAACCCCCCAGC
AGCAGTGTCAAAAAAAGTTTTTATAACAATGAGGATCTTGGGGGGGTCG
951 TGATGTTTGTATGGATCCTGAGCCCATAGTGCGTATCGTAGGTGCAAATG
ACTACAAACATACCTAGGACTCGGGTATCACGCATAGCATCCAGCTTTAC
1001 GTCTATGTGTTGATGTTAGGGATGGAAGATTCCACAACGGAAACGCAATA
CAGATACACAACCTACAATCCCTACCTTCTAAGGTGTTGCCTTTGCGTTAT
1051 CAGTTGTGGCCATGCAAGTCTAATACAGATGCAAATCAGCTCTGGACTTT
GTCAACACCGGTACGTTTACGATTATGTCTACGTTTAGTCGAGACCTGAAA
1101 GAAAAGAGACAATACTATTTCGATCTAATGGAAAGTGTTTAACTACTTACG
CTTTTCTCTGTTATGATAAGCTAGATTACCTTTCACAAATTGATGAATGC
1151 GGTACAGTCCGGGAGTCTATGTGATGATCTATGATTGCAATACTGCTGCA
CCATGTACAGGCCCTCAGATACACTACTAGATACTAACGTTATGACGACGT
1201 ACTGATGCCACCCGCTGGCAAATATGGGATAATGGAACCATCATAAATCC
TGACTACGGTGGGCGACCGTTTATACCTATTACCTTGGTAGTATTTAGG
1251 CAGATCTAGTCTAGTTTTAGCAGCGACATCAGGGAACAGTGGTACCACAC
GTCTAGATCAGATCAAATCGTCGCTGTAGTCCCTTGTCACCATGGTGTG
1301 TTACAGTGCAAACCAACATTTATGCCGTTAGTCAAGGTTGGCTTCCTACT
AATGTCACGTTTGGTTGTAAATACGGCAATCAGTTCCAACCGAAGGATGA
1351 AATAATACACAACCTTTTGTACAAACCATTTGTTGGGCTATATGGTCTGTG
TTATTATGTGTTGGAAACAATGTTGGTAACAACCCGATATACCAGACAC
1401 CTTGCAAGCAAATAGTGGACAAGTATGGATAGAGGACTGTAGCAGTGAAA
GAACGTTTCGTTTATCACCTGTTTCATACCTATCTCCTGACATCGTCACTTT
1451 AGGCTGAACAACAGTGGGCTCTTTATGCAGATGGTTCAATACGTCCTCAG
TCCGACTTGTTGTCACCCGAGAAATACGTCTACCAAGTTATGCAGGAGTC
1501 CAAAACCGAGATAATTGCCTTACAAGTGATTCTAATATACGGGAAACAGT

233/254

FIGURE 47C (P3)

GTTTTGGCTCTATTAACGGAATGTTCACTAAGATTATATGCCCTTTGTCA
1551 TGTTAAGATCCTCTCTTGTGGCCCTGCATCCTCTGGCCAACGATGGATGT
ACAATTCTAGGAGAGAACACCGGGACGTAGGAGACCGGTTGCTACCTACA
1601 TCAAGAATGATGGAACCATTTTAAATTTGTATAGTGGATTGGTGTAGAT
AGTTCTTACTACCTTGGTAAAATTTAAACATATCACCTAACCACAATCTA
1651 GTGAGGCGATCGGATCCGAGCCTTAAACAAATCATTCTTTACCCCTCTCCA
CACTCCGCTAGCCTAGGCTCGGAATTTGTTTAGTAAGAAATGGGAGAGGT
1701 TGGTGACCCAAACCAATATGGTTACCATTATTTTGATAGACAGATTACT
ACCACTGGGTTTGGTTTATACCAATGGTAATAAACTATCTGTCTAATGA
1751 CTCTTGCAGTGTGTGTGTCCTGCCATGAAAATAGATGGCTTAAATAAAAA
GAGAACGTCACACACACAGGACGGTACTTTTATCTACCGAATTTATTTTT
1801 GGACATTGTAAATTTTGTAAGTAAAGGACAGCAAGTTATATCGAATTCC
CCTGTAAACATTTAAACATTGACTTTCCTGTCGTTCAATATAGCTTAAGG
1851 TGCAG
ACGTC

Total number of bases is: 1855.

Sequence name: PAP296


234/254

FIGURE 47D


**Amino acid sequence Comparison of Mutant Preproricin Linker
region of Calpain to Wild Type**

Wild type ricin linker:	A chain- S L L I R P V V P N F N -B chain
pAP-296 (Calpain) linker:	A chain- F F K N I V T P R T P P -B chain

235/254


FIGURE 48**Cleavage of pAP 214 by Cathepsin B****A. Ricin standard****B. pAP 214****C. pAP 214 digested with 100 ng of Cathepsin B (18 hours)****D. pAP 214 digested with 618 ng of Cathepsin B (18 hours)**

236/254

FIGURE 49**Cleavage of pAP 220 with MMP-9**


- A.** pAP 220
- B.** pAP 220 digested with 200 ng of MMP-9 (16 hrs)
- C.** pAP 220 digested with 20 ng of MMP-9 (16hrs)
- D.** pAP 220 digested with 20 ng of MMP-9 (2hrs)

237/254

FIGURE 50**Activation of pAP 214**


- A. 41.7 pg of pAP 214 digested with Cathepsin B
- B. 291 pg of pAP 214 digested with Cathepsin B
- C. 2.0 ng of pAP 214 digested with Cathepsin B
- D. 14.2 ng of pAP 214 digested with Cathepsin B
- E. 100 ng of pAP 214 digested with Cathepsin B
- F. Negative control
- G. Ricin A chain
- H. 41.7 pg of pAP 214 variant
- I. 291 pg of pAP 214 variant
- J. 2.0 ng of pAP 214 variant
- K. 14.2 ng of pAP 214 variant
- L. 100ng of pAP 214 variant
- M. RNA ladder

238/254

FIGURE 51**Activation of pAP 220**


- A. 48.5 pg of pAP 220 variant
- B. 291 pg of pAP 220 variant
- C. 2.0 ng of pAP 220 variant
- D. 14.3 ng of pAP 220 variant
- E. 100 ng of pAP 220 variant
- F. Ricin A chain
- G. Negative Control
- H. 48.5 pg of pAP 220 variant digested with MMP-9
- I. 291 pg of pAP 220 variant digested with MMP-9
- J. 2.0 ng of pAP 220 variant digested with MMP-9
- K. 14.3 ng of pAP 220 variant digested with MMP-9
- L. 100 ng of pAP 220 variant digested with MMP-9
- M. RNA ladder

239/254

FIGURE 52**Cleavage of pAP-248 Protein by The Human Cytomegalovirus (HCMV) protease**


- A. pAP-248 (0.279 μ g)
B. pAP-248 protein (0.279 μ g) digested with 0.25 μ g of the HCMV protease
C. Ricin standard (20 ng)
D. Ricin standard (40 ng)

240/254

FIGURE 53**Activation of pAP-248 Protein**


- A. 90 ng of pAP-248 variant
- B. 12.8 ng of pAP-248 variant
- C. 1.8 ng of pAP-248 variant
- D. 260 pg pAP-248 variant
- E. 37 pg of pAP-248 variant
- F. Negative control
- G. Ricin A chain
- H. 37 pg of pAP-248 digested with HCMV protease
- I. 260 pg of pAP-248 digested with HCMV protease
- J. 1.8 ng of pAP-248 digested with HCMV protease
- K. 12.8 ng of pAP-248 digested with HCMV protease
- L. 90 ng of pAP-248 digested with HCMV protease
- M. RNA ladder

241/254

FIGURE 54**Cleavage of pAP-256 protein by The Hepatitis A Virus 3C (HAV 3C) Protease**

- A. Ricin standard (0.250 ug)
B. pAP-256 protein (0.378 ug)
C. pAP-256 protein digested (0.302 ug) with 1.25 μ g of the HAV 3C protease


242/254

FIGURE 55**Activation of pAP-256 Protein**

- A. 100 ng of pAP-256 variant
B. 14.2 ng of pAP-256 variant
C. 2.0 ng of pAP-256 variant
D. 291 pg of pAP-256 variant
E. 41.7 pg of pAP-256 variant
F. Negative control
G. Ricin A chain
H. 41.7 pg of pAP-256 digested with HAV 3C protease
I. 291 pg of pAP-256 digested with HAV 3C protease
J. 2.0 ng of pAP-256 digested with HAV 3C protease
K. 14.2 ng of pAP-256 digested with HAV 3C protease
L. 100 ng of pAP-256 digested with HAV 3C protease
M. RNA ladder

SUBSTITUTE SHEET (RULE 26)

243/254


FIGURE 56**Cytotoxicity of Digested and Undigested
pAP 214 with Cathepsin B to COS-1 Cells**

	Ricin	pAP 214	pAP 214 + Cathepsin B
IC ₅₀ (ng/ml)	0.11	1.9	0.078
Relative Toxicity	1X	17X	0.7X

244/254


FIGURE 57

Cytotoxicity of pAP220 Digested with MMP-9 Compared to Freshly Thawed pAP220 and Ricin on COS-1 Cells




	Ricin	pAP 220	pAP 220 + MMP-9
IC ₅₀ (ng/ml)	0.31	6.7	0.13
Relative Toxicity	1X	22X	0.4X

245/254

FIGURE 58**Cleavage of pAP-270 protein by The Matrix Metalloproteinase 2 (MMP-2)****A. pAP-270 (0.120 μ g) undigested****B. pAP-270 (0.120 μ g) digested with 0.250 μ g MMP-2****C. Ricin Standard (0.05 μ g)****SUBSTITUTE SHEET (RULE 26)**


246 /254

FIGURE 59**Activation of pAP-270 protein**


- A. 100 ng of digested pAP-270
- B. 14.2 ng of digested pAP-270
- C. 2.0 ng of digested pAP-270
- D. 290 pg of digested pAP-270
- E. 46 ng of digested pAP-270
- F. Ricin A chain
- G. Negative control
- H. 46 pg of pAP-270
- I. 290 pg of pAP-270
- J. 2.0 ng of pAP-270
- K. 14.2 ng of pAP-270
- L. 100 ng of pAP-270

SUBSTITUTE SHEET (RULE 26)

247/254


FIGURE 60**Cleavage of pAP-288 protein by Plasminogen Tissue Activator (t-PA)****A. Ricin Standard (0.05µg)****B. pAP-288 (0.66 µg) undigested****C. pAP-288 (0.60 µg) digested with 0.18 µg of t-PA protease**

248/254


FIGURE 61**Activation of pAP-288 protein**

- A. 200 ng of pAP-288
- B. 28.4 ng of pAP-288
- C. 4.0 ng of pAP-288
- D. 482 pg of pAP-288
- E. 83.4 pg of pAP-288
- F. Ricin A chain
- G. Negative control
- H. 83.4 pg of pAP-288 digested with tissue Plasminogen Activator (t-PA)
- I. 482 pg of pAP-288 digested with t-PA
- J. 4.0 ng of pAP-288 digested with t-PA
- K. 28.4 ng of pAP-288 digested with t-PA
- L. 200 ng of pAP-288 digested with t-PA
- M. RNA ladder

249/254

FIGURE 62**Cleavage of pAP 294 With Human Neutrophil Elastase****A. Ricin Standard (0.050 μ g)****B. pAP 294 protein (0.171 μ g) digested with 1.42 μ g of Human Neutrophil Elastase****C. pAP 294 protein (0.121 μ g)**


250/254

FIGURE 63**Activation of pAP 294 Protein**


- A. 60 ng of pAP 294
- B. 8..57 ng of pAP 294
- C. 1.22 ng of pAP 294
- D. 175 pg of pAP 294
- E. 25 pg of pAP 294
- F. Ricin A chain
- G. Negative Control
- H. 360 ng of pAP 294 digested with Human Neutrophil Elastase
- I. 51 ng of pAP 294 digested with Human Neutrophil Elastase
- J. 7.3 ng of pAP 294 digested with Human Neutrophil Elastase
- K. 1.0 ng of pAP 294 digested with Human Neutrophil Elastase
- L. 150 pg of pAP 294 digested with Human Neutrophil Elastase
- M. RNA ladder

SUBSTITUTE SHEET (RULE 26)

251/254

FIGURE 64**Cleavage of pAP 296 with Calpain****A. Ricin Standard (0.05 μ g)****B. pAP 296 (0.761 μ g) undigested****C. pAP 296 (0.761 μ g) digested with 4.0 μ g of Calpain**


252/254

FIGURE 65**Activation of pAP 296 Protein**

- A. 100 ng of pAP 296 variant
- B. 14.2 ng of pAP 296 variant
- C. 2.0 ng of pAP 296 variant
- D. 290 pg of pAP 296 variant
- E. 46 pg of pAP 296 variant
- F. Ricin A chain
- G. Negative control
- H. 46 pg of pAP 296 variant digested with Calpain
- I. 290 pg of pAP 296 variant digested with Calpain
- J. 2.0 ng of pAP 296 variant digested with Calpain
- K. 14.2 ng of pAP 296 variant digested with Calpain
- L. 100 ng of pAP 296 variant digested with Calpain
- M. RNA ladder


SUBSTITUTE SHEET (RULE 26)

253/254

FIGURE 66**Cleavage of pAP-222 Protein by The Matrix Metalloproteinase 2 (MMP-2)**

- A. Ricin Standard (0.250 ug)**
B. pAP-222 Protein (0.250 ug)
C. pAP-222 protein (0.250 ug) digested with 0.28 ug of MMP-2

254/254

FIGURE 67**Activation of pAP-222 Protein**

- A. 100 ng of pAP-222 variant
- B. 14.2 ng of pAP-222 variant
- C. 2.0 ng of pAP-222 variant
- D. 291 pg of pAP-222 variant
- E. 41.7 pg of pAP-222 variant
- F. Ricin A chain
- G. Ricin A chain
- H. 41.7 pg of pAP-222 digested with MMP-2
- I. 291 pg of pAP-222 digested with MMP-2
- J. 2.0 ng of pAP-222 digested with MMP-2
- K. 14.2 ng of pAP-222 digested with MMP-2
- L. 100 ng of pAP-222 digested with MMP-2
- M. RNA ladder




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/29, 15/62, 15/70, 15/86, A61K 38/16, 48/00		(11) International Publication Number: WO 98/49311
A3		(43) International Publication Date: 5 November 1998 (05.11.98)
(21) International Application Number: PCT/CA98/00394 (22) International Filing Date: 30 April 1998 (30.04.98) (30) Priority Data: 60/045,148 30 April 1997 (30.04.97) US 60/063,715 29 October 1997 (29.10.97) US (71) Applicant (for all designated States except US): DE NOVO ENZYME CORPORATION [CA/CA]; #2 Suite SFU Discovery Park, Burnaby, British Columbia V5A 1S6 (CA). (72) Inventor; and (75) Inventor/Applicant (for US only): BORGFORD, Thor [CA/CA]; 443 Fadar Street, New Westminster, British Columbia V3L 3T2 (CA). (74) Agent: BERESKIN & PARR; 40th floor, 40 King Street West, Toronto, Ontario M5H 3Y2 (CA).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 11 February 1999 (11.02.99)

(54) Title: RICIN-LIKE TOXIN VARIANTS FOR TREATMENT OF CANCER, VIRAL OR PARASITIC INFECTIONS

(57) Abstract

The present invention provides a protein having an A chain of a ricin-like toxin, a B chain of a ricin-like toxin and a heterologous linker amino acid sequence, linking the A and B chains. The linker sequence contains a cleavage recognition site for a disease specific protease such as a cancer, fungal, viral or parasitic protease. The invention also relates to a nucleic acid molecule encoding the protein and to expression vectors incorporating the nucleic acid molecule. Also provided is a method of inhibiting or destroying mammalian cancer cells, cells infected with a virus, a fungus, or parasite, or parasites utilizing the nucleic acid molecules and proteins of the invention and pharmaceutical compositions for treating human cancer, viral infection, fungal infection, or parasitic infection.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

Int. Application No.
PCT/CA 98/00394

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C12N15/29 C12N15/62 C12N15/70 C12N15/86 A61K38/16 A61K48/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 C07K C12N A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 94 18332 A (US HEALTH) 18 August 1994 * see the whole document, esp. p.16 l.30 - p.20 l.2 *	1-10, 12-23, 25-35
Y	--- WESTBY M ET AL: "PREPARATION AND CHARACTERIZATION OF RECOMBINANT PRORICIN CONTAINING AN ALTERNATIVE PROTEASE-SENSITIVE LINKER SEQUENCE" BIOCONJUGATE CHEMISTRY, vol. 3, no. 5, 1 January 1992, pages 375-381, XP000578216 cited in the application * see the whole document, esp. last paragraph * --- -/--	1-10, 12-23, 25-35
<div style="display: flex; justify-content: space-between;"> <input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex. </div>		
* Special categories of cited documents : <div style="display: flex;"> <div style="flex: 1;"> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="flex: 1;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <div style="text-align: center; font-weight: bold;">27 October 1998</div>		Date of mailing of the international search report <div style="text-align: center; font-weight: bold;">10/11/1998</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer <div style="text-align: center; font-weight: bold;">Kania, T</div>

INTERNATIONAL SEARCH REPORT

ational Application No

PCT/CA 98/00394

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>LEPPLA S H ET AL: "DEVELOPMENT OF ANTHRAX-TOXIN BASED FUSION PROTEINS FOR TARGETING OF HIV-1-INFECTED CELLS" ZENTRALBLATT FUER BAKTERIOLOGIE. SUPPLEMENT, vol. 24, 1994, pages 431-442, XP002041056 cited in the application * see the whole document, esp. pp.437-39 *</p>	1-35
A	<p>COOK J P ET AL: "BIOLOGICALLY ACTIVE INTERLEUKIN 2-RICIN A CHAIN FUSION PROTEINS MAY REQUIRE INTRACELLULAR PROTEOLYTIC CLEAVAGE TO EXHIBIT A CYTOTOXIC EFFECT" BIOCONJUGATE CHEMISTRY, vol. 4, no. 6, 1 November 1993, pages 440-447, XP000417282 see the whole document</p>	1-35
A	<p>O'HARE M ET AL: "CYTOTOXICITY OF A RECOMBINANT RICIN-A-CHAIN FUSION PROTEIN CONTAINING A PROTEOLYTICALLY-CLEAVABLE SPACER SEQUENCE" FEBS LETTERS, vol. 273, no. 1/02, 29 October 1990, pages 200-204, XP002041057 cited in the application see the whole document</p>	1-35
A	<p>PANCHAL R. ET AL.: "Tumor protease-activated, pore-forming toxins from a combinatorial library" NATURE BIOTECHNOLOGY, vol. 14, no. 7, 14 July 1996, pages 852-856, XP002082096 cited in the application see the whole document</p>	1-35
A	<p>EP 0 466 222 A (DOWELANCO) 15 January 1992 cited in the application see the whole document</p>	1-35
P,X	<p>WO 97 41233 A (NOVO ENZYME CORP DE ;BORG FORD THOR (CA)) 6 November 1997 see the whole document</p>	1-10, 12-23, 25-35

INTERNATIONAL SEARCH REPORT

International application No.

PCT/CA 98/00394

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 25-30, 32, 33
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims: it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 98/00394

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9418332	A	18-08-1994	US 5591631 A	07-01-1997
			US 5677274 A	14-10-1997
			AT 169959 T	15-09-1998
			AU 682500 B	09-10-1997
			AU 6392294 A	29-08-1994
			DE 69412593 D	24-09-1998
			EP 0684997 A	06-12-1995
EP 0466222	A	15-01-1992	US 5248606 A	28-09-1993
			AU 638133 B	17-06-1993
			AU 7832991 A	12-12-1991
			CA 2044201 A	12-12-1991
			CN 1062172 A	24-06-1992
			JP 4279599 A	05-10-1992
			US 5635384 A	03-06-1997
			US 5646026 A	08-07-1997
WO 9741233	A	06-11-1997	AU 2377097 A	19-11-1997